

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/308128643>

# Efficacy of inspiratory muscle training as a practical and minimally intrusive technique to aid functional fitness among adults with obesity

Article in *Respiratory Physiology & Neurobiology* · September 2016

DOI: 10.1016/j.resp.2016.09.007

CITATIONS

10

READS

318

4 authors:



**Andrew M Edwards**

Canterbury Christ Church University

70 PUBLICATIONS 1,972 CITATIONS

[SEE PROFILE](#)



**Deborah Graham**

The Cairns Institute

36 PUBLICATIONS 442 CITATIONS

[SEE PROFILE](#)



**Saul Bloxham**

University of St Mark & St John

31 PUBLICATIONS 282 CITATIONS

[SEE PROFILE](#)



**Graeme P Maguire**

Western Health

158 PUBLICATIONS 3,702 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Drug and Alcohol Service Provision in Northern Australia [View project](#)



Indigenous Health [View project](#)



# Efficacy of inspiratory muscle training as a practical and minimally intrusive technique to aid functional fitness among adults with obesity



A.M. Edwards<sup>a,b,\*</sup>, D. Graham<sup>b</sup>, S. Bloxham<sup>a</sup>, G.P. Maguire<sup>b,c</sup>

<sup>a</sup> University of St Mark & St John, Plymouth, United Kingdom, United Kingdom

<sup>b</sup> James Cook University, Cairns, Australia

<sup>c</sup> Baker IDI Heart and Diabetes Research Institute, Melbourne, Australia

## ARTICLE INFO

### Article history:

Received 30 March 2016

Received in revised form 4 August 2016

Accepted 12 September 2016

Available online 13 September 2016

### Keywords:

Obesity

Physical activity

Respiratory disorders

Chronic disease

## ABSTRACT

**Objective:** To examine the efficacy of inspiratory muscle training (IMT) as a non-intrusive and practical intervention to stimulate improved functional fitness in adults with obesity. As excess adiposity of the chest impedes the mechanics of breathing, targeted re-training of the inspiratory muscles may ameliorate sensations of breathlessness, improve physical performance and lead to greater engagement in physical activity.

**Methods:** Sixty seven adults (BMI =  $36 \pm 6.5$ ) were randomized into either an experimental (EXP: n = 35) or placebo (PLA: n = 32) group with both groups undertaking a 4-week IMT intervention, comprising daily use of an inspiratory resistance device set to 55% (EXP), or 10% (PLA) of maximum inspiratory effort.

**Results:** Inspiratory muscle strength was significantly improved in EXP (19.1 cmH<sub>2</sub>O gain; P < 0.01) but did not change in PLA. Additionally, the post training walking distance covered was significantly extended for EXP (P < 0.01), but not for PLA. Bivariate analysis demonstrated a positive association between the change (%) of performance in the walking test and BMI (r = 0.78; P < 0.01) for EXP.

**Conclusion:** The findings from this study suggest IMT provides a practical, self-administered intervention for use in a home setting. This could be a useful strategy to improve the functional fitness of obese adults and perhaps lead to better preparedness for engagement in physical activity initiatives.

© 2016 Published by Elsevier B.V.

## 1. Introduction

Adults with obesity commonly experience shortness of breath at rest and during exercise compared to healthy normal weight adults (Ladosky and Botelho, 2001; Luce, 1980; Mandal and Hart, 2012; Salome et al., 2008; Villiot-Danger et al., 2011). This is typically due to excess adiposity of the chest which impedes the actions of the inspiratory muscles, leading to an inability to exercise effectively and is associated with conditions such as obesity hypoventilation syndrome and sleep apnea (Al Dabal and Bahammam, 2009; Olsen and Zwilllich, 2005). As physical inactivity exacerbates breathing inadequacy by detraining inspiratory and skeletal muscles (Edwards et al., 2008; Salome et al., 2008; Villiot-Danger et al., 2011) the primary purpose of this study was to examine whether or not an inspiratory muscle training (IMT) pro-

gramme undertaken in a home setting might both strengthen the muscles of respiration of adults with obesity and thereby increase their capacity to perform exercise (Edwards et al., 2012). The application of such an unobtrusive, self-administered and practical intervention might prove a meaningful intervention for wider scale implementation. Improved performance of detrained inspiratory muscles in people with obesity would be expected to enable greater capacity to engage and perform exercise through improvements to breathing (Ladosky and Botelho, 2001) but as yet few studies have examined this issue among out-patients (Arena and Cahalin, 2014; Edwards et al., 2012), although, encouraging gains have been demonstrated among athletic groups (Edwards et al., 2008; Griffiths and McConnell, 2007; Romer et al., 2002).

Many physical activity interventions have been developed which aim to improve health outcomes for adults with obesity by reducing excess body weight (Villiot-Danger et al., 2011). However, the effectiveness of exercise is often restricted by factors associated with premature fatigue, such as breathlessness (Luce, 1980; Salome et al., 2008). Such sensations of fatigue could diminish the

\* Corresponding author at: Faculty of Sport & Health Sciences, University of St Mark & St John, Plymouth, United Kingdom.

E-mail address: [aedwards@marjon.ac.uk](mailto:aedwards@marjon.ac.uk) (A.M. Edwards).

**Table 1**  
Participants' characteristics.

	EXP (n = 35)	PLA (n = 32)
Age (years)	46 ± 7.5	48 ± 11.0
Height (cm)	172.4 ± 9.1	169.9 ± 10.8
Mass (kg)	107.3 ± 33.6	101.5 ± 26.6
BMI (kg/m <sup>2</sup> )	36.8 ± 7.4	35.2 ± 5.9
Systolic blood pressure (mm Hg)	135.9 ± 15.1	135.7 ± 13.7
Diastolic blood pressure (mm Hg)	87.8 ± 13.5	88.7 ± 20.9

Mean ± SD. There were no statistically significant differences between the physical characteristics of EXP and PLA groups.

motivational drive to commence a physical training programme or affect the sustainability of participation (Edwards and Polman, 2013; Edwards and Walker 2009; Ekkekakis, 2009).

The act of inspiration is the primary cause of work when breathing. This occurs whereby the chest and lungs expand to accommodate an increased volume of air, while expiration is largely passive, particularly when resting or only exercising at moderate intensity (Otis et al., 1950). Consequently, a pre-exercise training programme specifically designed to enhance the performance of inspiratory muscles for adults with obesity might lessen subconscious inhibition of exercise performance (Ekkekakis et al., 2009), reduce respiratory muscle fatigue (Salome et al., 2008) and promote improved performance in response to exercise challenges (Edwards et al., 2008; Edwards and Cooke, 2004). In support of this perspective, a study of hospitalised obese adults demonstrated an aggressive two month intervention of supervised respiratory (inspiratory and expiratory) muscle training coupled with diet and physical training significantly improved both respiratory muscle endurance and the distance covered in a 6-min walking test (~11% gain) (Villiot-Danger et al., 2011). While the results of that experiment strongly suggest respiratory muscle training may be of value to obese individuals, its findings are not directly applicable to non-hospitalised individuals due to the multidimensional nature of the intervention and the supervisory requirements of such an intense protocol. A less aggressive, but potentially equally effective strategy, is via inspiratory muscle training (IMT) using a portable inspiratory-resistance training device (Edwards, 2013; Edwards et al., 2012).

As obese individuals are well known to experience shortness of breath to a greater extent than healthy normal subjects (Salome et al., 2008) it is therefore likely that a programme of IMT training will be particularly meaningful for obese individuals. The aim of this study is therefore to investigate whether a programme of IMT will improve inspiratory muscle strength and functional performance as assessed by the self-paced 6-min walk test (Enright, 2003).

## 2. Material and methods

### 2.1. Participants

Sixty seven adults (37 males and 30 females) volunteered for this study, provided written informed consent prior to participation and were randomly allocated to either experimental (EXP: n = 35; m = 19, f = 16) or placebo (PLA: n = 32; m = 18, f = 14) group as matched parallel pairs based on body mass index (BMI) and history of smoking. Inclusion criteria were (i) BMI > 27 kg/m<sup>2</sup> and (ii) being free of respiratory or cardiovascular diseases. The physical characteristics of the two groups are shown in Table 1. Ethical clearance for this study was provided by the Research and Ethics committee of James Cook University.

### 2.2. Study overview

Baseline physical assessments were made of mass, height, blood pressure, standard spirometry (FVC, FEV<sub>1</sub>), maximal inspiratory muscle pressure (MIP), 6-min walk test performance and estimation of maximal aerobic power ( $\dot{V} O_2 \text{ max}$ ). Following these measures, all individuals undertook familiarization with a portable inspiratory-resistance training device (PowerBREATHE, UK). This device was pre-set to either 55% of individualized maximal inspiratory effort (EXP) or to the minimum device setting equivalent to approximately 10% of maximal inspiratory effort (PLA) and thereafter used during the experiment (Edwards, 2013). Over the 4-week period, both groups performed 2 × 30 daily inspiratory efforts (Edwards et al., 2012; Volianitis et al., 2001). The assessments were then repeated following the 4-week intervention. Adherence and compliance to the training protocol were regularly checked and no participants reported experiencing issues or difficulties.

### 2.3. Study procedures

#### 2.3.1. Lung function and inspiratory muscle performance

Spirometry measurements were undertaken at baseline and repeated post-programme. These included forced vital capacity (FVC) and forced expiratory volume in 1 s (FEV<sub>1</sub>). These procedures were completed using a hand held training device (Microlab-Spirometry SN M20364, USA).

In addition to standard spirometry measures, maximal static inspiratory mouth pressure (MIP) was also measured. This was assessed at residual volume following a slow and complete expiration using a mouth pressure meter (PowerBREATHE KH1 INSPIRATORY METER, Gaiam, UK). The best of three maximal efforts were analysed for all measures. These procedures were completed similarly with our earlier methodology (Edwards et al., 2012).

#### 2.3.2. Functional exercise capacity

Participants were instructed to “walk as far as you can in six minutes without running or jogging” in accordance with previously validated techniques for a 6-min walk test (Gibbons et al., 2001). Distance covered (m) and heart rates were recorded at the conclusion of the 6-min period. This test is a clinically relevant and common procedure which provides an effective measure of functional walking capacity in untrained, sedentary adults (Enright, 2003; Hulens et al., 2003; Gibbons et al., 2001).


Using a validated heart rate derived algorithm, maximal aerobic power ( $\dot{V} O_2 \text{ max}$ ) was estimated from a sub-maximal single stage 4-min walking test (Ebbeling et al., 1991). All participants were requested to perform an individually determined brisk and constant walking pace ranging from 3.5 to 5 km/h for 4-min on a treadmill in accordance with the protocol.

The study participants were required to wear a heart rate watch and a chest strap transmitter (Polar, T31 Coded Transmitter, Australia) during exercise testing.

The CR10 Borg Scale was used to ascertain ratings of perceived exertion RPE as an index of fatigue perception in response to exercise (Borg, 1982).

### 2.4. Statistical analyses

Statistical software package SPSS (version 18.0, SPSS, Chicago, Illinois) was used for all statistical analyses. Parametric pre- and post-training results and group interactions were statistically compared using two-way repeated measures analyses of variance (group × time) (ANOVA). Post hoc Tukey tests were used to examine differences between data sets where indicated by ANOVA. Associations were examined using Pearson Product Moment Correlations. To ascertain an appropriate sample size for the study, analysis was



**Fig 1.** Distance covered (metres) in response to the 6-min walk test for both experimental (EXP;  $n=35$ ) and placebo (PLA;  $n=32$ ) groups. \* = significant difference between baseline and post-training distance covered ( $P<0.01$ ). Means  $\pm$  SD and individual (before and after training) results are displayed.

**Table 2**

Lung function and maximal aerobic power variables prior to and following the 4-week intervention.

	EXP		PLA	
	Pre ( $n=35$ )	Post ( $n=35$ )	Pre ( $n=32$ )	Post ( $n=32$ )
FVC (l)	$3.3 \pm 0.9$	$3.2 \pm 0.7$	$3.1 \pm 0.8$	$3.1 \pm 0.7$
FEV <sub>1</sub> (l)	$2.7 \pm 0.8$	$2.8 \pm 0.9$	$2.6 \pm 0.7$	$2.7 \pm 0.5$
$\dot{V}O_2$ max (ml/kg/min)	$39.4 \pm 10.1$	$39.5 \pm 10.3$	$38.3 \pm 11.1$	$38.5 \pm 11.2$

Mean  $\pm$  SD.

based on an anticipated mean improvement (SD) in the six minute walk test of those in the EXP in the PLA group (Edwards et al., 2012; Edwards, 2013). Probability values of  $<0.05$  were considered significant and all tests were two sided. All results are expressed as means (SD) unless otherwise stated.

### 3. Results

Evaluation of distance covered in response to the 6-min walking test revealed a significant group  $\times$  time ANOVA interaction. As expected, there was no difference between groups at baseline. Within-group comparisons for time (pre- to -post-training) indicated EXP significantly improved distance covered (m) in response to the 6-min walk test from baseline to post-training ( $60.6 \pm 25.7$  m gain;  $P<0.01$ ). Conversely, the distance covered by PLA was not significantly extended over the 4-week intervention period ( $13.3 \pm 35.9$  m gain; NS) (Fig. 1).

The estimation of  $\dot{V}O_2$  max in response to treadmill walking did not identify a significant difference between EXP and PLA at either baseline or after the intervention (Table 2). Additionally, assessment of standard spirometry variables (FVC and FEV<sub>1</sub>) also did not identify differences between groups at either baseline or post-training (Table 2).

The MIP assessment revealed a significant group  $\times$  time ANOVA interaction effect ( $P<0.01$ ). Subsequent post hoc Tukey HSD test evaluation demonstrated MIP improved significantly over the 4-week intervention for EXP ( $66.7 \pm 10.5$ – $85.8 \pm 9.3$  cm H<sub>2</sub>O;  $P<0.01$ ). However, MIP did not significantly change for PLA ( $68.4 \pm 11.7$ – $77.7 \pm 10.8$  cmH<sub>2</sub>O; NS). There was a between group difference following the intervention where EXP demonstrated significantly greater MIP than PLA ( $P<0.01$ ).

Heart rate responses to the 6-min walk test were unchanged for both EXP ( $124 \pm 14$  and  $121 \pm 15$  b/min) and PLA ( $118 \pm 15$  and  $116 \pm 11$  b/min) from pre- to post-training.

RPE evaluations undertaken after exercise were not different between groups and did not change significantly from baseline to post-training in either EXP ( $2.7 \pm 0.7$ – $2.7 \pm 0.8$ ) or PLA ( $2.7 \pm 1.7$ – $2.9 \pm 1.8$ ).

A significant correlation was observed between % change of distance covered in the 6-min walk test (pre- to post-training) and baseline BMI ( $r=0.78$ ;  $P<0.01$ ). This effect between a participant factor and intervention response was specific to EXP. There were no meaningful associations identified in PLA.

### 4. Discussion

The main finding of this study was that a 4-week period of inspiratory muscle training (IMT) appears efficacious for improving inspiratory muscle strength and the functional fitness of obese and overweight participants. As these effects were not evident in PLA, it suggests that IMT may be a meaningful intervention with which to augment physical performance outcomes for overweight and obese individuals. The results of our study support and exceed those from our earlier pilot data (Edwards et al., 2012) and also from hospitalised obese individuals (Villiot-Danger et al., 2011). As these results were achieved with a considerably less aggressive intervention it seems likely that such a practical technique might be suitable for wider implementation.

In our study, post-test evaluations of perceived exertion did not differentiate the groups, despite a significant improvement in walking performance for EXP. This suggests individuals may have paced themselves according to physical sensations (Suzuki et al., 1995, such as a tolerable level of physical discomfort the individuals were prepared to endure in the 6-min task (Ekkakakis, 2009; Edwards and Polman, 2013). As such, participants would (and did) experience the same level of tolerable physical discomfort during the 6-min walk test at both baseline and post-training. The observed performance difference would therefore not be expected to be evident in a change to the perceived exertion, but rather in a changed (improved) outcome of a greater walking distance covered for the same perceived level of exertion.

Bivariate analysis revealed an interesting association between data sets whereby the (%) change from baseline to post-training in distance covered for the 6-min walk test was positively related to BMI for EXP ( $r=0.78$ ;  $P<0.01$ ). This shows that individuals with a higher BMI gained the biggest change in performance from an IMT intervention. This is possibly due to greater post-intervention resistance to fatigue of inspiratory muscles, although further work is required to confirm the mechanisms responsible for this effect (Verges et al., 2007; Zerah et al., 1993).

MIP results for EXP remained beneath levels reported for healthy subjects (Volianitis et al., 2001) suggesting that continuation of IMT beyond a 4-week period could be meaningful to an obese population where detraining effects may be substantial. There are very limited data in this area and, therefore, further studies may elucidate whether extending the period of IMT prior to physical training and also utilising concurrent (IMT and physical) training improve performance outcomes for obese individuals.

Although this study was much larger and robust than our previous pilot study (Edwards et al., 2012) it did not include a concurrent IMT strategy with exercise intervention. The use of IMT in conjunction with an exercise training intervention could be expected to further augment performance as has been the case in athletes (Edwards et al., 2008). Nevertheless, a 4-week period of IMT demonstrates the usefulness of the technique over a short period, but a longitudinal intervention with and without concurrent exercise would be worthwhile to determine whether training effects are sustainable.

In summary, IMT may provide a practical, minimally intrusive intervention to augment both inspiratory muscle strength and walking distance among overweight and obese adults. The beneficial effects of this treatment were similar to those previously reported from vigorous, supervised training among hospitalised obese patients (Villiot-Danger et al., 2011). Our findings indicate similar effects could be expected without the need for hospitalisation and indicate that IMT can easily be performed in the home environment. Therefore, IMT appears a useful strategy to enhance walking performance in overweight and obese individuals which may prove to be a meaningful priming (pre-exercise) intervention with which to stimulate performance adaptations and greater future engagement with physical activity.

### Funding

No external funding was provided for this project.

### Ethical approval

Ethical approval for this project was provided by the Human Research and Ethics Committee of James Cook University (ref: H5450).

### Conflict of interest

None of the authors had a conflict of interest regarding any aspect of this work.

### Acknowledgement

The authors would like to thank research students for their support in data collection.

### References

- Al Dabal, L., Bahammam, A.S., 2009. Obesity hypoventilation syndrome. *Ann. Thorac. Med.* 4, 41–49.
- Arena, R., Cahalin, L.P., 2014. Evaluation of cardiorespiratory fitness and respiratory muscle function in the obese population. *Prog. Cardiovasc. Dis.* 56, 457–464.
- Borg, G., 1982. Psychophysiological bases of perceived exertion. *Med. Sci. Sports Exerc.* 14, 377–387.
- Ebbeling, C., Ward, A., Puleo, E., Widrick, J., Rippe, J.M., 1991. Development of a single-stage submaximal treadmill walking test. *Med. Sci. Sports Exerc.* 23, 966–973.
- Edwards, A.M., Cooke, C., 2004. Oxygen uptake kinetics and maximal aerobic power are unaffected by inspiratory muscle training in healthy subjects where time to exhaustion is extended. *Eur. J. Appl. Physiol.* 93, 139–144.
- Edwards, A.M., Polman, R.C.J., 2013. Pacing and awareness: brain regulation of physical activity. *Sports Med.* 43, 1057–1063.
- Edwards, A.M., Walker, R., 2009. Inspiratory muscle training and endurance: a central metabolic control perspective. *Int. J. Sports Physiol. Perform.* 4, 122–128.
- Edwards, A.M., Wells, C., Butterly, R.J., 2008. Concurrent inspiratory muscle and cardiovascular training differentially improves both perceptions of effort and 5000 m running performance compared with cardiovascular training alone. *Br. J. Sports Med.* 42, 523–527.
- Edwards, A.M., Graham, D., Maguire, G., Boland, V., Richardson, G., 2012. The effect of inspiratory muscle training on walking performance in overweight and obese individuals. *J. Obes.*, 1–6.
- Edwards, A.M., 2013. Respiratory muscle training extends exercise tolerance without concomitant change to peak oxygen uptake: physiological, performance and perceptual responses derived from the same incremental exercise test. *Respirology* 18, 1022–1027.
- Ekkekakis, P., Lind, E., Vazou, S., 2009. Affective responses to increasing levels of exercise intensity in normal-weight, overweight, and obese middle-aged women. *Obesity* 18, 79–85.
- Ekkekakis, P., 2009. Let them roam free: physiological and psychological evidence for the potential of self-selected exercise intensity in public health. *Sports Med.* 39, 857–888.
- Enright, P., 2003. The six-minute walk test. *Respir. Care* 48, 783–785.
- Gibbons, W., Fruchter, N., Sloan, S., Levy, R., 2001. Reference values for a multiple repetition 6-minute walk test in healthy adults older than 20 years. *J. Cardiopulm. Rehabil.* 21, 87–93.
- Griffiths, L.A., McConnell, A.K., 2007. The influence of inspiratory and expiratory muscle training upon rowing performance. *Eur. J. Appl. Physiol.* 99, 457–466.
- Hulens, M., Vansant, G., Claessens, A., Lysens, R., Muls, E., 2003. Predictors of 6-minute walk test results in lean: obese and morbidly obese women. *Scand. J. Med. Sci. Sports* 13, 98–105.
- Luce, J., 1980. Respiratory complications of obesity. *Chest* 78, 626–631.
- Mandal, S., Hart, N.I., 2012. Respiratory complications of obesity. *Clin. Med.* 12, 75–78.
- Olsen, A.L., Zwillich, C., 2005. The obesity hypoventilation syndrome. *Am. J. Med.* 118, 948–956.
- Otis, A., Fenn, W., Rahn, H., 1950. Mechanics of breathing in man. *J. Appl. Physiol.* 2, 592–607.
- Romer, L.M., McConnell, A.K., Jones, D.A., 2002. Effects of inspiratory muscle training on time-trial performance in trained cyclists. *J. Sports Sci.* 20, 547–590.
- Salome, C., Munoz, P., Berend, N., Thorpe, C.W., Schachter, L.M., King, G.G., 2008. Effect of obesity on breathlessness and airway responsiveness to methacholine in non-asthmatic subjects. *Int. J. Obes.* 32, 502–509.
- Suzuki, S., Sato, M., Okubo, T., 1995. Expiratory muscle training and sensation of respiratory effort during exercise in normal subjects. *Thorax* 50, 366–370.
- Verges, S., Lenherr, O., Haner, A., Schulz, C., Spengler, C., 2007. Increased fatigue resistance of respiratory muscles during exercise after respiratory muscle endurance training. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 292, R1246–R1253.
- Villiot-Danger, J., Villiot-Danger, E., Borel, J., Pepin, J.L., Wuyam, B., Verges, S., 2011. Respiratory muscle endurance training in obese patients. *Int. J. Obes.* 35, 692–699.
- Volianitis, S., McConnell, A.K., Koutedakis, Y., McNaughton, L.R., Backx, K., Jones, D.A., 2001. Inspiratory muscle training improves rowing performance. *Med. Sci. Sports Exerc.* 33, 803–809.
- Zerah, F., Harf, A., Perlemuter, L., Lorino, H., Lorino, A., Atlan, G., 1993. Effects of obesity on respiratory resistance. *Chest* 103, 1470–1476.