

Health & Research Journal

Vol 8, No 3 (2022)

Volume 8 Issue 3 July - September 2022



Volume 8 Issue 3 July - September 2022

EDITORIAL

DIGITAL HEALTH: IS IT THE FUTURE FOR OUR LIVES?

RESEARCH ARTICLES

RENAL BIOMARKER COMBINATIONS PREDICT EARLY CRRT NEED IN A MIXED ICU POPULATION

TRANSLATION AND PRELIMINARY VALIDATION OF THE MODIFIED MINI MENTAL STATE (3MS) IN A GREEK SAMPLE

CLINICAL OUTCOMES OF ENHANCED RECOVERY AFTER SURGERY PROTOCOL FOR HEPATO-PANCREATO-BILIARY SURGERY: A FIVE-YEAR EXPERIENCE FROM A HELLENIC ONCOLOGICAL HOSPITAL

USE OF HERBAL PRODUCTS AND FOOD SUPPLEMENTS OF TYPE 2 DIABETES PATIENTS

SUBJECTIVE SLEEP QUALITY AND DAYTIME SLEEPINESS AMONG GREEK NURSING STAFF: A MULTICENTER CROSS-SECTIONAL STUDY

SYSTEMIC REVIEW

IMMERSIVE VIRTUAL REALITY IN COGNITIVE REHABILITATION: A SYSTEMATIC REVIEW

THE BENEFITS OF INSPIRATORY MUSCLE TRAINING IN HEART FAILURE PATIENTS - A SYSTEMATIC REVIEW

Published in cooperation with the Postgraduate Program "Intensive Care Units", the Hellenic Society of Nursing Research and Education and the Hellenic Society of Intensive Care Medicine

The Benefits of Inspiratory Muscle Training in Heart Failure Patients – A Systematic Review

Christina Katsoula, Ioannis Vasileiadis, Christina Routsis, Serafeim Nanas, Eleftherios Karatzanos

doi: [10.12681/healthresj.28185](https://doi.org/10.12681/healthresj.28185)

To cite this article:

Katsoula, C., Vasileiadis, I., Routsis, C., Nanas, S., & Karatzanos, E. (2022). The Benefits of Inspiratory Muscle Training in Heart Failure Patients – A Systematic Review. *Health & Research Journal*, 8(3), 242–266. <https://doi.org/10.12681/healthresj.28185>

SYSTEMATIC REVIEW

THE BENEFITS OF INSPIRATORY MUSCLE TRAINING IN HEART FAILURE PATIENTS: A SYSTEMATIC REVIEW

Christina Katsoula¹, Ioannis Vasileiadis¹, Christina Routsis¹, Serafeim Nanas¹, Eleftherios Karatzanos¹

1. Clinical Ergospirometry, Exercise and Rehabilitation Laboratory, 1st Critical Care Department, Evangelismos Hospital, School of Medicine, National & Kapodistrian University of Athens, Greece

Abstract

Background: Patients with heart failure (HF) have impaired function of respiratory system and frequently experience dyspnea. Inspiratory muscle training (IMT) offers an alternative way of exercise with a lot of benefits for HF patients.

Aim: The aim of this review was to summarize and to reveal the effects of IMT in HF patients.

Methods: Electronic searches were performed using Pubmed Database, Physiotherapy Evidence Database (PEDro) and Cochrane Library. Inclusion criteria were: RCTs, patients with HF, full text articles after 2010 and at least one intervention group with IMT. Methodological quality was assessed using the PEDro (Physiotherapy Evidence Database) scale.

Results: Nineteen articles met the inclusion criteria and were included in this review. In most studies training protocols involved 3 to 7 sessions per week with intensity $\leq 30-60\%$ of maximal inspiratory pressure (MIP), for a total duration of 4-12 weeks. Respiratory muscle strength improved in 11/12 studies, peripheral muscle strength in 4/5 studies, exercise capacity (peak VO_2 , VE/VCO_2 slope) in 4/10, pulmonary function (FEV_1 , FEV_1/FVC , FVC) in 0/5, functional capacity (6MWT) in 6/8, echocardiography parameters in 1/6, quality of life and dyspnea in 9/16. Control groups followed sham IMT, usual care, no intervention, intervention without exercise, or aerobic training, which could explain in some cases the conflicting results.

Conclusions: IMT is beneficial for HF patients and should be included as a complementary method in cardiac rehabilitation programs. The optimal characteristics of IMT as well as the benefits when combined with common forms of exercise need further research.

Keywords: Inspiratory muscle training, respiratory muscle training, inspiratory resistance training, breathing exercise, heart failure.

Corresponding Author: Ioannis Vasileiadis, M.D., Ph.D., Associate Professor of Internal Medicine and Critical Care National and Kapodistrian University of Athens, School of Medicine, 1st Critical Care Department, Evangelismos Hospital, Ipsilantou 45-47, 10676, Athens, Greece, Email: ioannisvmed@yahoo.gr, ivasileiadis@med.uoa.gr, Tel.: +30 2132043385, +30 6977644866, ORCID ID: 0000-0002-9529-9361

Cite as: Katsoula, C., Vasileiadis, I., Routsis, C., Nanas, S., Karatzanos, E. (2022). The benefits of inspiratory muscle training in heart failure patients: a systematic review. *Health and Research Journal*, 8(3), 242-263. <https://ejournals.epublishing.ekt.gr/index.php/HealthRes/>

INTRODUCTION

Chronic heart failure (CHF) is a complex disease characterized by a wide range of symptoms (e.g. dyspnea, muscle fatigue) and signs (e.g. pulmonary dysfunction, abnormal breathing patterns).^{1,2} Several skeletal muscle and respiratory abnormalities including impaired resting lung function, peripheral muscle microcirculation and inefficient ventilation have been identified in CHF patients that probably play an important role in the pathophysiology of exercise intolerance.³⁻⁵ Also, CHF patients demonstrate increased respiratory drive at rest and abnormal breathing pattern during exercise which is associated with disease severity.⁶ Moreover, patients with CHF present impaired inspiratory muscle strength and endurance, an additional factor which is associated with limited exercise and poor prognosis.⁷ There is evidence of benefits from IMT applied in CHF patients, including improvement of respiratory muscle strength, functional capacity, ventilation, quality of life and decreased dyspnea.⁸⁻¹²

The aim of this review is to summarize and shed light on the effects of IMT in CHF patients, as well as to provide more reliable estimates regarding the usefulness of respiratory training as a complimentary method in the HF rehabilitation programs.

METHODOLOGY

Data sources

The literature research was performed in the following electronic databases: Pubmed, Cochrane library and Physiotherapy Evidence Database (Pedro). The keywords used in the search were: inspiratory muscle training, IMT, respiratory muscle training, respiratory exercise, inspiratory resistance training, breathing exercise, heart failure, CHF. These terms were appropriately combined using the booleans OR and AND. For example, the search strategy in Pubmed database was as follows: (Inspiratory muscle training OR IMT OR Respiratory muscle training OR Respiratory training OR Respiratory exercise OR Inspiratory resistance training OR Breathing exercise) AND (Heart failure OR CHF). The search was restricted by the following criteria: Clinical Trials, Full Text studies, humans and English language, using appropriate filters.

Eligibility Criteria and Data Collection

We included studies with or without combined exercise programs. Inclusion criteria were the following: 1) RCTs, 2) HF patients, 3) full text articles, 4) articles published after 2010. Exclusion criteria were: 1) animal studies, 2) languages other than English.

Our data presented patient characteristics (number of patients included, age, gender, diagnosis and disease severity), intervention (type, duration, frequency, resting period) and outcomes related to respiratory muscle strength, pulmonary function, exercise capacity, peripheral muscle function, functional capacity, quality of life and dyspnea.

Methodological quality

The methodological quality of the studies was evaluated by the PEDro scale. The Pedro scale is based on 11 items to assess scientific rigor related to eligibility criteria, allocation, baseline between-group comparability, blindness of subjects, therapists, and assessors, follow-up, statistical analysis, and results reporting¹³. Based on the total score, studies were categorized as high- (> 5), moderate- (4-5), or low- (< 4) quality.

RESULTS

Study selection

The initial search led to retrieval of 1798 articles. After 1350 duplicates were removed, 450 records were screened at abstract level. The full text of 31 studies was assessed for eligibility. Of those 31 studies, 12 were excluded being published before 2010. Finally, 19 RCTs were included in the present review.¹⁴⁻³² The PRISMA flow diagram is presented in figure 1.

Methodological quality

The Pedro score for the included studies is presented in Table 1. The score ranged from 5 to 8 (that is, moderate to high quality). The allocation of subjects was concealed in 6 studies.^{14,20,21,24,27,31} Furthermore, all the studies provided baseline data ensuring between-group comparability. All the studies met the follow up criteria as described by Pedro scale. Ten of the studies were of high methodological quality (PEDro score>5)^{14-17,20,23,26,27,31} and nine studies were of moderate quality (PEDro score

=5).^{18,19,22,24,25,28,29,30,32} The level of evidence could be considered as moderate to strong.

Participants

The characteristics of the patients included in this systematic review are presented in Table 2. We included data that were relevant for at least one of the main outcomes of interest. Data from 726 patients were analyzed. Disease severity was categorized by the NYHA classification. Two studies had CHF patients with preserved ejection fraction.^{26, 27} The remaining 17 studies had CHF patients with reduced ejection fraction. One study investigated stroke patients with stable CHF²³ and another study included patients with implanted left ventricular assist device.¹⁶ Inspiratory muscle weakness was identified in some of the patients included in the study (MIP < 70% of the predicted value).^{14,19, 21,23}

Interventions

The characteristics of the interventions are shown in Table 2. IMT duration ranged, in most studies, from 4 to 12 weeks.^{14-24,26,28} One study lasted 4 months.²⁵ Two studies started with 10-12 weeks IMT, followed by 10-12 weeks usual care (crossover studies).^{29,31} Three studies used follow up evaluation: one study at 5 months³⁰ and two studies at 6 months.^{27,32} Concerning the load, 9 studies used loads > 30% to 60% of maximal inspiratory pressure (MIP) or sustained MIP (SMIP).^{15-18,20-23,25} One study used training loads which allowed the performance of 10 consecutive maximal repetitions (10RM), so the training intensity was 100% of their 10 RM.¹⁴ Five studies used load ≤ 30% of MIP.^{19, 21, 24, 26, 27} Frequency of training ranged from 3 to 7 times per week with duration of each session, mostly, up to 30 minutes, with 1 or 2 sessions per day and with a common respiratory rate of 15-20 breaths/min. In relation to the training methods employed, IMT was performed using an inspiratory muscle trainer with training loads adjusted to the inspiratory pressure,¹⁴⁻²⁷ or using a device-guided slow breathing via the introduction of low and high-frequency tones to entrain exhalation and inhalation,^{28,29,31,32} or using deep and slow diaphragm breathing training according to voice-guided directions.³⁰

Concerning the control groups, patients performed IMT with low or no respiratory load,^{14,17,19-21} aerobic training,^{15,16} combined

training (aerobic with IMT or with strength exercises),^{18,22} usual care treatment,^{26,27,29,31} intervention without exercise^{23,28,30} and no intervention.^{24,25}

Functional assessments

Respiratory muscle strength

Five studies evaluated the effect of combined exercise/IMT vs. control group on respiratory muscle strength;^{15,16,18,22,24} MIP improved in two studies,^{22,24} SMIP^{15,18} improved also in two studies, maximal expiratory pressure (MEP) improved in one study,²⁴ whereas nonsignificant differences were observed in MIP in three studies^{15,16,18} and in SMIP in one study.¹⁶ Seven studies^{14,17,19,21,23,25,26} examined IMT alone with no other form of exercise training; intervention groups demonstrated significant improvement in MIP (all studies) and in MEP (two studies),^{14,17} compared to control groups, while no significant differences were found in MEP in one study.²³

Peripheral muscle strength

Five studies assessed peripheral muscle strength.^{14,15,17,22,24} Three studies^{15,22,24} used combined exercise/IMT vs. control group and evaluated lower limb muscle strength (quadriceps femoris). The results revealed significantly higher muscle strength and endurance in the treatment groups compared with the control groups. The remaining two studies^{14,17} examined IMT alone with no other form of exercise training, with conflicting results. One study¹⁴ evaluated upper limb muscle strength (handgrip strength) and revealed no significant differences between groups, while the other one¹⁷ found greater improvement of peripheral muscle strength and functional balance in the treatment group compared to controls.

Pulmonary function

Five studies^{16-18,23,24} evaluated the effect of IMT on pulmonary function, mostly employing: forced expiratory volume in 1 s (FEV₁), forced vital capacity (FVC), FEV₁/FVC; no significant differences were observed between the intervention and control groups.

Exercise capacity

Six studies assessed the effects of combined exercise/IMT vs. control group using cardiopulmonary exercise testing (CPX).^{15,16,18,22,26,27} Variables mostly employed were peak oxygen uptake (peak VO₂), ventilation/carbon dioxide slope, VE/VCO₂ slope and VE. Four of these studies^{15,16,18,22} revealed no significant differences in most CPX parameters between the intervention and control groups, whereas the other two studies^{26,27} found significant improvement of CPX parameters in the trained patients.

Five studies investigated the effects of IMT alone with no other form of exercise training vs. control group^{19,21,25,27,32} on cardiopulmonary function. Two studies revealed no significant differences between the intervention and control groups in most CPX parameters.^{19,32} In the remaining three studies, trained patients presented significantly improved CPX parameters compared to controls.^{21,25,27}

Echocardiographic parameters

Four studies evaluated the effects of combined exercise/IMT vs. control group on echocardiographic parameters and reported no significant differences between the intervention and control groups.^{15,18,26,27} Three studies^{27,31,32} evaluated the effects of IMT alone with conflicting results; one study³¹ reported significant improvement in most echocardiographic parameters, while the other two studies^{27,32} reported no significant changes in echocardiographic parameters.

Functional capacity

Four studies^{16,24,26,27} assessed the effect of combined exercise/IMT vs. control group on functional capacity using the six minute walk test (6MWT). Conflicting results emerged. Three studies^{24,26,27} found that the distance covered during the 6MWT significantly increased in the treatment groups compared to control groups, whereas one study¹⁶ found no significant differences between groups. Five studies^{17,27,30-32} used IMT alone and all but one³⁰ showed significant improvement in the treatment groups compared to control groups.

Quality of life and dyspnea

Seven studies^{15,16,18,22,24,26,27} evaluated the effect of combined exercise/IMT vs. control group on quality of life (QoL) and dyspnea using several assessment methods. Three studies found no significant changes in QoL^{16,22,24} and dyspnea¹⁶ between the intervention and control groups, whereas four studies reported significant improvement in QoL^{15,18,26,27} and dyspnea^{15,18} in the intervention compared to control groups. Ten studies used IMT alone^{14,17,19-21,25,27-30} with conflicting results; four studies found no significant differences between the intervention and control groups regarding QoL^{14,17,29} and dyspnea^{14,28,30} while in six studies the intervention groups showed significant improvement in QoL^{19,21,25,27} and dyspnea^{17,19,20} compared to control groups.

Functional classification

Three studies^{15,18,24} assessed the effect of combined exercise/IMT vs. control group on functional status using the New York Heart Association (NYHA) classification and reported no significant differences between the intervention and control groups. Two studies^{20,28} used IMT alone with conflicting results; NYHA class improved in the training group compared to the control group in one study,²⁰ while the other one²⁸ did not show significant differences between groups.

DISCUSSION

This systematic review indicates that IMT, isolated or combined with other forms of exercise training, improves respiratory and peripheral muscle strength, whereas it seems to have no effect on pulmonary function. Furthermore, the included studies demonstrate conflicting results after IMT application regarding the other parameters of interest: exercise and functional capacity, echocardiographic parameters, quality of life and dyspnea. The results were varying among the studies likely because of the different loads (adjusted for MIP/SMIP), the method employed, i.e., *threshold IMT, device-guided breathing and diaphragmatic exercises*, the frequency and duration of intervention as well as the small sample size of the studies. The best mode and training method of IMT is not clear.

All the studies which performed IMT based on SMIP revealed non-significant differences in MIP; utilization of SMIP adjusted load seems to improve the respiratory muscle endurance and

not the MIP.^{15,16,18} This could possibly explain, at least partly, the conflicting results. Analysis of the studies showed that IMT at a low percentage load ($\leq 15\%$ of MIP)^{14,17,19} was not enough for the control groups to reach the same improvements as the intervention groups which used training loads $\geq 30\%$ of MIP. Another study came to the same conclusion, comparing the effect of low intensity training (15% of MIP) and moderate intensity training (30% of MIP); it was shown that only the moderate intensity exercise training improved MEP and NYHA functional classification.²⁴

IMT when combined with a specific program of lower limb muscle resistance training improved peripheral muscle strength.^{15,22,24} IMT when performed without other form of intervention improved also peripheral muscle strength; although this result is a clear indication of an autonomous beneficial effect of this type of exercise training (i.e. IMT) on peripheral muscle strength,¹⁷ there is a need for further research for its final documentation. In another study which used IMT alone, upper limb muscle strength (handgrip strength) did not correlate with MIP.¹⁴

All studies, regardless of using IMT alone or combined with other form exercise training, presented no differences in pulmonary function in between-group comparisons.^{16,17,18,23,24} This result may be due to the relatively good functional status of the lungs at baseline or because of an inappropriate training method, unable to improve the spirometric parameters (FEV₁, FEV₁/FVC, FVC). NYHA status improved in all studies but one;²⁸ the duration and respiratory rate of the device-guided breathing exercise technique could probably account for the result.

Although studies revealed within group improvements in exercise capacity, functional capacity, quality of life and dyspnea after IMT, results were varied in terms of between-group comparisons; in several studies no differences were noted between the intervention and control groups regarding peak VO₂ and VE/VCO₂ slope,^{15,16,18,19,22,32} 6MWT,^{16,30} as well as quality of life and dyspnea;^{14,16,22,24,28-30} this may be due to the fact that in some cases not only the intervention but the control groups also incorporated aerobic exercise in their training programs or they had almost the same training characteristics (duration, rest,

sham-IMT). Furthermore, echocardiographic parameters^{15,18,26,27,32} were not improved in almost all studies.

Patients with preserved ejection fraction^{26,27} showed remarkable improvements in functional capacity, exercise capacity and quality of life, as patients with reduced ejection fraction. Follow up evaluations^{27,30,32} also, revealed that IMT beneficial effects are maintained in most outcomes of interest.

Different training methods may lead to different results: some studies utilize a Threshold IMT breathing trainer for continuously providing a specific resistance (e.g. 30% of MIP) which is controlled, i.e. determined by the training supervisor; in contrast, other studies use other exercise techniques that focus on slow and deep inhalations according to recorded instructions.

Heart and lungs are interconnected, being parts of the system that supplies oxygen to the body organs and tissues. However, peripheral muscle dysfunction and inspiratory muscle weakness seems to be involved in the underlying mechanisms for fatigue, dyspnea and exercise intolerance in patients with HF. At rest and during exercise HF patients hyperventilate, causing structural and biochemical alterations in diaphragmatic muscle.³³ The effect of IMT on respiratory system, functional status and quality of life in patients with HF has been studied by many researchers. Potential mechanisms, underlying the effects of respiratory training, include increased ventilatory efficiency, lower oscillatory ventilation during incremental exercise, improved recovery oxygen uptake kinetics, as well as reduced sympathetic nervous activity.^{8,21} The use of IMT, also, attenuates the mechanisms involved in respiratory muscle metaboreflex in CHF patients, improving blood flow in peripheral muscles during inspiratory loading, in resting and exercising limbs.^{19,34}

In a previous review by Lin et al,³⁵ which included studies with isolated or combined IMT, it was shown that IMT significantly improves respiratory muscle strength and functional capacity, whereas, the effect of IMT on quality of life was inconsistent; this could be due to the different evaluation questionnaires being used. The improvement in inspiratory muscle strength could be translated to an improvement in functional capacity as CHF patients demonstrate low functional capacity due to respiratory fatigue and dyspnea.^{8,36}

In another review by Neto et al,³⁷ combined exercise along with IMT offered additive benefits in quality of life compared with conventional aerobic training alone although, no extra benefits were established regarding the exercise capacity. This improvement in quality of life may be related to the benefits of IMT in cardiovascular and respiratory response as well as in the reduction of dyspnea.

Limitations

There were several limitations in the articles we reviewed. Initially, the sample size in all studies was small. There is a need for more randomized controlled trials in future with well-defined protocols and larger samples. Articles with device-guided paced breathing in HF patients were limited, as well as studies in patients with preserved ejection fraction and follow up evaluations. In addition, the improvement identified in the control groups in respiratory muscle strength probably occurred because they used IMT at a low percentage of MIP as sham therapy. It would be better if all the studies were designed without any training load in their control groups.

CONCLUSIONS

There is substantial evidence that IMT in patients with HF increases respiratory and peripheral muscle strength and seems to be beneficial regarding functional capacity, exercise capacity, quality of life and dyspnea. IMT should be part of a cardiac rehabilitation program as a complementary method, considering its safety and the demonstrated positive effects. The optimal characteristics of IMT as well as benefits when combined with common forms of exercise need further research. Also, more follow up evaluations are needed to explore the long term effects of respiratory training in patients with heart failure.

Conflict of interest:

Authors declare that there is no conflict of interest.

REFERENCES

1. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. 2016;37(27):2129-200.
2. Tsai MF, Hwang SL, Tsay SL, Wang CL, Tsai FC, Chen CC, et al. Predicting Trends in Dyspnea and Fatigue in Heart Failure Patients' Outcomes. *Acta Cardiologica Sinica*. 2013;29:488-495.
3. Manetos C, Dimopoulos S, Tzannis G, Vakrou S, Tasoulis A, Kapelios C, et al. Skeletal muscle microcirculatory abnormalities are associated with exercise intolerance, ventilator inefficiency, and impaired autonomic control in heart failure. *J Heart and Lung Transplant*. 2011;30(12):1403-8.
4. Nanas S, Nanas J, Papazachou O, Kassiotis C, Papamichalopoulos A, Millic-Emili J, et al. Resting lung function and hemodynamic parameters as predictors of exercise capacity in patients with chronic heart failure. *Chest*. 2003;123(5):1386-93.
5. Meyer FJ, Zugck C, Haass M, Otterspoor L, Strasser RH, Kubler W, et al. Inefficient ventilation and reduced respiratory muscle capacity in congestive heart failure. *Basic Res Cardiol*. 2000;95(4):333-42.
6. Tasoulis A, Dimopoulos S, Repasos E, Manetos C, Tzannis G, Sousonis V, et al. Respiratory drive and breathing pattern abnormalities are related to exercise intolerance in chronic heart failure patients. *Respir Physiol Neurobiol*. 2014;192:90-94.
7. Meyer FJ, Borst MM, Zugck C, Kirschke A, Schellberg D, Kubler W, et al. Respiratory muscle dysfunction in congestive heart failure: clinical correlation and prognostic significance. *Circulation*. 2001;103:2153-4.
8. Dall'Ago P, Chiappa GRS, Guthus H, Stein R, Ribeiro JP. Inspiratory Muscle Training in Patients With Heart Failure and Inspiratory Muscle Weakness: A Randomized Trial. *J Am Coll Cardiol*. 2006;47(4):757-63.
9. Winkelmann ER, Chiappa GR, Lima COC, Vecili PRN, Stein R, Ribeiro JP. Addition of inspiratory muscle training to aerobic training improves cardiorespiratory responses to exercise in patients with heart failure and inspiratory muscle weakness. *Am Heart J*. 2009; 158(5):768.e1-7.

10. Stein R, Chiappa GR, Guths H, Dall'Ago P, Ribeiro J. Inspiratory Muscle Training Improves Oxygen Uptake Efficiency Slope in Patients With Chronic Heart Failure. *J Cardiopulm Rehabil Prev.* 2009;29:392-5.
11. Padula CA, Yeaw E, Mistry S. A home-based nurse-coached inspiratory muscle training intervention in heart failure. *Appl Nurs Res.* 2009;22(1):18-25.
12. Parati G, Malfatto G, Boarin S, Branzi G, Caldara G, Giglio A, et al. Device-Guided Paced Breathing in the Home Setting: Effects on Exercise Capacity, Pulmonary and Ventricular Function in Patients With Chronic Heart Failure: A Pilot Study. *Circ Heart Fail.* 2008;1(3):178-83.
13. Moseley AM, Herbert RD, Sherrington C, Maher CG. Evidence for physiotherapy practice: A survey of the Physiotherapy Evidence Database (PEDro). *Aust J Physiother.* 2002;48(1):43-9.
14. Marco E, Ramirez-Sarmiento AL, Coloma A, Sartor M, Comin-Colet J, Vila J, et al. High-intensity vs. sham inspiratory muscle training in patients with chronic heart failure: a prospective randomized trial. *Eur J Heart Fail.* 2013;15(8):892-901.
15. Laoutaris ID, Adamopoulos S, Manginas A, Panagiotakos DB, Kallistratos MS, Doulaptsis C, et al. Benefits of combined aerobic/resistance/inspiratory training in patients with chronic heart failure. A complete exercise model? A prospective randomized study. *Int J Cardiol.* 2013;167(5):1967-72.
16. Laoutaris ID, Dritsas A, Adamopoulos S, Manginas A, Gouzouta A, Kallistratos MS, et al. Benefits of physical training on exercise capacity, inspiratory muscle function, and quality of life in patients with ventricular assist devices long-term postimplantation. *Eur J Cardiovasc Prev Rehabil.* 2011;18(1):33-40.
17. Bosnak GM, Arikan H, Savci S, Inal ID, Tulumen E, Aytemir K, et al. Effects of inspiratory muscle training in patients with heart failure. *Respir Med.* 2011;105:1671-81.
18. Adamopoulos S, Schmid JP, Dendale P, Poerschke D, Hansen D, Dritsas A, et al. Combined aerobic/inspiratory muscle training vs. aerobic training in patients with chronic heart failure. *Eur J Heart Fail.* 2014;16:574-582.
19. Moreno AM, Toledo-Arruda AC, Lima JS, Duarte CS, Villacorta H, Nobrega ACL. Inspiratory Muscle Training Improves Intercostal and Forearm Muscle Oxygenation in Patients With Chronic Heart Failure: Evidence of the Origin of the Respiratory Metaboreflex. *J Card Fail.* 2017;23(9):672-679.
20. Hossein Pour AH, Gholami M, Saki M, Birjandi M. The effect of inspiratory muscle training on fatigue and dyspnea in patients with heart failure: A randomized, controlled trial. *Jpn J Nurs Sci.* 2019;17(2):12290.
21. Mello PR, Guerra GM, Borile S, Rondon MU, Alves MJ, Negrao CE, et al. Inspiratory Muscle Training Reduces Sympathetic Nervous Activity and Improves Inspiratory Muscle Weakness and Quality of Life in Patients With Chronic Heart Failure: A Clinical Trial. *J Cardiopulm Rehabil Prev.* 2012;32(5):255-61.
22. Hornikx M, Buys R, Cornelissen V, Deroma M, Goetschalckx K. Effectiveness of high intensity interval training supplemented with peripheral and inspiratory resistance training in chronic heart failure: a pilot study. *Acta Cardiol.* 2020;75(4):339-347.
23. Chen PC, Liaw MY, Wang LY, Tsai YC, Hsin YJ, Chen YC, et al. Inspiratory muscle training in stroke patients with congestive heart failure: A CONSORT-compliant Prospective Randomized Single-Blind Controlled Trial. *Medicine.* 2016;95(37):4856.
24. Kawauchi TS, Umeda IIK, Braga LM, Mansur AP, Rossi-Neto JM, de Moraes Rego Sousa AG, et al. Is there any benefit using low-intensity inspiratory and peripheral muscle training in heart failure? A randomized clinical trial. *Clin Res Cardiol.* 2017;106(9):676-685.
25. Antunes-Correa LM, Trevizan PF, Bacurau AVN, Ferreira-Santos L, Gomes JLP, Urias U, et al. Effects of aerobic and inspiratory training on skeletal muscle microRNA-1 and downstream-associated pathways in patients with heart failure. *J Cachexia Sarcopenia Muscle.* 2020;11(1):89-102.
26. Palau P, Dominquez E, Nunez E, Schmid JP, Vergara P, Ramon MJ, et al. Effects of inspiratory muscle training in patients with heart failure with preserved ejection fraction. *Eur J Prev Cardiol.* 2014;21(12):1465-73.

27. Palau P, Dominguez E, Lopez L, Ramon JM, Heredia R, Gonzalez J, et al. Inspiratory Muscle Training and Functional Electrical Stimulation for Treatment of Heart Failure With Preserved Ejection Fraction: The TRAINING-HF Trial. *Rev Esp Cardiol (Engl Ed)*. 2019;72(4):288-297.
28. Ekman I, Kjellstrom B, Falk K, Norman J, Swedberg K. Impact of device-guided slow breathing on symptoms of chronic heart failure: a randomized, controlled feasibility study. *Eur J Heart Fail*. 2011;13(9):1000-5.
29. Drozd T, Bilo G, Debicka-Dabrowska D, Klocek M, Malfatto G, Kielbasa G, et al. Blood pressure changes in patients with chronic heart failure undergoing slow breathing training. *Blood Press*. 2016;25(1):4-10.
30. Seo Y, Yates B, LaFramboise L, Pozehl B, Norman JF, Hertzog M. A Home-Based Diaphragmatic Breathing Retraining in Rural Patients With Heart Failure. *West J Nurs Res*. 2016;38(3):270-91.
31. Kawecka-Jaszcz K, Bilo G, Drozd T, Debicka-Dabrowska D, Kielbasa G, Malfatto G, et al. Effects of device-guided slow breathing training on exercise capacity, cardiac function, and respiratory patterns during sleep in male and female patients with chronic heart failure. *Pol Arch Intern Med*. 2017;127(1):8-15.
32. Lachowska K, Bellwon J, Narkiewicz K, Gruchala M, Hering D. Long-term effects of device-guided slow breathing in stable heart failure patients with reduced ejection fraction. *Clin Res Cardiol*. 2019;108(1):48-60.
33. Tikunov B, Levine S, Mancini D. Chronic Congestive Heart Failure Elicits Adaptions of Endurance Exercise in Diaphragmatic muscle. *Circulation*. 1997;95(4):910-6.
34. Chiappa GR, Roseguini BT, Vieira PJC, Alves CN, Tavares A, Winkelman ER, et al. Inspiratory Muscle Training Improves Blood Flow to Resting and Exercising Limbs in Patients With Chronic Heart Failure. *J Am Coll Cardiol*. 2008;51(17):1663-71.
35. Lin SL, McElfresh J, Hall B, Bloom R, Farrell K. Inspiratory muscle training in patients with heart failure: a systematic review. *Cardiopulm Phys Ther J*. 2012;23(3):29-36.
36. Working Group on Cardiac Rehabilitation and Exercise Physiology and Working Group on Heart Failure of the European Society of Cardiology. Recommendations for exercise training in chronic heart failure patients. *Eur Heart J*. 2001;22:125-135.
37. Neto MG, Martinez BP, Conceicao CS, Silva PE, Carvalho VO. Combined exercise and inspiratory muscle training in patients with heart failure: a systematic review and meta-analysis. *J Cardiopulm Rehabil Prev*. 2016;36(6):395.

ANNEX


Figure 1. Flow diagram of search strategy.

TABLE 1. Methodological quality of the studies included in the systemic review (Pedro scores)

	Random allocation	Concealed allocation	Baseline comparability	Blinded subjects	Blinded therapists	Blinded assessors	Follow-up	Intention-to-treat	Between-group analysis	Point estimates and variability	Total score
Bosnak-Guclu et al, ¹⁷ 2011	√		√	√		√	√		√	√	7
Ekman et al, ²⁸ 2011	√		√				√		√	√	5
Laoutaris et al, ¹⁶ 2011	√		√			√	√		√	√	6
Mello et al, ²¹ 2012		√	√		√	√	√			√	6
Laoutaris et al, ¹⁵ 2013	√		√			√	√		√	√	6
Marco et al, ¹⁴ 2013		√	√	√	√	√	√		√	√	8
Palau et al, ²⁶ 2014	√		√			√	√		√	√	6
Adamopoulos et al, ¹⁸ 2014	√		√				√		√	√	5
Drozd et al, ²⁹ 2015	√		√				√		√	√	5
Seo et al, ³⁰ 2015	√		√				√		√	√	5
Chen et al, ²³ 2016	√		√			√	√		√	√	6
Kawauchi et al, ²⁴ 2017		√	√				√		√	√	5
Kawecka et al, ³¹ 2017		√	√		√		√		√	√	6
Moreno et al, ¹⁹ 2017	√		√				√		√	√	5
Palau et al, ²⁷ 2020		√	√			√	√	√	√	√	7
Hornikx et al, ²² 2020	√		√				√		√	√	5
Hosseini Pour et al, ²⁰ 2019		√	√	√	√	√	√		√	√	8
Lachowska et al, ³² 2019	√		√				√		√	√	5

Antunes-Correra et al, ²⁵ 2020	√		√				√		√	√	5
---	---	--	---	--	--	--	---	--	---	---	---

TABLE 2. Intervention characteristics, parameters, outcomes and main results of the studies which included in this review

Study	Sample of intervention/control group	Interventions by group/side		IMT parameters	Study Measurements				
		Control group	Experimental group		Functional assessments				
					Peripheral muscle strength	Respiratory muscle strength	Exercise and functional capacity	Pulmonary function	Quality of life, dyspnea & echocardiography parameters
Marco et al,¹⁴ 2013	22 CHF Exp= 11 Age: 68.5±8.88 yr Con= 11 Age: 70.1±10.7 5 years NYHA: II-III	Sham-IMT	IMT	4 wk IMT F: 2x/d, 7d/wk 5 set/10 rep R: 1-2 min RR:15-20 breaths/min Exp: I: 10 consecutive max inspirations (10RM)-100% of their 10RM Con: I: 10 cmH ₂ O & ↑ 2.5 cmH ₂ O/wk	Exp: ↔ in upper limb muscles Con: ↔ in upper limb muscles Between groups: ↔ in upper limb muscles	Exp: ↑MIP, ↑MEP Con: ↔ MIP, MEP Between groups: Exp ↑MIP, ↑MEP Vs Con			Exp: ↓dyspnea ↔ Qol Con: ↔ Qol, dyspnea Between groups ↔ Qol, dyspnea

<p>Laoutaris et al,¹⁵ 2013</p>	<p>27 CHF</p> <p>Exp= 13 Age: 57.1±11 yr</p> <p>Con= 14 Age: 58.6±8 yr</p> <p>NYHA: II/III</p>	<p>AT</p>	<p>ARIS (AT/RT/ IMT)</p>	<p>12 wk F: 3x/wk</p> <p>AT: bike exercise 70-80% of max HR</p> <p>T: 30 min for Exp and 45 min for Con group</p> <p>Exp: RT: I: 50% of 1RM 3set/10-12 rep (quadriceps resistance exercises) 2set/10-12 rep (upper limb exercises) T: 15 min R: 2 min</p> <p>IMT: I: 60% of SMIP T: 20 min</p>	<p>Exp: ↑QMT peak, ↑QME, ↑1RM</p> <p>Con: ↑QME ↔ QMT peak, 1RM,</p> <p>Between groups:</p> <p>Exp ↑QMT peak, ↑QME, ↑1RM Vs Con</p>	<p>Exp: ↑MIP, ↑SMIP</p> <p>Con: ↑MIP ↔ SMIP</p> <p>Between groups:</p> <p>Exp ↑SMIP, ↔ MIP Vs Con</p>	<p>Exp: ↑peak VO₂, ↑exercise time</p> <p>↑VT, ↑CP, ↓VE/VCO₂ slope</p> <p>↔ RER, VE,</p> <p>Con: ↑peak VO₂, ↑exercise time, ↑CP, ↑VT, ↔ VE, VE/VCO₂ slope, RER</p> <p>Between groups:</p> <p>Exp ↑Exercise time, ↑CP Vs Con</p> <p>↔ peak VO₂, VE/VCO₂ slope, VT, VE, RER, peak VO₂</p>		<p>Exp: ↑Qol, ↓NYHA, ↑LVEF (%), ↑LVEDD, ↑LVESD ↔ dyspnea</p> <p>Con: ↑LVEF (%), ↑LVESD ↔ Qol, LVEDD, NYHA, dyspnea</p> <p>Between groups:</p> <p>Exp ↑Qol, ↓dyspnea Vs Con</p> <p>↔ NYHA, peak VO₂, LVEF (%), LVEDD, LVESD</p>
<p>Laoutaris et al,¹⁶ 2011</p>	<p>15 CHF</p> <p>Exp= 10</p> <p>Age: 37.2±17.7 yr</p> <p>Con= 5</p> <p>Age: 41.8±14.6 yr</p>	<p>AT</p>	<p>IMT/AT</p>	<p>10 wk</p> <p>Both groups walked every day for 30-45 min</p> <p>Exp: Bike or treadmill I: moderate 12-14 of Borg scale T: 45 min F: 3-5x/wk</p> <p>IMT: I: 60% of SMIP F: 2-3x/wk</p>		<p>Exp: ↑MIP, ↑SMIP</p> <p>Con: ↔ MIP, SMIP</p> <p>Between groups:</p> <p>Exp ↔ MIP, SMIP Vs Con</p>	<p>Exp: ↑peak VO₂, ↑VO₂ at ventilatory threshold, ↑6MWT, ↓VE/VCO₂ slope</p> <p>↔ VE, RER, exercise time</p> <p>Con: ↔ peak VO₂, VE/VCO₂ slope, VO₂ at ventilatory threshold,</p>	<p>Exp: ↑IC lung, ↔ FVC (%), FEV₁ (%), FEV₁/FVC</p> <p>Con: ↔ IC lung, FVC (%), FEV₁ (%), FEV₁/FVC</p> <p>Between groups: ↔ IC lung, FVC (%), FEV₁ (%), FEV₁/FVC</p>	<p>Exp: ↑Qol, ↔ dyspnea</p> <p>Con: ↔ Qol, dyspnea</p> <p>Between groups: ↔ dyspnea, Qol</p>

							VE, RER, exercise time Between groups: ↔ 6MWT, peak VO ₂ , VO ₂ at ventilatory threshold, VE/VCO ₂ slope, VE, RER, exercise time		
Bosnak-Guclu et al,¹⁷ 2011	30 CHF Exp= 16 Age: 69.50±7.9 6 yr Con= 14 Age: 65.71±10.52 yr NYHA: II-III	Sham-IMT	IMT	6 wk Exp: I: 40% of MIP adjusted weekly to maintain 40% of MIP Con: I: 15% of MIP T:30 min/d F: 7x/wk RR: 25-30 breaths at each workload	Exp: ↑balance, ↑QFMS, ↑QFMS (%) Con: ↔ QFMS, QFMS (%), balance Between groups: Exp ↑QFMS ↑balance Vs Con	Exp: ↑MIP, ↑MIP (%), ↑MEP, ↑MEP (%) Con: ↑MIP, ↑MIP (%), ↑MEP ↔ MEP (%) Between groups: Exp ↑MIP, ↑MIP (%), ↑MEP Vs Con	Exp: ↑6MWT, 6MWT (%) Con: ↔ 6MWT Between groups: Exp ↑6MWT, 6MWT (%) Vs Con	Exp: ↑FVC (%), ↑FEV ₁ (%), ↔ FEV ₁ /FVC Con: ↑FVC (%), ↔ FEV ₁ /FVC Between groups: Exp ↑FEV ₁ /FVC Vs Con ↔ FEV ₁ (%), FVC(%)	Exp: ↓fatigue, ↑QoL, ↓dyspnea, Con: ↑QoL ↔ dyspnea, fatigue Between groups: Exp ↓dyspnea Vs Con ↔ fatigue, QoL
Adamopoulos et al,¹⁸ 2014	43 CHF Exp= 21 Age: 57.8 ± 11.7 yr Con= 22 Age: 58.3±13.2 yr NYHA: II-III	AT/SHAM-IMT	AT/IMT	12 wk F: 3x/wk AT for both groups: I: 70-80% max HR T: 45 min Exp: I: 60% of SMIP T:30 min		Exp: ↑SMIP, ↑MIP Con: ↑MIP, ↔SMIP Between groups: Exp: ↑SMIP, ↔MIP Vs Con	Exp: ↑peak VO ₂ , ↑exercise time, ↑VE, ↑RER ↔ VE/VCO ₂ slope, VT, CP Con: ↑peak VO ₂ ↔ VE/VCO ₂ slope, VT, VE, RER,	Exp: ↔ FEV ₁ , FEV ₁ /FVC, FVC Con: ↔ FEV ₁ , FEV ₁ /FVC, FVC Between groups: ↔ FEV ₁ , FEV ₁ /FVC, FVC	Exp: ↑QoL, ↑LVEF (%), ↓LVESD, ↓NYHA, ↓dyspnea ↔ LVEDD Con: ↑LVEF (%), ↓LVESD, ↓NYHA ↔ QoL, LVEDD, dyspnea Between groups:

				<p>TIRE Protocol: 6 inspiratory efforts at each level:</p> <p>Level 1-60s R Level 2-45s R Level 3-30s R Level 4-15s R Level 5-10s R Level 6-5s (to exhaustion)</p> <p>Con I: 10% of SMIP T: 30 min</p>			<p>exercise time, CP</p> <p>Between groups: ↔ VE/VCO₂ slope, VT, VE, RER, exercise time, CP, peak VO₂</p>		<p>Exp ↑QoL, ↓Dyspnea Vs Con ↔ LVEDD, LVESD, LVEF (%), NYHA</p>
Moreno et al,¹⁹ 2017	<p>26 CHF</p> <p>Exp= 13 Age: 61±14 yr</p> <p>Con= 13 Age: 60±13 yr</p> <p>NYHA: II-III</p>	Sham -IMT	IMT	<p>8 wk</p> <p>F: 6x/wk T: 30min/d</p> <p>Exp: I: 30% of MIP RR: 15 breaths/min</p> <p>Con: I: 2% of MIP IMMP: IRT set at 60% of MIP for 1 min after ↑ to 70%, 80% and 90% of MIP until fatigue</p>		<p>Exp: ↑MIP</p> <p>Con: ↔ MIP</p> <p>Between groups: IMT ↑ MIP Vs Con</p>	<p>Exp: ↑VO₂</p> <p>Con: ↑VO₂</p> <p>Between groups: ↔ VO₂</p>		<p>Exp: ↑QoL</p> <p>Con: ↔ QoL</p> <p>Between groups: Exp ↑QoL Vs Con</p>
Hossein Pour et al,²⁰ 2019	<p>84 CHF</p> <p>Exp= 42 Age: 55.97±9.4 3 yr</p> <p>Con= 42 Age: 57.28±9.0 6 yr</p> <p>NYHA: II-III/IV</p>	Sham -IMT	IMT	<p>6 wk</p> <p>F: 1x/d, 7d/wk T: 30 min, 3 min sets of training R: 1 min/set</p> <p>Exp: I: 40% of MIP</p>					<p>Exp: ↓NYHA, ↓dyspnea</p> <p>Con: ↑dyspnea, ↔ NYHA</p> <p>Between groups: Exp ↓dyspnea, fatigue, NYHA Vs Con</p>

				Training load adjusted to maintain 40% of MIP weekly Con: I: 10% of MIP					
Mello et al,²¹ 2012	27 CHF Exp= 15 Age: 54.3 ±2 yr Con= 12 Age: 53.3 ±2 yr	No intervention	IMT	12 wk F: 3x/d, 7x/wk T: 10 min Exp: I: 30% of MIP Con: no inspiratory load	Exp: ↑MSNA Con:↔ MSNA Between groups: Exp ↔ MSNA Vs Con	Exp: ↑MIP Con ↔ MIP Between groups: Exp ↑MIP Vs Con	Exp: ↑peak VO ₂ , ↓VE/VCO ₂ peak ↓VE/VCO ₂ slope Con ↔ peak VO ₂ , VE/VCO ₂ slope, VE/VCO ₂ peak Between groups: Exp ↑peak VO ₂ , ↓VE/VCO ₂ peak, ↓VE/VCO ₂ slope Vs Con		Exp: ↑Qol Con: ↔ Qol Between groups: Exp ↑Qol Vs Con
Hornikx et al,²² 2020	20 CHF Exp=10 Age: 64±8 yr Con=10 Age: 58±11 yr	AT/RT	RHIIT (RT/HIT/IRT)	12 wk F: 3x/wk Con: I: 50% Wpeak (3 min warm up, 2x7 min cycling, 2x7 min walking in treadmill, 12 min: row- ing, step and armergometry) RT: callis-	Exp: ↑QFMS Con: ↔QFMS Between groups: Exp ↑QFMS Vs Con	Exp: ↑MIP Con: ↔ MIP Between groups: Exp ↑MIP Vs Con	Exp: ↑peak VO ₂ , ↔ VE/VCO ₂ Con: ↑VO ₂ peak, ↔ VE/VCO ₂ Between groups: ↔ peak VO ₂ , VE/VCO ₂		Exp: ↑Qol Con: ↔ Qol Between groups: ↔ Qol

				<p>thenics exercises (20 min)</p> <p>Exp: HIIT: cycling I: 80% of W_{peak} T: 33 min RT: I: 65 % of 1RM (2 set/10 rep on a leg press) IRT: F: 2x/d I:50% of MIP RR: breath in & out 30 times</p>					
Chen et al,²³ 2016	<p>21 CHF</p> <p>Exp= 11 Age: 63.73 ±14.64 yr</p> <p>Con= 10 Age: 67.50±10.35 yr</p>	Stroke rehabilitation program	IMT	<p>10 wk F: 1x/d, at least 5x/wk T:30 min</p> <p>Exp: I: 30%of MIP with ↑ 2cmH₂O each wk Exp+ Con: participated in a conventional stroke rehabilitation program</p> <p>Con: did not receive any IMT</p>		<p>Exp: ↑MIP, ↔ MEP</p> <p>Con: ↔ MIP, MEP</p> <p>Between groups:</p> <p>Exp ↑MIP, ↔ MEP Vs Con</p>		<p>Exp: ↑FVC, ↑FEV₁, ↔ FEV₁/FVC</p> <p>Con: ↔ FEV₁, FEV₁/FVC</p> <p>Between groups: ↔ FEV₁, FVC, FEV₁/FVC</p>	
Kawauchi et al,²⁴ 2017	<p>35 CHF</p> <p>Exp1= 13 Age: 54±10 yr</p> <p>Exp2= 13 Age: 56±7 yr</p>	No intervention	<p>Exp₁= LIPRT (IMT/RT)</p> <p>Exp₂= MIPRT</p>	<p>8 wk F: 7d/wk LIPRT: IMT: I: 15% of MIP</p> <p>RT: 0.5 kg (upper & lower limbs)</p>	<p>Exp₁: ↑Quadri-ceps strength</p> <p>Exp_{1,2}: ↑Quadri-ceps strength</p>	<p>Exp₁: ↑MEP, ↑MIP</p> <p>Exp₂: ↑MIP, ↑MEP</p> <p>Con: ↔ MIP, MEP</p>	<p>Exp₁: ↑6MWT</p> <p>Exp₂: ↑6MWT</p> <p>Con: ↔ 6MWT</p>	<p>Within & Between groups: ↔ FEV₁ (%), FVC (%), FEV₁/FVC</p>	<p>Exp₁: ↑QoL, ↔ NYHA</p> <p>Exp₂: ↑QoL, ↓NYHA</p> <p>Con: ↑QoL, ↔NYHA</p>

	Con= 9 Age: 56±7 yr NYHA: II/III		(IMT/RT)	exercises), 10 rep/exercise during first 2 wk, 2set/10 rep for the remaining 6 wk MIPRT: IMT: I: 30% of MIP RT: 50% of 1RM	(1RM) Vs Con	Between groups: Exp ₁ ↑MIP, ↔ MEP Vs Con Exp ₂ ↑MIP, ↑MEP Vs Con	Between groups: Exp _{1,2} ↑6MWT Vs Con		Between groups: ↔ QoI, NYHA
Antunes-Correa et al,²⁵ 2020	33 CHF Exp1= 11 Age: 55±3 yr Exp2= 12 Age: 57±2 yr Con= 10 Age: 57±3 yr NYHA: II-III	No intervention	Exp ₁ = IMT Exp ₂ = AT	4 mo Exp ₁ : I: 60% of MIP F: 5x/wk T: 30 min RR: 15-20 breaths/min Exp ₂ : F: 3x/week Each session included: 5 min stretching exercises, 40 min of cycling, 10 min strengthening exercises R: 5 min		Exp ₁ : ↑MIP Exp ₂ : ↔ MIP Con: ↔ MIP Between groups: Exp ₁ ↑MIP Vs Con Exp ₂ ↔ MIP Vs Con	Exp ₁ : ↑peak VO ₂ , ↑peak workload Exp ₂ : ↑peak VO ₂ , ↑peak workload Con: ↑peak workload, ↔ peak VO ₂ Between groups: Exp _{1,2} ↑VO ₂ peak Vs Con Exp ₂ ↑peak workload Vs Con Exp ₁ ↔ peak workload Vs Con		Exp ₁ : ↑QoI Exp ₂ : ↑QoI Con: ↑QoI Between groups: Exp _{1,2} ↑QoI Vs Con
Palau et al,²⁶ 2014	26 CHF (HFpEF) Exp= 14Age: 68 (60-76) y	Usual care	IMT +Usual care	12 wk F: 2x/d T: 20 min Exp: I: 25-30% of MIP		Exp: ↑ MIP Between: Exp: ↑MIP Vs Con	Exp: ↑peak VO ₂ , ↑VO ₂ AT, ↓VE/VCO ₂ slope, ↑RER, ↑6MWT		Exp: ↑QoI, ↔ LVEF (%) Con ↔ LVEF (%), QoI Between groups: Exp

	Con= 12Age: 74 (73–77) yr NYHA: III/IV						Con ↔ VE/VCO ₂ slope, VO ₂ AT, ↓peak VO ₂ , 6MWT Between groups: Exp ↑peak VO ₂ , ↑VO ₂ AT, VE/VCO ₂ slope ↑ 6MWT, ↑RER Vs Con	↑QoL Vs Con, ↔ LVEF (%)
Palau et al,²⁷ 2019	59 (HFpEF) Con= 13 Exp1= 15 Exp2= 15 Exp3= 16 Age: 74±9 yr NYHA: II- III/IV	Usual care	Exp ₁ : IMT (home based) Exp ₂ : FES Exp ₃ : IMT +FES	12 wk & 6 mo follow up Exp ₁ : F: 2x/d T: 20 min/session I: 25% to 30% of their MIP Exp ₂ : FES program for both legs T: 45 min F: 2d/wk for a total of 12 wk Stimulator: F: 10 to 50Hz, for 5s R: 5s Exp ₃ : re- ceived IMT and FES training at the same time			Exp ₁ : ↑peak VO ₂ (3 & 6 mo) ↓VE/VCO ₂ slope (6 mo) ↑6MWT (3 & 6 mo), ↔ VE/VCO ₂ slope (3 mo) Exp ₂ : ↑peak VO ₂ (3 & 6 mo) ↑6MWT (3 & 6 mo), ↔ VE/VCO ₂ slope (3 & 6 mo) Exp ₃ : ↑peak VO ₂ (3 & 6 mo) ↓VE/VCO ₂ slope (3 mo), ↑6MWT (3 & 6 mo), ↔ VE/VCO ₂	Exp ₁ : ↑QoL (3 & 6 mo) Exp ₂ : ↑QoL (3 mo), ↔ QoL (6 mo) Exp ₃ : ↑QoL (3 & 6 mo) Between groups: Exp _{1,2,3} groups ↑ QoL (3 mo) Vs Con ↔ LVEF (%), LVEDD, LVESD

							<p>slope (6 mo)</p> <p>Between groups: Exp_{1,2,3} groups ↑peak VO₂ (3 & 6 mo) Vs Con</p> <p>Exp_{1,2,3} groups ↑6MWT (3 mo) Vs Con, at 6 mo this beneficial effects persisted for Exp_{2,3}</p> <p>↔ peak VO₂, VE/VCO₂ slope between Exp_{1,2,3} groups</p>	
Ekman et al,²⁸ 2011	<p>65 CHF</p> <p>Exp= 30</p> <p>Con= 35</p> <p>Age: 73±11 yr</p> <p>NYHA: II-IV</p>	Music listening	DGB	<p>4 wk</p> <p>Exp: DGB exercises F: 2x/d T: 20 min/session RR: <10 breaths/min</p> <p>Con: listened music</p>				<p>Exp: ↓dyspnea, ↓NYHA Con: ↔ dyspnea, NYHA Responders in Exp (an average increase in Tex/Tin of >0.2 and a reduction in the average respiration rate): ↓breathlessness, ↓NYHA compared with no-responders or controls</p> <p>Between groups ↔</p>

									dyspnea, NYHA
Drozd et al,²⁹ 2016	40 CHF Exp1= 20 Exp2= 20 Age: 63.3 ± 13.4 yr NYHA: II-III	Usual care	DGB	Exp ₁ : started with 10-12 wk SBT followed by 10-12 wk usual care Exp ₂ : started with 10-12 wk usual care followed by 10-12 wk SBT SBT: F: 2x/d T: 15 min/session RR: 6 breaths/min					Exp1: ↑QoL Exp2: ↔ QoL Between groups ↔ QoL
Seo et al,³⁰ 2015	36 CHF Exp=18 Age: 65.2±11.3 4 yr Con= 18 Age: 66.6±13.6 9 yr NYHA: II-IV	Attention group	DBR	8 wk & 5 mo follow up Exp: received 3 audio CDs (1 each for wk 1, wk 2, & wk 3-8) RR: 6 breaths/min F: 2x/day at least 5 d/wk Week 1 goal: 5 min of CD-guided DBR Week 2 goal: 10 min of CD-guided DBR Weeks 3-8 goal: 15 min of CD-guided DBR + 4 telephone calls (feedback & encourage)			Exp: ↑6MWT (8 wk & sustained at 5 mo follow up) Con: ↑6MWT (8 wk) Between groups: ↔ 6MWT (8 weeks & 5 mo)		Exp & Con: ↓dyspnea (8 wk & 5 mo) Between groups: ↔ dyspnea (8 weeks & 5 mo)

				Con: received 4 telephone calls with general health topics					
Kawecka-Jaszcz et al,³¹ 2017	96 CHF Age: 64.5 (57.0–71.5) yr NYHA: I-III	Usual care	DGB	Exp ₁ : started with 10-12 wk SBT followed by 10-12 wk usual care Exp ₂ : started with 10-12 wk usual care followed by 10-12 wk SBT SBT: F: 2 x/day T: 15 min/session RR: 6 breaths/min			After SBT: ↑6MWT		After SBT: ↑EF, ↑LVEF (%), ↓LVEDD
Lachowska et al,³² 2019	21 HFrEF Exp= 11 Con= 10 Age: 52±17 yr NYHA: I-III		DGB	12 wk & 6 mo follow up SBT: F: 2x/d T: 15 min/session (totally 30 min) RR: 10 breaths/ min At 3 mo follow up- Group ₁ : continue SBT Group ₂ : no SBT			SLOWB from baseline to 6 mo follow up: ↑6MWT & peak RER in Group 1 but no Group 2 ↔ peak VO ₂ , VE/VCO ₂ from baseline to 3 mo follow up		SLOWB: ↔ LVEF (%), LVEDD (3 & 6 mo)

Abbreviations:

Exp: experimental, **Con:** control, **IMT:** inspiratory muscle training, **SBT:** slow breathing training, **DGB:** device-guided breathing, **IMMP:** inspiratory muscle metaboreflex protocol, **LVEF:** left ventricular ejection fraction, **LVESD:** left ventricular end-systolic diameter, **LVEDD:** left ventricular end-diastolic diameter, **HR:** heart rate, **I:** intensity, **R:** rest, **F:** frequency, **T:** time, **RR:** respiratory rate, **MSNA:** muscle sympathetic nerve activity, **LIPRT:** low intensity inspiratory and peripheral resistance training, **RT:** resistance training, **IRT:** inspiratory resistance training, **HIIT:** high intensity interval training, **MIPRT:** moderate-intensity inspiratory and peripheral resistance training, **FES:** functional electrical stimulation **AT:** aerobic training, **DBR:** diaphragmatic breathing retraining, **RER:** respiratory exchange ratio, **SMIP:** sustained maximal inspiratory pressure, **MEP:** maximal expiratory pressure, **FEV₁:** forced expiratory volume in 1 s, **FVC:** forced vital capacity, **QFMS:** quadriceps femoris muscle strength, **QMT:** quadriceps muscle torque, **QME:** quadriceps muscle endurance, **CP:** circulatory power, **VT:** ventilatory threshold, **VE:** minute ventilation, **TG:** training group, **IC:** inspiratory capacity, **EF:** ejection fraction, **peak VO₂:** peak oxygen consumption, **VE/VCO₂:** ventilation/carbon dioxide, **6MWT:** 6 min walk test, **1RM:** 1 repetition maximum, **Rep:** repetition, **NYHA:** New York Heart Association, **QoL:** quality of life, **↔:** no changes, **↑:** increased, **↓:** reduced, **%:** predicted