



Cochrane
Library

Cochrane Database of Systematic Reviews

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

Ammous O, Feki W, Lotfi T, Khamis AM, Gosselink R, Rebai A, Kammoun S

Ammous O, Feki W, Lotfi T, Khamis AM, Gosselink R, Rebai A, Kammoun S.
Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD).

Cochrane Database of Systematic Reviews 2023, Issue 1. Art. No.: CD013778.

DOI: [10.1002/14651858.CD013778.pub2](https://doi.org/10.1002/14651858.CD013778.pub2).

www.cochranelibrary.com

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

WILEY

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	3
SUMMARY OF FINDINGS	5
BACKGROUND	10
OBJECTIVES	11
METHODS	11
Figure 1.	14
RESULTS	17
DISCUSSION	24
Figure 2.	26
Figure 3.	27
Figure 4.	28
Figure 5.	29
AUTHORS' CONCLUSIONS	30
ACKNOWLEDGEMENTS	30
REFERENCES	32
CHARACTERISTICS OF STUDIES	50
RISK OF BIAS	164
DATA AND ANALYSES	193
Analysis 1.1. Comparison 1: PR+IMT vs PR, Outcome 1: Dyspnea: Borg (at submaximal exercise: 50% to 80% of Wmax)	195
Analysis 1.2. Comparison 1: PR+IMT vs PR, Outcome 2: Dyspnea: Modified Medical Research Council (mMRC)	196
Analysis 1.3. Comparison 1: PR+IMT vs PR, Outcome 3: Functional exercise capacity: 6-minute walk distance (6MWD) (meters) ..	196
Analysis 1.4. Comparison 1: PR+IMT vs PR, Outcome 4: Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: duration of the intervention)	197
Analysis 1.5. Comparison 1: PR+IMT vs PR, Outcome 5: Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: with or without respiratory muscle weakness)	198
Analysis 1.6. Comparison 1: PR+IMT vs PR, Outcome 6: Functional exercise capacity: 12-minute walk distance (12MWD) (meters)	198
Analysis 1.7. Comparison 1: PR+IMT vs PR, Outcome 7: Functional exercise capacity: Wmax (watt)	199
Analysis 1.8. Comparison 1: PR+IMT vs PR, Outcome 8: Functional exercise capacity: exercise time (seconds)	199
Analysis 1.9. Comparison 1: PR+IMT vs PR, Outcome 9: Health-related quality of life (HRQoL): St George's Respiratory Questionnaire (SGRQ)	200
Analysis 1.10. Comparison 1: PR+IMT vs PR, Outcome 10: Health-related quality of life (HRQoL): Chronic Respiratory Disease Questionnaire (CRQ)	201
Analysis 1.11. Comparison 1: PR+IMT vs PR, Outcome 11: Health-related quality of life (HRQoL): COPD Assessment Test (CAT) ..	202
Analysis 1.12. Comparison 1: PR+IMT vs PR, Outcome 12: Inspiratory muscle strength: P _I max (cmH ₂ O)	202
Analysis 1.13. Comparison 1: PR+IMT vs PR, Outcome 13: Inspiratory muscle strength: P _I max (cmH ₂ O) (subgroup analysis: duration of the intervention)	203
Analysis 1.14. Comparison 1: PR+IMT vs PR, Outcome 14: Inspiratory muscle strength: P _I max (cmH ₂ O) (subgroup analysis: with or without respiratory muscle weakness)	204
Analysis 1.15. Comparison 1: PR+IMT vs PR, Outcome 15: Laboratory exercise test: VO ₂ peak (L/min)	204
Analysis 1.16. Comparison 1: PR+IMT vs PR, Outcome 16: Respiratory muscle endurance: respiratory muscle endurance pressure (P _{th} max) (cmH ₂ O)	205
Analysis 1.17. Comparison 1: PR+IMT vs PR, Outcome 17: Respiratory muscle endurance time: T _{lim} (seconds) (sustained ventilation according to P _I max)	205
Analysis 1.18. Comparison 1: PR+IMT vs PR, Outcome 18: Respiratory muscle endurance time: T _{lim} (seconds) (sustained ventilation according to MVV)	205
Analysis 1.19. Comparison 1: PR+IMT vs PR, Outcome 19: Maximal voluntary ventilation (MVV)	206
Analysis 1.20. Comparison 1: PR+IMT vs PR, Outcome 20: Respiratory function: forced expiratory volume at 1 second (FEV ₁) (%Pred)	206
Analysis 1.21. Comparison 1: PR+IMT vs PR, Outcome 21: Respiratory function: forced expiratory volume at 1 second (FEV ₁) (Liters)	206
Analysis 2.1. Comparison 2: IMT vs control/sham, Outcome 1: Dyspnea: Borg (at submaximal exercise capacity)	210

Analysis 2.2. Comparison 2: IMT vs control/sham, Outcome 2: Dyspnea: Baseline and Transition Dyspnea Indexes (BDI-TDI) ...	211
Analysis 2.3. Comparison 2: IMT vs control/sham, Outcome 3: Dyspnea: Transition Dyspnea Index (TDI): Focal score (subgroup analysis: with or without respiratory muscle weakness)	212
Analysis 2.4. Comparison 2: IMT vs control/sham, Outcome 4: Dyspnea: Modified Medical Research Council (mMRC)	212
Analysis 2.5. Comparison 2: IMT vs control/sham, Outcome 5: Functional exercise capacity: 6-minute walk distance (6MWD) (meters)	213
Analysis 2.6. Comparison 2: IMT vs control/sham, Outcome 6: Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: duration of the intervention)	214
Analysis 2.7. Comparison 2: IMT vs control/sham, Outcome 7: Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: with or without respiratory muscle weakness)	215
Analysis 2.8. Comparison 2: IMT vs control/sham, Outcome 8: Functional exercise capacity: 12-minute walk distance (12MWD) (meters)	215
Analysis 2.9. Comparison 2: IMT vs control/sham, Outcome 9: Functional exercise capacity: Wmax (watt)	216
Analysis 2.10. Comparison 2: IMT vs control/sham, Outcome 10: Functional exercise capacity: exercise time (seconds)	216
Analysis 2.11. Comparison 2: IMT vs control/sham, Outcome 11: Functional exercise capacity: shuttle walk test (SWT) (meters)	217
Analysis 2.12. Comparison 2: IMT vs control/sham, Outcome 12: Health-related quality of life (HRQoL): St George Respiratory Questionnaire (SGRQ)	217
Analysis 2.13. Comparison 2: IMT vs control/sham, Outcome 13: Health-related quality of life (HRQoL): Chronic Respiratory Disease Questionnaire (CRQ)	218
Analysis 2.14. Comparison 2: IMT vs control/sham, Outcome 14: Health-related quality of life (HRQoL): COPD Assessment Test (CAT)	219
Analysis 2.15. Comparison 2: IMT vs control/sham, Outcome 15: Inspiratory muscle strength: P _{lmax} (cmH ₂ O)	220
Analysis 2.16. Comparison 2: IMT vs control/sham, Outcome 16: Inspiratory muscle strength: P _{lmax} (cmH ₂ O) (subgroup analysis: duration of the intervention)	221
Analysis 2.17. Comparison 2: IMT vs control/sham, Outcome 17: Inspiratory muscle strength: P _{lmax} (cmH ₂ O) (subgroup analysis: with or without respiratory muscle weakness)	222
Analysis 2.18. Comparison 2: IMT vs control/sham, Outcome 18: Inspiratory muscle strength: P _{lmax} (cmH ₂ O) (subgroup analysis: method of measurement)	223
Analysis 2.19. Comparison 2: IMT vs control/sham, Outcome 19: Laboratory exercise test: VO _{2peak}	224
Analysis 2.20. Comparison 2: IMT vs control/sham, Outcome 20: Respiratory muscle endurance: respiratory muscle endurance pressure (P _{thmax}) (cmH ₂ O)	224
Analysis 2.21. Comparison 2: IMT vs control/sham, Outcome 21: Respiratory muscle endurance time: T _{lim} (seconds)	225
Analysis 2.22. Comparison 2: IMT vs control/sham, Outcome 22: Maximal voluntary ventilation (MVV)	225
Analysis 2.23. Comparison 2: IMT vs control/sham, Outcome 23: Respiratory function: forced expiratory volume at 1 second (FEV ₁) (%pred)	226
Analysis 2.24. Comparison 2: IMT vs control/sham, Outcome 24: Respiratory function: forced expiratory volume at 1 second (FEV ₁) (Liters)	226
APPENDICES	226
WHAT'S NEW	231
HISTORY	231
CONTRIBUTIONS OF AUTHORS	231
DECLARATIONS OF INTEREST	231
SOURCES OF SUPPORT	232
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	232
INDEX TERMS	232

[Intervention Review]

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD)

Omar Ammous¹, Walid Feki², Tamara Lotfi³, Assem M Khamis⁴, Rik Gosselink⁵, Ahmed Rebai⁶, Samy Kammoun²

¹Faculty of Medicine, University of Sfax, Sfax, Tunisia. ²Department of Respiratory Medicine, Hedi Chaker University Hospital, University of Sfax, Sfax, Tunisia. ³Department of Health Research Methods, Evidence and Impact, McMaster University, Hamilton, Canada. ⁴Hull York Medical School, University of Hull, Hull, UK. ⁵Department of Rehabilitation Sciences, Faculty of Movement and Rehabilitation Sciences, University Hospitals Leuven, Leuven, Belgium. ⁶Centre of Biotechnology of Sfax, University of Sfax, Sfax, Tunisia

Contact: Omar Ammous, omar.ammous@outlook.com.**Editorial group:** Cochrane Airways Group.**Publication status and date:** Edited (no change to conclusions), published in Issue 1, 2023.**Citation:** Ammous O, Feki W, Lotfi T, Khamis AM, Gosselink R, Rebai A, Kammoun S. Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD). *Cochrane Database of Systematic Reviews* 2023, Issue 1. Art. No.: CD013778. DOI: [10.1002/14651858.CD013778.pub2](https://doi.org/10.1002/14651858.CD013778.pub2).

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration. This is an open access article under the terms of the [Creative Commons Attribution-Non-Commercial-No-Derivatives Licence](https://creativecommons.org/licenses/by-nc-nd/4.0/), which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

ABSTRACT

Background

Inspiratory muscle training (IMT) aims to improve respiratory muscle strength and endurance. Clinical trials used various training protocols, devices and respiratory measurements to check the effectiveness of this intervention. The current guidelines reported a possible advantage of IMT, particularly in people with respiratory muscle weakness. However, it remains unclear to what extent IMT is clinically beneficial, especially when associated with pulmonary rehabilitation (PR).

Objectives

To assess the effect of inspiratory muscle training (IMT) on chronic obstructive pulmonary disease (COPD), as a stand-alone intervention and when combined with pulmonary rehabilitation (PR).

Search methods

We searched the Cochrane Airways trials register, CENTRAL, MEDLINE, Embase, PsycINFO, Cumulative Index to Nursing and Allied Health Literature (CINAHL) EBSCO, Physiotherapy Evidence Database (PEDro) ClinicalTrials.gov, and the World Health Organization International Clinical Trials Registry Platform on 20 October 2022. We also checked reference lists of all primary studies and review articles.

Selection criteria

We included randomized controlled trials (RCTs) that compared IMT in combination with PR versus PR alone and IMT versus control/sham. We included different types of IMT irrespective of the mode of delivery. We excluded trials that used resistive devices without controlling the breathing pattern or a training load of less than 30% of maximal inspiratory pressure (P_{Imax}), or both.

Data collection and analysis

We used standard methods recommended by Cochrane including assessment of risk of bias with RoB 2. Our primary outcomes were dyspnea, functional exercise capacity and health-related quality of life.

Main results

We included 55 RCTs in this review. Both IMT and PR protocols varied significantly across the trials, especially in training duration, loads, devices, number/ frequency of sessions and the PR programs. Only eight trials were at low risk of bias.

PR+IMT versus PR

We included 22 trials (1446 participants) in this comparison. Based on a minimal clinically important difference (MCID) of -1 unit, we did not find an improvement in dyspnea assessed with the **Borg** scale at submaximal exercise capacity (mean difference (MD) 0.19, 95% confidence interval (CI) -0.42 to 0.79; 2 RCTs, 202 participants; moderate-certainty evidence).

We also found no improvement in dyspnea assessed with the **modified Medical Research Council dyspnea scale (mMRC)** according to an MCID between -0.5 and -1 unit (MD -0.12, 95% CI -0.39 to 0.14; 2 RCTs, 204 participants; very low-certainty evidence).

Pooling evidence for the **6-minute walk distance (6MWD)** showed an increase of 5.95 meters (95% CI -5.73 to 17.63; 12 RCTs, 1199 participants; very low-certainty evidence) and failed to reach the MCID of 26 meters. In subgroup analysis, we divided the RCTs according to the training duration and mean baseline P_{lmax}. The test for subgroup differences was not significant. Trials at low risk of bias (n = 3) demonstrated a larger effect estimate than the overall.

The summary effect of the **St George's Respiratory Questionnaire (SGRQ)** revealed an overall total score below the MCID of 4 units (MD 0.13, 95% CI -0.93 to 1.20; 7 RCTs, 908 participants; low-certainty evidence).

The summary effect of **COPD Assessment Test (CAT)** did not show an improvement in the HRQoL (MD 0.13, 95% CI -0.80 to 1.06; 2 RCTs, 657 participants; very low-certainty evidence), according to an MCID of -1.6 units.

Pooling the RCTs that reported **P_{lmax}** showed an increase of 11.46 cmH₂O (95% CI 7.42 to 15.50; 17 RCTs, 1329 participants; moderate-certainty evidence) but failed to reach the MCID of 17.2 cmH₂O. In subgroup analysis, we did not find a difference between different training durations and between studies judged with and without respiratory muscle weakness.

One abstract reported some **adverse effects** that were considered "minor and self-limited".

IMT versus control/sham

Thirty-seven RCTs with 1021 participants contributed to our second comparison. There was a trend towards an improvement when **Borg** was calculated at submaximal exercise capacity (MD -0.94, 95% CI -1.36 to -0.51; 6 RCTs, 144 participants; very low-certainty evidence). Only one trial was at a low risk of bias.

Eight studies (nine arms) used the **Baseline Dyspnea Index - Transition Dyspnea Index (BDI-TDI)**. Based on an MCID of +1 unit, they showed an improvement only with the 'total score' of the TDI (MD 2.98, 95% CI 2.07 to 3.89; 8 RCTs, 238 participants; very low-certainty evidence). We did not find a difference between studies classified as with and without respiratory muscle weakness. Only one trial was at low risk of bias.

Four studies reported the **mMRC**, revealing a possible improvement in dyspnea in the IMT group (MD -0.59, 95% CI -0.76 to -0.43; 4 RCTs, 150 participants; low-certainty evidence). Two trials were at low risk of bias.

Compared to control/sham, the MD in the **6MWD** following IMT was 35.71 (95% CI 25.68 to 45.74; 16 RCTs, 501 participants; moderate-certainty evidence). Two studies were at low risk of bias. In subgroup analysis, we did not find a difference between different training durations and between studies judged with and without respiratory muscle weakness.

Six studies reported the **SGRQ** total score, showing a larger effect in the IMT group (MD -3.85, 95% CI -8.18 to 0.48; 6 RCTs, 182 participants; very low-certainty evidence). The lower limit of the 95% CI exceeded the MCID of -4 units. Only one study was at low risk of bias.

There was an improvement in life quality with **CAT** (MD -2.97, 95% CI -3.85 to -2.10; 2 RCTs, 86 participants; moderate-certainty evidence). One trial was at low risk of bias.

Thirty-two RCTs reported **P_{lmax}**, showing an improvement without reaching the MCID (MD 14.57 cmH₂O, 95% CI 9.85 to 19.29; 32 RCTs, 916 participants; low-certainty evidence). In subgroup analysis, we did not find a difference between different training durations and between studies judged with and without respiratory muscle weakness.

None of the included RCTs reported **adverse events**.

Authors' conclusions

IMT may not improve dyspnea, functional exercise capacity and life quality when associated with PR. However, IMT is likely to improve these outcomes when provided alone.

For both interventions, a larger effect in participants with respiratory muscle weakness and with longer training durations is still to be confirmed.

PLAIN LANGUAGE SUMMARY

Are exercises for strengthening breathing muscles effective for people with chronic obstructive pulmonary disease?

Key messages

- Exercise combined with specific exercises to strengthen breathing muscles may not improve breathlessness, physical fitness and life quality. Strength of breathing muscles and endurance increased but not enough to make a difference to patients.
- Specific exercises to strengthen breathing muscles compared to no exercise may improve breathlessness, physical fitness and life quality. Strength of breathing muscles and endurance increased, but we don't know if this benefitted patients.
- We don't know whether exercise or specific exercises to strengthen breathing muscles is better for people with weakened breathing muscles who trained for several weeks.
- Future research should focus on people with weakened breathing muscles and studies should include more people.

What is chronic obstructive pulmonary disease (COPD)?

Chronic obstructive pulmonary disease (COPD) is a lung condition characterized by blockages in the airways, which cause shortness of breath and a cough. It appears after the long-term inhalation of irritating gases like cigarette smoke and chemicals. Training and strengthening the breathing muscles is thought to improve breathing and reduce air obstruction.

What exercise treatments do people with COPD use?

Health professionals use various exercises to help improve people's COPD.

- Some people undertake a program of general exercise and education to help reduce symptoms and improve their exercise capacity and life quality.
- Other people try to improve the strength and endurance of the breathing muscles through a series of breathing exercises using specific devices. This is called 'inspiratory muscle training' (IMT). The devices add resistance to breathing to strengthen the diaphragm and the intercostal muscles between the ribs - the muscles used for breathing. People may then be able to breathe in more air with each breath and be active for longer. The devices are also used by people with healthy lungs to improve their sports performance.

What did we want to find out?

We wanted to find out if exercise combined with IMT compared to exercise alone, and IMT compared to no exercise or sham IMT has a better effect on breathlessness, physical fitness and life quality. (A sham device has no effect on breathing muscles. It allows a fair test of the real devices, because people don't know which they are using.)

We also wanted to check whether IMT was associated with any unwanted effects.

What did we do?

We searched for studies that compared

- exercise combined with IMT with exercise alone; and
- IMT with no exercise or sham IMT.

We compared and summarized the results of the studies and rated our confidence in the evidence, based on factors such as study methods and sizes.

What did we find?

Exercise plus IMT compared with exercise alone

We found 22 studies with 1446 participants, which lasted between 2 and 24 weeks. Exercise ranged from training only on a treadmill, with only a cycle, and a combination of exercises (training with a cycle and treadmill, muscle strengthening, stair climbing, and education). The duration and devices of IMT also varied across the studies.

We found that this combination:

- probably makes little to no difference to breathlessness (measured with different scales);
- has an unknown effect on physical fitness;

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

- may make little to no difference to life quality (measured with different scales);
- probably makes little to no difference to strength of breathing muscles.

IMT versus no training or sham device

We found 37 studies with 1021 participants, which lasted from 2 weeks to a year. IMT varied across the studies regarding devices, resistance, frequency and supervision.

We found out that IMT alone:

- may reduce breathlessness measured with one scale, but it is unclear if it has an effect when measured with two other scales;
- probably improves physical fitness;
- probably improves life quality when measured with one scale, but it is unclear if it has a benefit when measured with another one;
- may make little to no difference to strength of breathing muscles.

What are the limitations of the evidence?

The studies used different training durations, resistance, devices, number and frequency of sessions, and physical training programs. This makes it hard to draw firm conclusions. Overall our confidence in the conclusions is reduced because the studies were small, some participants may have been aware of which treatment they were receiving, and generally, there was some diversity in the studies.

How up to date is the evidence?

The evidence is up-to-date to 20 October 2022.

SUMMARY OF FINDINGS

Summary of findings 1. Pulmonary rehabilitation plus inspiratory muscle training compared to pulmonary rehabilitation alone for people with chronic obstructive pulmonary disease

Pulmonary rehabilitation plus inspiratory muscle training compared to pulmonary rehabilitation alone for people with chronic obstructive pulmonary disease

Patient or population: people with chronic obstructive pulmonary disease (COPD)

Setting: community

Intervention: pulmonary rehabilitation (PR) + inspiratory muscle training (IMT)

Comparison: PR

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with PR	Risk with PR +IMT				
<p>Dyspnea assessed with Borg scale at submaximal exercise capacity</p> <p>Scale from 0 to 10 (worse)</p> <p>Follow-up: range 3 months to 4 months</p>	The mean dyspnea was 4.65	The mean dyspnea was 0.19 points higher (0.42 lower to 0.79 higher)	-	202 (2 RCTs)	⊕⊕⊕⊕ Moderate ^d	The combination of PR+IMT probably results in little to no difference in dyspnea measured with Borg at submaximal exercise capacity compared to PR alone, considering an MCID of -1 unit
<p>Dyspnea assessed with mMRC</p> <p>Scale from 0 to 4 (worse)</p> <p>Follow-up: range 1 month to 2 months</p>	The mean dyspnea ranged from -0.8 to -0.33	MD 0.12 lower (0.39 lower to 0.14 higher)	-	204 (2 RCTs)	⊕⊕⊕⊕ Very low ^{a,b}	The evidence is very uncertain about the effect of the combination of PR+IMT on dyspnea measured with the mMRC compared to PR alone, considering an MCID between -0.5 and -1 unit
<p>Functional exercise capacity assessed with 6MWD</p> <p>Follow-up: range 2 weeks to 6 months</p>	The mean functional exercise capacity was 304.72 meters ^c	MD 5.95 meters higher (5.73 lower to 17.63 higher)	-	1199 (12 RCTs)	⊕⊕⊕⊕ Very low ^{d,e}	The evidence is very uncertain about the effect of the combination of PR+IMT on the 6MWD compared to PR alone, considering an MCID of 26 meters

<p>Health-related quality of life assessed with SGRQ total score</p> <p>Scale from 0 to 100 (worse) Follow-up: range 3 weeks to 6 months</p>	<p>The mean health-related quality of life was 14.9^c</p>	<p>MD 0.13 higher (0.93 lower to 1.2 higher)</p>	<p>-</p>	<p>908 (7 RCTs)</p>	<p>⊕⊕⊕⊕ Low^f</p>	<p>The combination of PR+IMT may result in little to no difference in health-related quality of life measured with the SGRQ compared to PR alone, considering an MCID of -4 units</p>
<p>Health-related quality of life assessed with CAT</p> <p>Scale from 0 to 40 (worse) Follow-up: range 3 weeks to 6 months</p>	<p>The mean health-related quality of life ranged from -3.42 to -3</p>	<p>MD 0.13 higher (0.8 lower to 1.06 higher)</p>	<p>-</p>	<p>657 (2 RCTs)</p>	<p>⊕⊕⊕⊕ Very low^{g,h}</p>	<p>The evidence is very uncertain about the effect of the combination of PR+IMT on health-related quality of life measured with the CAT compared to PR alone, considering an MCID of about -1.6 units</p>
<p>Inspiratory muscle strength assessed with PImax</p> <p>Follow-up: range 3 weeks to 6 months</p>	<p>The mean inspiratory muscle strength was 67.37 cmH₂O^c</p>	<p>MD 11.46 cmH₂O higher (7.42 higher to 15.50 higher)</p>	<p>-</p>	<p>1329 (17 RCTs)</p>	<p>⊕⊕⊕⊕ Moderate^d</p>	<p>The combination of IMT+PR probably slightly increases inspiratory muscle strength (PImax) compared to PR alone, without reaching the MCID of 17.2 cmH₂O</p>

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **CAT:** COPD [chronic obstructive pulmonary disease] Assessment Test; **IMT:** inspiratory muscle training; **MD:** mean difference; **mMRC:** modified Medical Research Council dyspnoea scale; **MCID:** minimum clinically important difference; **PImax:** Maximal Inspiratory Pressure; **PR:** pulmonary rehabilitation; **RCT:** randomized controlled trial; **SGRQ:** St George's Respiratory Questionnaire; **6MWD:** six-minute walk distance

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

See interactive version of this table: https://gdt.grade.pro.org/presentations/#/isof/isof_question_revman_web_423482112637765988.

^aDowngraded by one level for imprecision due to small sample size (rule of thumb: at least 400 participants).

^bDowngraded by two levels for risk of bias because all the trials are at high risk of bias.

^cIncluding change and endpoint scores.

- ^dDowngraded by one level for risk of bias because most of the evidence is from studies at high risk of bias and with some concern.
- ^eDowngraded by two levels for inconsistency due to considerable statistical heterogeneity (I^2 statistic), confidence intervals not overlapping, and significant variations in the direction of the effects.
- ^fDowngraded by two levels for risk of bias due to a considerable bias across the studies in the measurement of the outcome (lack of blinding) and all the trials are at high risk of bias and some concern.
- ^gDowngraded by two levels for risk of bias due to considerable bias across the studies in the measurement of the outcome (lack of blinding) and most of the evidence is from studies at high risk of bias and some concern.
- ^hDowngraded by one level for inconsistency due to considerable statistical heterogeneity and confidence intervals not overlapping.

Summary of findings 2. Inspiratory muscle training compared to control or sham for people with chronic obstructive pulmonary disease

Inspiratory muscle training compared to control or sham for people with chronic obstructive pulmonary disease

Patient or population: people with chronic obstructive pulmonary disease (COPD)

Setting: community

Intervention: inspiratory muscle training (IMT)

Comparison: control or sham

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Nº of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with control or sham	Risk with IMT				
Dyspnea assessed with Borg scale at submaximal exercise capacity Scale from 1 to 10 (worse) Follow-up: range 5 weeks to 4 months	The median dyspnea was 1.5	MD 0.94 lower (1.36 lower to 0.51 lower)	-	144 (6 RCTs)	⊕⊕⊕⊕ Very low ^{a,b}	IMT may improve dyspnea measured with Borg scale at submaximal exercise capacity compared to control/sham but the evidence is very uncertain, considering an MCID of -1 unit (only the lower limit of the 95% CI exceeded the MCID)
Dyspnea assessed with BDI-TDI: focal score (TDI) Scale from -9 to +9 (better) follow-up: range 2 months to 6 months	The median dyspnea was 1.2	MD 2.98 higher (2.07 higher to 3.89 higher)	-	238 (8 RCTs)	⊕⊕⊕⊕ Very low ^{b,c}	IMT may improve dyspnea measured with the BDI-TDI (Focal score) compared to control/sham but the evidence is very uncertain, considering an MCID of +1 unit
Dyspnea assessed with mMRC Scale from 0 to 4 (worse)	The median dyspnea was 0.62	MD 0.59 lower (0.76 lower to 0.43 lower)	-	150 (4 RCTs)	⊕⊕⊕⊕ Low ^{b,d}	IMT may improve dyspnea measured with the modified mMRC compared to con-

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Follow-up: range 8 months to 8 months						trol/sham, considering an MCID between -0.5 and -1 unit
Functional exercise capacity assessed with 6MWD Follow-up: range 2 weeks to 12 months	The mean functional exercise capacity was 298.4 meters	MD 35.71 meters higher (25.68 higher to 45.74 higher)	-	501 (16 RCTs)	⊕⊕⊕⊕ Moderate ^d	IMT probably improves functional exercise capacity measured with the 6MWD compared to control/sham, considering an MCID of 26 meters
Health-related quality of life assessed with SGRQ total score Scale from 0 to 100 (worse) Follow-up: range 2 months to 12 months	The median health-related quality of life was 23.61	MD 3.85 lower (8.18 lower to 0.48 higher)	-	182 (6 RCTs)	⊕⊕⊕⊕ Very low ^{e,f}	IMT may improve health-related quality of life measured with the SGRQ compared to control/sham but the evidence is very uncertain, considering an MCID of -4 units (only the lower limit of the 95% CI exceeded the MCID)
Health-related quality of life assessed with CAT Scale from 0 to 40 (worse) Follow-up: mean 2 months	The mean health-related quality of life ranged from -0.5 to 0.3	MD 2.97 lower (3.85 lower to 2.1 lower)	-	86 (2 RCTs)	⊕⊕⊕⊕ Moderate ^b	IMT probably improves health-related quality of life measured with CAT compared to control/sham, considering an MCID of about -1.6 units
Inspiratory muscle strength assessed with P _{lmax} Follow-up: range 2 weeks to 12 months	The mean inspiratory muscle strength was 51.23 cmH ₂ O	MD 14.57 cmH₂O higher (9.85 higher to 19.29 higher)	-	916 (32 RCTs)	⊕⊕⊕⊕ Low ^{d,g,h}	IMT may increase inspiratory muscle strength (P _{lmax}) slightly compared to control/sham considering the MCID of 17.2 cmH ₂ O

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

BDI-TDI: Baseline Dyspnea Index - Transition Dyspnea Index; **CI:** confidence interval; **CAT:** COPD Assessment Test; **IMT:** inspiratory muscle training; **MD:** mean difference; **mMRC:** modified Medical Research Council dyspnoea scale; **MCID:** minimum clinically important difference; **P_{lmax}:** Maximal Inspiratory Pressure; **PR:** pulmonary rehabilitation; **RCT:** randomized controlled trial; **SGRQ:** St George's Respiratory Questionnaire; **6MWD:** six-minute walk distance

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

See interactive version of this table: https://gdt.gradepr.org/presentations/#/isof/isof_question_revman_web_423482164894599533.

^aDowngraded by two levels for risk of bias because most of the evidence is from studies at high risk of bias and some concern, there is an issue with blinding, and high risk of bias studies show different estimates to studies at low risk of bias and some concern.

^bDowngraded by one level for imprecision due to small sample size (rule of thumb: less than 400).

^cDowngraded by two levels for risk of bias due to considerable bias across the studies in the measurement of the outcome (lack of blinding) and most of the evidence is from studies at high risk of bias and some concern.

^dDowngraded by one level for risk of bias because most of the evidence is from studies at high risk of bias and with some concern.

^eDowngraded by two levels for risk of bias because most of the evidence is from studies at high risk of bias and some concern, and high risk of bias studies show different estimate to studies at low risk of bias and some concern.

^fDowngraded by two levels for imprecision due to small sample size (rule of thumb: less than 400) and because the 95% CI includes benefit and harm.

^gWe did not downgrade inconsistency although substantial statistical heterogeneity because the studies are on one side of the line of no effect. So we are more confident about the direction of the effect.

^hDowngraded by one level for publication bias because the funnel plot and the number of studies give rise to serious suspicions about publication bias.

BACKGROUND

Description of the condition

Chronic obstructive pulmonary disease (COPD) is a respiratory condition that includes bronchitis and emphysema. Chronic bronchitis is defined by the presence of a productive cough for at least three months per year for two consecutive years, during which other causes of cough have been excluded (GOLD 2022). Emphysema is damage to the portion of the lungs responsible for gas transfer called alveoli (Berg 2016). COPD is a significant public health issue, especially in low- and middle-income countries, where nearly 90% of deaths from COPD occur (WHO 2020). COPD was the third leading cause of global deaths in 2016 with about 3 million deaths (WHO 2018), and it is expected to remain in the third ranking until 2030 (WHO 2013).

The main risk factors for COPD are tobacco smoking, second-hand smoking, air pollution, and exposure to fuel oil fumes (GOLD 2022). COPD is characterized by a non-reversible airflow obstruction in the lungs. Exposure to irritants stimulates mucus production and damages cilia that clear away mucus and dirt; this causes air to be trapped inside airways, leading to hyperinflation (GOLD 2022; Ramos 2014). Airflow obstruction in emphysema is due to the loss of elastin, which increases lung compliance and decreases elastic recoil (Costanzo 2019). In other words, the lungs lose their ability to return spontaneously to their resting position after inhalation. COPD is a cause of disability as it affects people's ability to breathe normally and has systemic, severe, and long-term effects (Agustí 2005). Post-bronchodilator spirometry is the primary test to measure airflow obstruction. It confirms airflow limitation if the ratio between forced expiratory volume in 1 second (FEV1) and forced vital capacity (FVC) is less than 70% (GOLD 2022).

Clinically, the most common symptoms are dyspnea, chronic cough, wheezing, and sputum production. Dyspnea — also known as shortness of breath or breathlessness — is the most common symptom reported by patients with COPD and is associated with a deterioration in their quality of life and physical activity (Anzueto 2017). Dyspnea results from multiple mechanisms, such as air trapping and dynamic hyperinflation, and it is associated with a significant load on the respiratory muscles (Padula 2006; O'Donnell 2007). The Global Initiative for Chronic Obstructive Lung Disease (GOLD) suggests a therapeutic strategy based on medical history, clinical symptoms, and life quality (ABCD stages) (GOLD 2022).

Description of the intervention

Inspiratory muscle training (IMT), also known as respiratory or ventilatory muscle training, aims to improve inspiratory muscle strength and endurance through a series of breathing exercises. It was developed in the late 1970s (Andersen 1979; Belman 1980; Leith 1976), and it has been used in people with respiratory diseases such as COPD and asthma. IMT focuses on enhancing the performance of respiratory muscles and on improving respiratory symptoms and exercise capacity (Padula 2006).

There are three main categories of IMT devices: threshold loading devices, passive and electronic flow resistive devices, and isocapnic hyperpnea devices (Belman 1994). Other devices exist, and they are reported in Menzes 2018. Most threshold trainers have an adjustable spring-loaded valve to set the resistance level from 9 cm of water (cmH₂O) to 41 cmH₂O (or from 7 cmH₂O to 41 cmH₂O)

and allow changes in resistance by 2 cmH₂O (Menzes 2018). The threshold pressure is independent of the breathing pattern (Geddes 2005). The passive-resistive trainer contains holes of different diameters: the biggest hole provides the lowest resistance, whereas the narrowest hole offers the highest resistance (McConnell 2004; Menzes 2018). The respiratory load can be selected by turning the dial towards the chosen hole. However, unlike the threshold device, passive-resistive trainers depend on the inspiratory flow (Wu 2017). The electronic resistive device is similar to the passive-resistive trainer, and it has the advantage of dynamically adapting the flow resistance (Menzes 2018). Isocapnic hyperpnea trainers are based on low load and high respiratory flow (60% to 90% of maximal voluntary ventilation (MVV)) so that respiratory muscles contract at a higher speed (for an extended time) (McConnell 2004). That device contains a rebreathing bag to maintain physiological rates of CO₂, so patients breathe both fresh air and some of the expired CO₂. In addition to multiple devices of IMT, protocols for this therapy differ between teams in terms of frequency, duration, and supervision (Langer 2015).

Various measures are used to evaluate respiratory muscle strength (Laveneziana 2019). Maximal static inspiratory mouth pressure (P_{imax}) is the most commonly used technique to assess the strength of the diaphragm and other inspiratory muscles (Pessoa 2014). It is calculated through a mouthpiece connected to a manometer, either at residual volume or at functional residual capacity (Laveneziana 2019). However, this technique requires the co-operation of patients. Other approaches to assess inspiratory muscle strength exist, such as sniff nasal inspiratory pressure (SNIP) based on a pressure sensor attached to a catheter placed in the nostril (Maillard 1998), phrenic nerve electric transcutaneous stimulation, and phrenic nerve magnetic stimulation (Caruso 2015).

How the intervention might work

Unlike inspiration, expiration is a passive process. In people with COPD, elastin in the lungs can be reduced, leading to incomplete expiration; this means air is trapped in the airways and leads to hyperinflation (Papandrinopoulou 2012). Static and dynamic distention caused by hyperinflation explains, in part, the pathophysiology of respiratory muscle dysfunction in COPD (O'Donnell 2006). Indeed, inspiratory muscles (the diaphragm, intercostal muscles, and the sternocleidomastoid muscle (SCM)) are exposed to an important load generated from hyperinflation and high airway resistance (Caron 2011). In the early stages of COPD, inspiratory muscles try to adapt to these circumstances. For example, type II muscle fibers of the diaphragm switch into type I, which are highly resistant to fatigue (Clanton 2009). There is also an increase in blood capillaries (Doucet 2004) as well as the oxidative capacity (since type I fibers have high mitochondrial density and enzymes that support the oxidative pathway, so the ability to use oxygen will be increased) (Ottenheijm 2008). At an advanced stage of the disease, oxidative stress, gas exchange abnormalities, and the changes in the chest cavity overcome adaptation mechanisms and the diaphragm will be in a position of impaired mechanical advantage. It loses up to 60% of its muscle tissue and becomes shorter and more horizontal, leading to ineffective mechanical function (Caron 2011; Ottenheijm 2008; Salito 2015). A study showed that IMT induced structural and anatomical changes in external intercostal muscles by changing the distribution of type I fibers and increasing the size of type II fibers

(Ramirez Sarmiento 2002). Overall, inspiratory muscle weakness is associated with dyspnea and respiratory failure, despite the ability of the diaphragm to adapt itself to hyperinflation (Bégin 1991; Caron 2011). IMT may improve the strength and endurance of these muscles.

In COPD, it is difficult to work on expiration flow and volume due to mechanical changes. However, it is possible to work on inspiration since it is an active process. In other words, strengthening of the inspiratory muscles increases the inspiratory flow (so there will be an increase in the tidal volume (TV)), decreases the inspiratory time, and improves the expiratory time (Beaumont 2018; Charususin 2016).

Why it is important to do this review

The clinical symptoms most often reported by patients with COPD are dyspnea, a decline in exercise capacity and an impairment in their quality of life (Spruit 2013). It is recommended to start pulmonary rehabilitation (PR) as soon as possible, ideally either during hospitalisation or soon after discharge from the hospital (Spruit 2013). Current guidelines recommend an optimal duration of eight weeks for a PR program (Rochester 2015). Usually, PR consists of physiotherapy, nutritional and psychosocial care, patient therapeutic education, and upper and lower limb training (Beaumont 2015; Spruit 2013).

The American Thoracic Society (ATS) reported that IMT may be beneficial as a stand-alone intervention and when added to PR in patients with respiratory muscle weakness (Spruit 2013). However, the potential effects of combining IMT and PR are still unclear. Indeed, some recent randomized controlled trials (RCTs) did not find significant improvements in patients with severe COPD (Beaumont 2018; Charususin 2018), and most of the published clinical trials and meta-analyses recommended further investigation (Beaumont 2018a; Gosselink 2011; Langer 2015; Lötters 2002). Furthermore, there is a lack of certainty in the linear relation between P_{Imax}, FEV₁, and clinical outcomes. In other words, many RCTs showed that IMT improves P_{Imax}, but the extent to which this improvement is clinically significant (i.e. the minimal clinically important difference) has not yet been proved (Beaumont 2018; Schultz 2018). The benefits of unsupervised IMT are also unclear (Langer 2015).

Although many meta-analyses have been published on different modes and modalities of IMT (Geddes 2005; Geddes 2008; Gosselink 2011; Lötters 2002; O'Brien 2008; Smith 1992), several questions remain unanswered. For example, the optimum duration of a training program has not been established, nor has the effect of IMT on dyspnea and quality of life. There is also a need to investigate its additional effect when added to PR. Moreover, the published meta-analyses are a few years old, and there are now other studies to be included. A recent clinical trial (Langer 2015), showed that factors other than inspiratory muscle weakness might influence the performance of IMT, such as the variety of protocols, and it is worth exploring these.

OBJECTIVES

To assess the effect of inspiratory muscle training (IMT) on chronic obstructive pulmonary disease (COPD) as a stand-alone intervention and when combined with pulmonary rehabilitation (PR).

METHODS

Criteria for considering studies for this review

Types of studies

We included published randomized controlled trials (RCTs), as full-text articles or abstracts, as well as unpublished RCTs. We included abstracts if they reported at least the number of participants in each group, the duration of the intervention, and the training load. We accepted trials with more than two arms. We excluded observational studies, quasi-RCTs and cross-over trials since no washout period for IMT has been established.

Types of participants

We included people with COPD diagnosed according to international standards (GOLD 2022), at any stage of the disease. We placed no restrictions on age, duration, setting, or the kind of pulmonary rehabilitation. We planned to include RCTs with different conditions for the same intervention of interest as long as we could obtain the data of participants with COPD separately.

We classified COPD according to GOLD 2022 stages based on the predicted value of Forced Expiratory Pressure in 1 Second (FEV₁):

- GOLD 1 - mild: FEV₁ ≥ 80% predicted
- GOLD 2 - moderate: 50% ≤ FEV₁ < 80% predicted
- GOLD 3 - severe: 30% ≤ FEV₁ < 50% predicted
- GOLD 4 - very severe: FEV₁ < 30% predicted

Types of interventions

The review consists of two comparisons, as follows.

- IMT plus PR versus PR
- IMT versus no treatment or sham

First, we included trials that explored the benefit of combining IMT and PR compared to PR only. PR consists of, but is not limited to, exercise training, physiotherapy, therapeutic education, and nutritional and psychosocial care (McCarthy 2015). We included different types of IMT irrespective of the mode of delivery: resistance training (high load, low frequency) or endurance training (low load, high frequency), device (i.e. threshold loading, resistive flow device, isocapnic hyperpnea). We made no restrictions on the duration, supervision (home-based or in a healthcare setting), or timing (during hospitalization or later) of the intervention. We excluded studies where the training was conducted only once per week (face-to-face or distance sessions), regardless of the total duration of the clinical trial. The minimum accepted training load was 30% of P_{Imax} or more (Hill 2010). We also excluded trials that used a resistive device without controlling the breathing pattern. If a study conducted an incremental training load that started less than 30% of P_{Imax}, we considered only the follow-up from which the load was equal to our threshold.

According to the proportion of supervised sessions, we defined supervision as:

- under 20%: unsupervised;
- 20% to 70%: partially supervised; and
- above 70%: fully supervised.

For the second comparison, we compared IMT with control or sham. We defined sham training as using a resistance of less than 30% of P_{Imax}. We accepted control groups if they did not receive any intervention or received an intervention other than exercise training to blind participants (e.g. therapeutic education). We made the exception for breathing exercises if participants in the control group did not receive more than one type of training (e.g. diaphragmatic breathing, pursed lips breathing), and the purpose was not to compare it with IMT.

Types of outcome measures

We analysed the following outcomes in the review, but we did not use them as a basis for including or excluding studies.

Primary outcomes

Dyspnea: the essential scale for our primary analysis is the Borg scale (Borg 1982). We only included the Borg score when it was measured at isotime. We analysed all the scales reported by the trials as long as they were validated, and when possible, we combined them in a meta-analysis. The other included scales were: Baseline and Transition Dyspnea Indexes (BDI-TDI) (Mahler 2005), and Modified Medical Research Council (mMRC) (Bestall 1999).

Functional exercise capacity: this can be assessed through multiple tests. We did not exclude studies based on the test used. However, for our analysis, we considered that the most important measurement is the six-minute walk distance (6MWD) (Holland 2014). Therefore, we included it in the summary of findings table and considered it for subgroup analysis. We included other tests and reported them either in qualitative or quantitative analysis.

Health-related quality of life: we accepted any scales as long as they were validated. This includes the St. George's Respiratory Questionnaire (SGRQ) (Jones 1992), the chronic respiratory questionnaire (CRQ) (Wijkstra 1994) and the COPD assessment test (CAT) (Jones 2009).

Secondary outcomes

Inspiratory muscle strength: measured by maximal static inspiratory mouth pressure (P_{Imax}) (Laveneziana 2019).

Laboratory exercise tests: we were primarily interested in the maximal oxygen uptake (VO_{2peak}), which could be measured through:

- incremental cycle ergometer test;
- endurance cycle ergometer test;
- treadmill test.

Respiratory muscle endurance:

- **Respiratory muscle endurance pressure (P_{thmax}):** measured by incremental load testing (Laveneziana 2019). It is the maximally tolerated pressure when breathing against a continuously increasing load.
- **Respiratory muscle endurance time (T_{lim}):** measured by constant load testing (Laveneziana 2019). It is the time an individual can maintain breathing against a fixed load. It can be carried out either through a threshold/resistive or isocapnic hyperpnea device.

- **Maximal voluntary ventilation (MVV):** this is the total volume of expired air between 12 seconds to 15 seconds of deep and fast respiration. MVV is usually compared to predicted MVV (calculated through the forced expiratory volume at 1 second) (Wood 2017).

Respiratory function:

- forced expiratory volume at 1 second (FEV1)
- residual volume

Adverse events: as defined by the trial authors.

We collected outcomes irrespective of the time frame and summary statistics (change from baseline or final values), with a preference for change score. For each trial, we analysed only the outcomes listed above and not all the outcomes reported in the trial. However, we included all the tests and measurements used for the same outcome.

Search methods for identification of studies

Electronic searches

We searched for all published and unpublished RCTs regarding IMT for COPD, in consultation with the Cochrane Airways Information Specialist. We did not apply restrictions on language or publication status (i.e. published, ongoing, or unpublished).

We searched the following databases for relevant trials in October 2020 and we updated our literature search on 23 August 2021 and on 20 October 2022.

- Cochrane Airways Trials Register (Cochrane Airways 2022), via the Cochrane Register of Studies, all years to date (searched 13 October 2020, 23 August 2021 and 20 October 2022)
- Cochrane Central Register of Controlled Trials (CENTRAL), via the Cochrane Register of Studies, all years to date (searched 13 October 2020, 23 August 2021 and 20 October 2022)
- MEDLINE Ovid SP ALL, 1946 to 12 October 2020 (searched 13 October 2020, 23 August 2021 and 20 October 2022)
- Embase Ovid SP, 1974 to week 41 2020 (searched 13 October 2020, 23 August 2021 and 20 October 2022)
- PsycINFO Ovid SP, 1967 to October week 1 2020 (searched 13 October 2020, 23 August 2021 and 20 October 2022)
- Cumulative Index to Nursing and Allied Health Literature (CINAHL) EBSCO, 1937 to 13 October 2020 (searched 13 October 2020, 23 August 2021 and 20 October 2022)
- Physiotherapy Evidence Database (PEDro), 1999 onwards (searched 13 October 2020, 23 August 2021 and 20 October 2022)

We searched the following trials registries.

- US National Institutes of Health Ongoing Trials Register, [ClinicalTrials.gov](https://clinicaltrials.gov) (searched 13 October 2020, 23 August 2021 and 20 October 2022)
- World Health Organization [International Clinical Trials Registry Platform](https://www.who.int/clinical-trials-registry) (searched 13 October 2020, 23 August 2021 and 20 October 2022)

The database search strategies are listed in [Appendix 1](#). The search strategy was developed in MEDLINE by the Cochrane Airways Information Specialist in collaboration with the authors and peer-

reviewed by another Cochrane Information Specialist using the PRESS checklist (McGowan 2016). The MEDLINE search strategy was then adapted appropriately for each database.

All databases and trial registries were searched from their inception to the present, with no restriction on language or type of publication. Hand-searched conference abstracts and grey literature were identified through the Cochrane Airways Trials Register and CENTRAL. When possible, A native-language speaker translated studies written in a language other than English.

Searching other resources

We checked the reference lists of all primary studies and reviews for additional references. We searched for relevant manufacturers' websites for device information. We searched on PubMed for errata or retractions from included studies published in full-text and, when possible, reported the date this was done within the review.

Data collection and analysis

Selection of studies


Two review authors from complementary disciplines (OA and WF) independently screened the titles and abstracts of the search

results using [Covidence](#), and they coded them as 'include' (eligible or potentially eligible/unclear) or 'exclude'. We retrieved full-text study reports of all potentially eligible studies, and two review authors (OA and WF) independently screened them for inclusion while also recording the reasons for excluding ineligible studies. We retrieved the full text of potentially relevant reports and removed duplicate records using [Covidence](#).

When appropriate, we contacted the study authors to request further information. We also contacted the manufacturer of the IMT device when we did not understand the concept of training. We resolved disagreements through discussion, without the need for a third review author. We identified and excluded duplicates and collated multiple reports of the same study so that each study, rather than each report, is the unit of interest in the review.

We recorded the selection process in sufficient detail to complete a PRISMA flow diagram ([Figure 1](#)) and the '[Characteristics of excluded studies](#)' table ([Moher 2009](#)).

Figure 1.



Data extraction and management

Two review authors (OA and AK) used a data collection form that we piloted on at least one study in the review to extract characteristics from included studies. We extracted the data using [Covidence](#) and an Excel spreadsheet. We planned to record any missing information as unclear or not described. Each form included the following information.

- General information: study ID, author contact detail, and the person who is completing the form
- Methods: aims of the study, study design, total study duration
- Participants: inclusion criteria, exclusion criteria, total number randomized, number randomized per group, mean age, age range, sex, COPD stage, clusters (if applicable), number missing, reasons for missing participants, number of participants moved from one group to another, reasons moved, baseline imbalances, and subgroup analysis
- Intervention/comparison groups: type of group, type of intervention, type of control, duration, supervision, setting, device, intensity, frequency, type of training (strength/endurance)
- Outcomes: primary and secondary outcomes, baseline characteristics, and time points
- Notes: funding for studies and notable conflicts of interest of trial authors

Two review authors (OA and AK) independently extracted outcome data from included studies. We noted in the [Characteristics of included studies](#) tables if outcome data were not reported in a usable way or if some of the data were missing. We resolved disagreements by reaching a consensus or by involving a third review author (SK). If we identified multiple reports from the same study, we would extract data from all reports directly into a single data collection form. One review author (OA) transferred data into the Review Manager 5 file ([Review Manager 2020](#)). We double-checked that data had been entered correctly by comparing data presented in the systematic review against the study reports. A second review author (WF) spot-checked study characteristics for accuracy against the study report.

Assessment of risk of bias in included studies

Two review authors (OA and TL) assessed the risk of bias independently for each study using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2021](#)). We resolved any disagreements by discussion or by involving another review author (SK). We assessed the risk of bias according to the following domains.

- Bias arising from the randomization process
- Bias due to deviations from intended interventions
- Bias due to missing outcome data
- Bias in measurement of the outcome
- Bias in selection of the reported result

We used RoB 2 to assess risk of bias in randomized studies ([Sterne 2019](#)). We used the RoB 2 Excel tool to complete the risk of bias assessment. We used RevMan Web to generate traffic light plots of the domain-level judgments for each outcome ([RevMan Web 2022](#)).

Our effect of interest was the assignment to the intervention at baseline and our main outcomes were those listed in the summary of findings tables. We judged each outcome as being at low risk, some concerns, or high risk according to the RoB 2 algorithm. We provided a quote from the study report, together with a justification for our judgment, in the risk of bias table.

Where information on risk of bias relates to unpublished data or correspondence with a trial author, we noted this in the risk of bias table. To detect reporting bias, we compared the study protocol with the published report, and we contacted the study authors to identify missing or partially reported data. If more than 10 studies were included in the meta-analysis, we created a funnel plot to explore publication bias. None of the included studies was a cluster-RCT.

We incorporated the risk of bias assessment in the [Results](#) section of the review and it was also part of the GRADE assessment of the certainty of evidence (along with precision, directness, consistency, and publication bias). When considering treatment effects, we took into account the risk of bias for the studies that contributed to that outcome. Our primary analysis included all the studies without taking the risk of bias judgments into account.

Assessment of bias in conducting the systematic review

We conducted the review according to the published protocol and reported any deviations from it in the '[Differences between protocol and review](#)' section of this systematic review.

Measures of treatment effect

All of our outcomes were continuous data. We calculated a mean difference (MD) with 95% confidence intervals (CIs) where studies used the same scale, and the standardised mean difference (SMD) with 95% CIs where studies used different scales to measure the same concept. We interpreted SMD analysis following the rule of thumb based on Cohen's d effect size ([Cohen 1988](#)):

- 0.2 represents a small effect;
- 0.5 represents a medium effect;
- 0.8 represents a large effect.

Depending on how studies reported ordinal data, we analysed the scales as continuous (since all of them were longer than five). We presented all results with a 95% CI.

We undertook a meta-analysis when it was meaningful. That is to say, it made sense to combine the data, and the populations, interventions and outcomes were similar enough to be pooled in the same forest plot. If both change-from-baseline and end-point scores were available for continuous data, we used change-from-baseline, unless there was a low correlation between measurements in individuals.

If adjusted analyses were available (analysis of variance (ANOVA) or analysis of covariance (ANCOVA)), we used these as a preference in our meta-analyses. If the adjusted MD was reported, we included it in the meta-analysis using the Generic Inverse Variance method unless adjusted and unadjusted analyses were similar.

Unit of analysis issues

- **Cluster-RCTs and dichotomous outcomes:** were not included in the review.

- **Repeated observations on participants:** if studies reported outcomes at multiple time points, we chose the longest follow-up period to keep consistent with the studies. We divided the duration of follow-up into categories to explore possible differences in the effect estimate. More information is in '[Subgroup analysis and investigation of heterogeneity](#)'.
- **Studies with more than two groups:** we included only the relevant arms.
- **Two comparisons from the same study within the same meta-analysis:** we combined the active arms or halved the control group to avoid double-counting.

Dealing with missing data

We requested missing or unclear numerical data from study authors (such as for conference abstracts; randomization, the training load). We did not use imputation because most of the data were available, and it was not possible to receive the participants' data. For studies that reported only the overall effect estimate without providing data for each intervention group, we used the generic inverse variance method to meta-analyse them.

We used the methods recommended by [McGrath 2020](#) to convert median to mean.

When the data were presented only graphically, and in case we could not get numerical information from the study authors, we used [WebPlotDigitizer](#) to extract them from the graphs.

Assessment of heterogeneity

We assessed statistical heterogeneity through visual inspection by detecting overlapping confidence intervals in the forest plot. We used the Chi² test with a P value of 0.05 to indicate the statistical significance and the I² test ([Higgins 2003](#)), to explore statistical heterogeneity (we considered a value over 50% to represent substantial heterogeneity). We performed subgroup analysis to investigate heterogeneity. We also discussed clinical heterogeneity (e.g. COPD stages) and methodological heterogeneity (e.g. duration of the intervention, number of sessions per week, the total number of sessions, the training load).

Assessment of reporting biases

We created funnel plots to explore possible small-study and publication biases for functional exercise capacity (6MWD) and respiratory muscle strength (P_{Imax}) since they were included in the summary of findings table and more than 10 trials explored these outcomes.

Data synthesis

We expected the included studies to have many variables that could influence the pooled effect estimate. Therefore, we used a random-effects model for the analysis. We ran a meta-analysis when it was appropriate; that is, where it made sense to combine different effect estimates and the studies were homogeneous enough to allow reliable interpretation of the analysis.

Ordinal outcomes were meta-analysed as continuous data following the guidance in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Deeks 2021](#)). Two authors (OA and AK) analysed the data using RevMan 5 ([Review Manager 2020](#)), and RevMan Web ([RevMan Web 2022](#)).

Subgroup analysis and investigation of heterogeneity

We carried out the following subgroup analyses.

- Duration of intervention: short-term (less than four weeks), medium-term (between four weeks and seven weeks and six days), long-term (eight weeks and longer).
- Respiratory muscle strength (P_{Imax}): studies with participants with respiratory muscle weakness (the mean baseline P_{Imax} of the participants was less than or equal to 60 cmH₂O) or without respiratory muscle weakness (the mean baseline P_{Imax} of the participants was greater than 60 cmH₂O).

We used the following outcomes in the subgroup analyses.

- Dyspnea: Baseline Dyspnea Index-Transition Dyspnea Index (BDI-TDI)
- Functional exercise capacity: 6-minute walk distance (6MWD)
- Respiratory muscle strength (P_{Imax})

We included within-study data when available. We used the formal test for subgroup interactions in RevMan Web ([RevMan Web 2022](#)).

Sensitivity analysis

We performed the following sensitivity analyses.

- We removed from the primary analysis studies judged to be at high risk of bias and some concerns.
- We compared the results of the random-effects and fixed-effect models for the BDI-TDI, the 6MWD, the SGRQ and P_{Imax}.

We considered studies to be at high risk of bias overall if we judged at least one of the domains to be high risk.

Summary of findings and assessment of the certainty of the evidence

We created summary of findings tables ([Summary of findings 1](#), [Summary of findings 2](#)) including the following outcomes.

- Dyspnea: Borg scale, mMRC and BDI-TDI
- Functional exercise capacity: 6MWD
- Health-related quality of life: SGRQ, COPD Assessment Test (CAT)
- Inspiratory muscle strength: P_{Imax}

We used the five GRADE considerations (risk of bias, consistency of effect, imprecision, indirectness, and publication bias) to assess the quality of the body of evidence as it relates to the studies that contribute data for the prespecified outcomes. Our time point was the end of the study. We used the methods and recommendations described in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Schünemann 2021](#)), and the *GRADE Handbook*, using [GRADEpro GDT](#) software. We justified all decisions to downgrade the quality of studies using footnotes and we made comments to aid the reader's understanding of the review where necessary.

RESULTS

Description of studies

Details are available in [Characteristics of included studies](#); [Characteristics of excluded studies](#); [Characteristics of studies awaiting classification](#); [Characteristics of ongoing studies](#) tables.

Results of the search

We identified from the literature search 7379 records through database searching and 378 from other sources ([International Clinical Trials Registry Platform \(ICTRP\)](#); [Epistemonikos](#)).

After removing duplicates and screening titles and abstracts, we checked the eligibility of 277 studies through full-text review (see [Figure 1](#)). Searching the reference lists of past published systematic reviews did not reveal further eligible records. In the end, we included 55 studies in this review, of which six RCTs had more than two arms. Of the 55 studies, 53 contributed data to meta-analyses.

When contact details were available, we contacted the study authors for clarification. We classified 42 studies as 'awaiting classification' because of insufficient data (although we contacted study authors) or because we couldn't find the abstract (only the title was available in Covidence), and 15 as ongoing studies.

One study in Chinese ([Zhou 2016](#)), and one in Spanish ([Bustamante 2007](#)), were translated into English by native speakers (see [Acknowledgements](#)). One study in Japanese ([Okura 2020](#)) was translated into English using [Google Translate](#).

Included studies

Comparison 1: pulmonary rehabilitation plus inspiratory muscle training versus pulmonary rehabilitation

Population

We included 22 RCTs with 1446 participants in this comparison. We classified the COPD stages according to post-bronchodilator forced expiratory volume in one second (FEV1) ([GOLD 2022](#)). They ranged from mild to moderate ([Tout 2013](#)), moderate to severe ([Abedi Yekta 2019](#); [Berry 1996](#); [Dekhuijzen 1991](#); [Larson 1999](#); [Magadle 2007](#); [Majewska-Pulsakowska 2016](#); [Schultz 2018](#); [Wang 2017](#)), mild to very severe ([Paneroni 2018](#)), moderate to very severe ([Beaumont 2015](#); [Charususin 2018](#); [Fanfa Bordin 2020](#); [Mador 2005](#); [Tounsi 2021](#)), and severe to very severe ([Beaumont 2018](#); [Dellweg 2017](#); [Weiner 1992](#); [Weiner 2000](#)). Three studies did not report COPD stage.

The number of participants in the intervention group (PR+IMT) was 742. The mean age ranged from 51.33 to 70.8 years, and the mean body mass index (BMI) ranged from 21.31 to 28.8 kg/m². The number of participants in the control group (PR only) was 704. The mean age ranged from 53.5 to 70.8 years, and the mean BMI ranged from 22.4 to 29.68 Kg/m². In the studies that reported gender, there were around 763 men and 482 women.

Intervention

Pulmonary rehabilitation

The rehabilitation programs varied across the studies. They ranged from training with only a treadmill ([Abedi Yekta 2019](#)), with only a cycle ergometer ([Larson 1999](#)), and a combination

of training protocols in the remaining studies (training with cycle and treadmill, limb muscles strengthening exercises, stair climbing, and therapeutic patient education programs). Five studies monitored training intensity according to heart rate: 50% ([Weiner 1992](#)), 60% ([Abedi Yekta 2019](#); [Fanfa Bordin 2020](#)), 80% ([Dekhuijzen 1991](#)) and 85% ([Larson 1999](#)). One study ([Tounsi 2021](#)) individualized the training program based on 60% to 80% of the average speed achieved during the six-minute walk test.

Inspiratory muscle training

Two studies trained their participants for two days a week ([Abedi Yekta 2019](#); [Tout 2013](#)), six studies for three days a week ([Fanfa Bordin 2020](#); [Mador 2005](#); [Magadle 2007](#); [Wang 2017](#); [Weiner 1992](#); [Weiner 2000](#)), eight studies reported five days a week ([Beaumont 2015](#); [Beaumont 2018](#); [De Farias 2019](#); [Dekhuijzen 1991](#); [Dellweg 2017](#); [Larson 1999](#); [Majewska-Pulsakowska 2016](#); [Paneroni 2018](#)), and four studies trained their participants for the whole week ([Berry 1996](#); [Charususin 2018](#); [Schultz 2018](#); [Tounsi 2021](#)). Two studies ([Masanga 2011](#); [Sykes 2005](#)) did not provide details about the training. The number of weeks ranged from two weeks ([Paneroni 2018](#)) to 24 weeks ([Magadle 2007](#)).

The training was unsupervised in one study ([Majewska-Pulsakowska 2016](#)), partially supervised in three studies ([De Farias 2019](#); [Larson 1999](#); [Schultz 2018](#)), and fully supervised in the remaining studies.

Three trials performed endurance training, using SpiroTiger ([De Farias 2019](#); [Paneroni 2018](#)), and normocapnic hyperpnea ([Mador 2005](#)), respectively. The other RCTs conducted strength training with Powerbreathe devices, Threshold IMT device, and Respifit S-Unit. Five studies did not change the training loads during the study, ranging from 30% to 80% of P_{Imax}. In the other RCTs, the training load increased from 15% to 60% of P_{Imax} ([Weiner 2000](#)), from 30% to 60% of P_{Imax} ([Abedi Yekta 2019](#); [Larson 1999](#); [Majewska-Pulsakowska 2016](#); [Schultz 2018](#); [Sykes 2005](#); [Tout 2013](#)), from 50% to 80% of P_{Imax} ([Berry 1996](#); [Tounsi 2021](#); [Weiner 1992](#)), from 35% to 80% ([De Farias 2019](#)), from 50% to 60% ([Beaumont 2018](#)), from 50% to 84% of P_{Imax} ([Charususin 2018](#)), and from 66% to 85% of Maximal Voluntary Ventilation (MVV) ([Paneroni 2018](#)).

In [Magadle 2007](#) and [Weiner 2000](#), participants received respectively three months and six months of PR before being allocated to continue with PR alone or to receive IMT.

Comparisons

All the studies focused on our main comparison (PR+IMT vs PR), and there were no indirect comparisons. Three RCTs ([Abedi Yekta 2019](#); [Larson 1999](#); [Majewska-Pulsakowska 2016](#)), had four arms, including both comparisons. One trial had three arms ([De Farias 2019](#)). We included the appropriate comparison separately.

Primary outcomes

Dyspnea: two studies explored dyspnea with the Borg scale at isotime and the Modified Medical Research Council (mMRC) scale, one study ([Schultz 2018](#)), with BDI-TDI, and two studies ([Beaumont 2015](#); [Beaumont 2018](#)), reported the Multidimensional Dyspnea Profile (MDP).

Functional exercise capacity: 12 studies measured functional exercise capacity with the 6MWD, three studies used the 12-

minute walk distance (12MWD), three studies used exercise time, and five studies used maximal exercise capacity (Wmax). For the latter measurement, [Charususin 2018](#) calculated Wmax by asking participants to cycle at a load of 20 watts (w) and then increasing it by 10 w/min until symptom limitation. [Mador 2005](#) followed a similar protocol starting at no load until the participant could no longer continue cycling for 30 seconds. [Dekhuijzen 1991](#) increased the load by 10% of the predicted Wmax, which was measured through the following formula: "Wmax predicted = 1.7x weight (kg) + 40x FEV₁(L)-25". [Wang 2017](#) chose an incremental load of 5 w/min or 10 w/min.

Health-related quality of life (HRQoL): seven studies assessed HRQoL with the SGRQ, three studies used the Chronic Respiratory Disease Questionnaire (CRQ), two studies used the COPD Assessment Test (CAT), and one study used the Clinical COPD Questionnaire (CCQ).

Secondary outcomes

Inspiratory muscle strength (P_{Imax}): 17 studies reported P_{Imax}. Eight of these studies measured it at residual volume.

Laboratory exercise test (VO_{2peak}): five trials measured VO_{2peak}. All studies except for [Berry 1996](#) reported VO_{2peak} in L/min. So we used the mean weight of each group in [Berry 1996](#) to convert from mL/kg/min to L/min.

Respiratory muscle endurance strength (P_{thmax}): two studies measured P_{thmax} ([Larson 1999](#); [Weiner 1992](#)).

Respiratory muscle endurance time (T_{lim}) was measured by asking participants to breathe until exhaustion against a load of 30% of P_{Imax} ([Paneroni 2018](#)), 50% to 60% of P_{Imax} ([Charususin 2018](#)), 70% of P_{Imax} ([Dekhuijzen 1991](#)), 70% of MVV ([Mador 2005](#)), and 70% to 75% of MVV ([Paneroni 2018](#)).

FEV₁: six studies reported FEV₁ as percentage of predicted and liters. Three studies ([Berry 1996](#); [Paneroni 2018](#); [Wang 2017](#)), reported MVV, and one study ([Charususin 2018](#)), reported residual volume

Only one abstract ([Masanga 2011](#)), reported adverse events.

For both our primary and secondary outcomes, we did not include data from [Sykes 2005](#) in our primary analysis.

Comparison 2: inspiratory muscle training versus control/sham

Population

We included 37 RCTs with 1021 participants in this comparison. As in comparison 1, we classified COPD stages according to post-bronchodilator FEV₁ ([GOLD 2022](#)). They ranged from moderate ([Leelarungrayub 2017](#)), severe ([Lisboa 1997](#), [Weiner 2003](#); [Weiner 2006](#)), mild to severe ([Bavarsad 2015](#)), mild to very severe ([Dacha 2019](#)), moderate to severe ([Abedi Yekta 2019](#); [Belman 1988](#); [Bustamante 2007](#); [Harver 1989](#); [Hsiao 2003](#); [Koppers 2006](#); [Larson 1999](#); [Majewska-Pulsakowska 2016](#); [Petrovic 2012](#); [Saka 2021](#); [Sanchez Riera 2001](#); [Scherer 2000](#); [Wu 2017](#); [Xu 2018](#)), moderate to very severe ([Berton 2015](#); [Chuang 2017](#); [Heijdra 1996](#); [Langer 2018](#); [Nikoleto 2016](#); [Saher 2021](#)), and severe to very severe ([Beckerman 2005](#); [Covey 2001](#); [Hill 2006](#); [Hill 2007](#); [Kim 1993](#); [Larson 1988](#); [Preusser 1994](#); [Ramirez Sarmiento 2002](#); [Zhou 2016](#)).

The number of participants in the IMT group was 526. The mean age ranged from 51.8 to 70.4 years, and the mean BMI ranged from 19.25 to 29 kg/m². For studies that reported gender, there were in total 268 men and 129 women. The number of participants in the control group (control/sham) was 495. The mean age ranged from 54.2 to 71.1 years, and the mean BMI ranged from 18.54 to 28.8 kg/m². For studies that reported gender, there were 269 men and 132 women

Intervention

Inspiratory muscle training

Participants trained from two days a week ([Abedi Yekta 2019](#)), to the whole week ([Beckerman 2005](#); [Berton 2015](#); [Bustamante 2007](#); [Dacha 2019](#); [Harver 1989](#); [Kim 1993](#); [Koppers 2006](#); [Langer 2018](#); [Larson 1988](#); [Petrovic 2012](#); [Xu 2018](#)). The duration of the intervention ranged from two weeks ([Saher 2021](#)) to a year ([Beckerman 2005](#)), and the total duration of training ranged from four hours ([Abedi Yekta 2019](#)) to 144 hours ([Beckerman 2005](#)).

Eight studies conducted training with resistive devices. Three studies ([Belman 1988](#); [Harver 1989](#); [Wu 2017](#)), used Pflx (Respironics Inc, Pittsburgh, PA, USA) device, one study ([Leelarungrayub 2017](#)), used Portex (Smith Medical ASD), one study ([Hsiao 2003](#)), used Respirix (Respirex[®]2, DHD 22-1000, Diemolding Healthcare Division, Canastota, NY, USA), two studies ([Heijdra 1996](#); [Sanchez Riera 2001](#)), used INSPIRx (Intertech Resources Inc, Ft. Myers, FL; Respirecare Medical Inc., The Hague, the Netherlands), and one study ([Bavarsad 2015](#)), used Respivol (Medinet, Milano, Italy).

Participants underwent endurance training with Normocapnic Hyperpnea in two RCTs ([Koppers 2006](#); [Scherer 2000](#)), and both endurance and strength training in one RCT ([Petrovic 2012](#)), using Respifit S (Mauerbach, Austria). The remaining 25 studies conducted IMT with either Threshold IMT or Powerbreathe devices.

Two trials trained their participants 'as tolerated' ([Belman 1988](#); [Bustamante 2007](#)), and one trial ([Bavarsad 2015](#)), used an incentive spirometer device at a load equal to or more than the inspiratory volume. The training load increased from 30% to 60% of P_{Imax} in six trials ([Abedi Yekta 2019](#); [Covey 2001](#); [Larson 1999](#); [Majewska-Pulsakowska 2016](#); [Nikoleto 2016](#); [Saher 2021](#)), 30% to 45% in one trial ([Xu 2018](#)), from 15% to 60% of P_{Imax} in two trials ([Weiner 2003](#); [Beckerman 2005](#)), from 9% to 100% in one trial ([Leelarungrayub 2017](#)), from 50% to 100% of P_{Imax} in two trials ([Hill 2006](#); [Langer 2018](#)) and approximately from 50% to 133% in one trial (15 to 40 cmH₂O) ([Chuang 2017](#)) The remaining 23 studies chose a fixed load that ranged from 30% to 80% of P_{Imax}.

The training was fully supervised in eight studies, partially supervised in two studies, and unsupervised in 20 studies. Six studies did not report details of supervision ([Harver 1989](#); [Hill 2007](#); [Petrovic 2012](#); [Saher 2021](#); [Weiner 2003](#); [Weiner 2006](#)).

Control/sham

Twenty-two studies used a sham IMT while 15 studies did not provide any intervention to the control group. One study ([Cutrim 2019](#)), provided diaphragmatic breathing at a rate of 15 to 20 breaths/min for both the intervention and the control groups. Participants in the control group underwent therapeutic patient education and pursed lips breathing ([Covey 2001](#)), and therapeutic patient education ([Larson 1999](#)).

Comparison

All the studies focused on our main comparison (IMT versus control/sham), and there were no indirect comparisons. Two studies (Hsiao 2003; Wu 2017), had three arms, including two intervention groups (each group used a different device or protocol) and a control group. When two arms from the same study were included in a forest plot, we halved the number of participants in the control group.

As for comparison 1, we extracted the appropriate arms from Abedi Yekta 2019, Larson 1999 and Majewska-Pulsakowska 2016.

Primary outcomes

Dyspnea: six studies measured dyspnea with the Borg scale at isotime. Eight studies assessed dyspnea with BDI-TDI and four studies with the mMRC.

Functional exercise capacity: 16 studies measured functional exercise capacity with the 6MWD, three studies with 12MWD, seven studies with Wmax, five studies with exercise time, and two studies with the shuttle walk test (SWT). Hill 2006 and Koppers 2006 measured Wmax by increasing the work rate by 10% per minute; Larson 1999 asked the participants to warm up by pedalling for 3 minutes at 10 w followed by 2 minutes at 20 w, and then they started the graded cycle at 30 w; and Lisboa 1997 increased the load by 75 kpm every 2 minutes. Sanchez Riera 2001, and Wu 2017 increased the work rate by 10 w/min after one minute of unloaded pedalling.

To measure exercise time, Berton 2015 asked the participants to cycle at 75% of Wmax; Koppers 2006 set the load at 50% of P_{lmax}; Scherer 2000 measured it on a treadmill set to 80% of the incline and to 100% of the speed reached at VO_{2peak}; and Wu 2017 considered it as the time to reach Wmax.

Health-Related Quality of Life (HRQoL): six studies assessed HRQoL with the SGRQ, five studies used CRQ, two studies used CAT, two studies used SF-36 (Chuang 2017; Nikolettou 2016), and one study used the CCQ (Leelarungrayub 2017).

Secondary outcomes

Inspiratory muscle strength (P_{lmax}): 32 studies reported P_{lmax}. Fourteen studies measured it at residual volume, 10 studies at functional residual capacity, one study reported both measurements, and the remaining studies did not report the measurement method.

Laboratory exercise test (VO_{2peak}): 11 studies reported VO_{2peak}.

Respiratory muscle endurance pressure (P_{thmax}): five studies (Hill 2006; Koppers 2006; Preusser 1994; Ramirez Sarmiento 2002; Weiner 2003), followed the protocol of Nickerson 1982. One study (Larson 1999), started with an initial load of 30% of P_{lmax} and increased by 5.7 cmH₂O until exhaustion.

Respiratory muscle endurance time (T_{lim}): 10 studies reported T_{lim}. Langer 2018, Nikolettou 2016 and Petrovic 2012 asked the participants to breathe as long as possible against 50% to 60% of P_{lmax}, and Hill 2006 and Ramirez Sarmiento 2002 asked them to breathe against 80% of P_{lmax}. Bustamante 2007 set the load at 66%

of P_{lmax}, Hsiao 2003 at 70%, and Scherer 2000 at 66% or 75% of MVV.

MVV: two studies measured MVV (Belman 1988; Harver 1989)

Residual volume: two studies measured residual volume (Ramirez Sarmiento 2002; Hill 2006).

Forced expiratory volume at 1 second (FEV1): 10 studies reported FEV1 in %Pred, and 12 studies in litres.

Adverse events: none of the included studies reported adverse events.

Excluded studies

We excluded 133 studies after the full-text review. For further details, please refer to [Characteristics of excluded studies](#).

Risk of bias in included studies

We present the risk of bias assessment for each outcome, including all domain judgments and support for judgments, in a spreadsheet (Ammous 2022). We generated traffic light plots in most forest plots of our primary outcomes and for P_{lmax} in the secondary outcomes.

Comparison 1. Pulmonary rehabilitation plus inspiratory muscle training versus pulmonary rehabilitation

- **Dyspnea:** one study was at low risk of bias for dyspnea (Borg and mMRC; Charususin 2018), while three others were at high risk of bias. Larson 1999 had issues with intention to treat (ITT) analysis (the number of participants that were not analyzed could impact the results), missing data could depend on its true value and participants were not blinded. Both Beaumont 2018 and Wang 2017 did not blind participants.
- **Functional exercise capacity:** three studies were at low risk of bias for functional exercise capacity (6MWD; Beaumont 2018; Charususin 2018; Dellweg 2017). Most studies that we judged at some concern did not provide sufficient details about allocation concealment, excluding participants from the analysis (less than 5%) and only the journal article was available. We considered one study at high risk of bias because missingness is likely to depend on its true value (Paneroni 2018), and another study because of a lack of details about the randomization process (Tout 2013).
- **Health-related quality of life:** no study was at low risk of bias for the SGRQ and CAT. The main issue across the studies was the lack of blinding. One study was at low risk of bias for CRQ (Charususin 2018), and the two others were at high risk of bias because of issues with ITT analysis (Larson 1999) and lack of blinding (Mador 2005).
- **Inspiratory muscle strength:** six studies were at low risk of bias (P_{lmax}; Beaumont 2018; Charususin 2018; Dellweg 2017; Fanfa Bordin 2020; Schultz 2018; Tounsi 2021). Most studies were at some concern because of a lack of details about allocation concealment and only the journal article was available. Two studies were considered at high risk of bias (Larson 1999; Paneroni 2018), because missingness is likely to depend on its true value and one study because of a lack of details about the randomization process (Tout 2013).

Comparison 2. Inspiratory muscle training versus control/sham

- **Dyspnea:** one study was at low risk of bias for dyspnea (Borg; [Langer 2018](#)), two studies were at high risk of bias because participants were not blinded ([Larson 1988](#); [Petrovic 2012](#)), and the others were at some concern because lack of details about allocation concealment. Similarly, only [Langer 2018](#) was at low risk of bias for BDI-TDI. We judged [Harver 1989](#) at high risk of bias because the authors did not mention the reasons behind missing data, [Weiner 2003](#) because the data in the graph were different from the text and [Wu 2017](#) because participants were not blinded. For mMRC, from the four included studies, two studies were at low risk of bias ([Langer 2018](#); [Xu 2018](#)), one study at some concern because lack of details about allocation concealment ([Saka 2021](#)), and one study at high risk of bias because participants were not blinded ([ZhouL 2016](#)).
- **Functional exercise capacity:** two studies were at low risk of bias (6MWD; [Cutrim 2019](#); [Xu 2018](#)). Four studies were at high risk of bias because of issues with ITT analysis ([Beckerman 2005](#); [Hsiao 2003](#); [Ramirez Sarmiento 2002](#); [Saher 2021](#)), lack of details about missing data ([Leelarungrayub 2017](#)), and the data in the graph were different from the text ([Weiner 2003](#)). The remaining studies were at some concern because of the lack of details about allocation concealment.
- **Health-related quality of life:** only [Xu 2018](#) was at low risk of bias for the SGRQ and CAT. Most of the studies that reported the SGRQ were at some concern because of the lack of details about allocation concealment ([Berton 2015](#); [Saka 2021](#)), lack of details in the trial register ([Saka 2021](#)), and no information about whether adjusted analysis was planned in advance ([Berton 2015](#)). All the studies that reported CRQ were at high risk of bias (mainly because participants were not blinded) and some concern (lack of details about allocation concealment and issues with the reported results).
- **Inspiratory muscle strength:** only three studies were at low risk of bias (P_{lmax}; [Cutrim 2019](#); [Langer 2018](#); [Xu 2018](#)). All the remaining studies did not provide enough details about allocation concealment and studies at high risk of bias had issues with ITT analysis or missingness, or both.

Overall risk of bias

The main issues we found across the studies were the lack of details about allocation concealment (randomization process), lack of blinding (measurement of the outcome, although we considered using a sham IMT equal to blinding participants), and only journal articles were available (selection of the reported results).

Effects of interventions

See: [Summary of findings 1 Pulmonary rehabilitation plus inspiratory muscle training compared to pulmonary rehabilitation alone for people with chronic obstructive pulmonary disease](#); [Summary of findings 2 Inspiratory muscle training compared to control or sham for people with chronic obstructive pulmonary disease](#)

Comparison 1: pulmonary rehabilitation plus inspiratory muscle training versus pulmonary rehabilitation

See: [Summary of findings 1](#)

Primary outcomes

Dyspnea

Borg scale

Two studies used the Borg scale at isotime to evaluate dyspnea. Considering a MCID of -1 unit ([Ries 2005](#)), there was no improvement in dyspnea with an overall effect estimate (MD 0.19, 95% CI -0.42 to 0.79; $I^2 = 0\%$; 2 studies, 202 participants; [Analysis 1.1](#)). We judged only one study ([Charususin 2018](#)), to be at low risk of bias, which revealed a similar effect estimate. One study ([Sykes 2005](#)), reported greater improvement in dyspnea at "heavy load" in the IMT+ exercise group but did not report numerical data.

Using GRADE, we downgraded the certainty of the evidence for Borg by 1 point due to serious concerns regarding imprecision.

mMRC scale

We did not find any improvement in dyspnea using the mMRC scale (MD -0.12, 95% CI -0.39 to 0.14; $I^2 = 0\%$; 2 studies, 204 participants; [Analysis 1.2](#)). The MCID is estimated to be between -0.5 to -1 unit ([Araújo 2017](#); [Cazzola 2015](#)).

We downgraded the certainty of evidence to very low due to very serious concerns with risk of bias and serious concerns with imprecision.

BDI-TDI and MDP

There was no significant difference between the two arms in the studies that reported the BDI-TDI ([Schultz 2018](#)), and the MDP ([Beaumont 2015](#); [Beaumont 2018](#)).

Functional exercise capacity

6MWD

We pooled the 12 studies that reported the 6MWD test in a meta-analysis that showed no evidence of a difference between groups (MD 5.95, 95% CI -5.73 to 17.63; $I^2 = 61\%$; 12 studies, 1199 participants; [Analysis 1.3](#)). The mean and upper bounds of the 95% CI were lower than the MCID of 26 meters indicating that the mean change was not clinically relevant ([Puhan 2011](#)). [Sykes 2005](#) narratively reported a better 6MWD in the intervention group.

We had very low confidence in the results due to serious concerns with risk of bias and very serious concerns with inconsistency.

We conducted a subgroup analysis according to the duration of the intervention (short, medium and long-term interventions). The test for subgroup differences was not significant ($\text{Chi}^2 = 0.30$, $\text{df} = 2$ ($P = 0.86$), $I^2 = 0\%$, [Analysis 1.4](#)).

In the following subgroup analysis, we divided the studies according to their baseline P_{lmax} and the test for subgroup differences was not significant ($\text{Chi}^2 = 1.94$, $\text{df} = 1$ ($P = 0.16$), $I^2 = 48.3\%$, [Analysis 1.5](#)).

One study ([Wang 2017](#)), reported a subgroup analysis for intervention group participants with or without respiratory muscle weakness, with greater improvement in 6MWD for the weakened respiratory muscle group.

In sensitivity analysis, keeping just the studies at low risk of bias ([Beaumont 2018](#); [Charususin 2018](#); [Dellweg 2017](#)), increased the mean difference without exceeding the MCID (MD 8.90, 95% CI

-11.86 to 29.65; $I^2 = 80\%$; 4 studies, 379 participants). However, it decreased when switching to the fixed-effect model (MD 0.73, 95% CI -4.80 to 6.26, $I^2 = 61\%$).

12MWD

Three studies reported the 12MWD and showed a larger effect in the intervention group (MD 155.77 meters, 95% CI -84.53 to 396.08; $I^2 = 79\%$; 80 participants; [Analysis 1.6](#))

Wmax

Five studies reported Wmax and showed no difference between the groups (MD -1.01 watts, 95% CI -6.96 to 4.94; $I^2 = 25\%$; 326 participants; [Analysis 1.7](#)).

Exercise time (seconds)

Four studies reported exercise time (seconds) and also showed a larger effect in the intervention group (MD 58.62 seconds, 95% CI -25.09 to 142.32; $I^2 = 0\%$; 192 participants; [Analysis 1.8](#)). We did not include [Berry 1996](#) in the meta-analysis because they did not specify the level of exercise at which they measured exercise time. However, it is unclear to what extent these changes are clinically relevant.

Health-related quality of life (HRQoL)

SGRQ

We pooled seven trials that explored the effect of the intervention on HRQoL using the SGRQ, which showed no difference in total scores (MD 0.13, 95% CI -0.93 to 1.20; $I^2 = 0\%$; 908 participants; [Analysis 1.9](#)). This mean difference did not exceed the MCID threshold of -4 units ([Welling 2015](#)).

We downgraded the certainty of evidence by two levels due to very serious concerns with risk of bias.

Two studies that reported the three SGRQ domains showed no differences in symptoms (MD -2.33, 95% CI -6.28 to 1.62; [Analysis 1.9](#)), activity (MD 0.28, 95% CI -1.65 to 2.20) or impact (MD -1.63, 95% CI -5.38 to 2.11) scores.

In sensitivity analysis, effect estimates remained unchanged with a fixed-effect model. We could not explore the summary effect of low risk of bias studies because all the studies were at high risk of bias and some concerns.

CRQ

There were no differences in CRQ domain scores ([Analysis 1.10](#)): three RCTs reported 'Dyspnea' (MD -0.30, 95% CI -1.90 to 1.29) and 'Fatigue' (MD 0.28, 95% CI -0.76 to 1.31). Two RCTs reported 'Emotion' (MD -0.63, 95% CI -2.53 to 1.26) and 'Mastery' (MD -0.05, 95% CI -1.18 to 1.08). We used the generic inverse variance method to pool the results because [Charususin 2018](#) showed a different adjusted mean difference compared to non-adjusted analysis. None of the differences exceeded the MCID threshold of +0.5 units ([Alma 2018](#)). One abstract ([Sykes 2005](#)), narratively reported an improved CRQ score in the intervention group.

CAT

Two studies that reported the CAT scale showed no difference between groups (MD 0.13, 95% CI -0.80 to 1.06; $I^2 = 50\%$; 657 participants; [Analysis 1.11](#)) and the difference did not exceed the threshold of clinical significance, MCID of -1.6 units ([American](#)

[Thoracic Society](#)). We considered the certainty of evidence to be very low due to very serious concerns with risk of bias and serious concerns with inconsistency.

Secondary outcomes

Inspiratory muscle strength: P_{lmax} (cmH₂O)

Aggregated data from 17 studies showed higher inspiratory muscle strength in the intervention group (MD 11.46, 95% CI 7.42 to 15.15; $I^2 = 84\%$; 1329 participants; [Analysis 1.12](#)), but this did not exceed the MCID of 17.2 ([Iwakura 2020](#)). The statistical heterogeneity is due to the studies that reported change from baseline and had narrow confidence intervals.

We downgraded the certainty of evidence by one level due to serious concerns with risk of bias.

We performed a subgroup analysis according to the duration of intervention, which did not reveal a difference between subgroups ($\text{Chi}^2 = 0.05$, $\text{df} = 2$ ($P = 0.98$), $I^2 = 0\%$; [Analysis 1.13](#)).

We carried out subgroup analysis between studies with participants with and without respiratory muscle weakness. The test for subgroup differences was not significant ($\text{Chi}^2 = 1.72$, $\text{df} = 1$ ($P = 0.19$), $I^2 = 41.8\%$; [Analysis 1.14](#)).

[Wang 2017](#) divided the participants of the intervention group according to the state of their respiratory muscles, and they reported a larger change in P_{lmax} with the normal respiratory muscle group. [Sykes 2005](#) reported a higher P_{lmax} narratively in the intervention group.

Sensitivity analysis showed minimal impact of fixed-effect models on the synthesised results (MD 10.53, 95% CI 9.25 to 11.81, $I^2 = 84\%$). Retaining studies at low risk of bias ([Beaumont 2018](#); [Charususin 2018](#); [Dellweg 2017](#); [Fanfa Bordin 2020](#); [Schultz 2018](#); [Tounsi 2021](#)), also had minimal impact (MD 13.43, 95% CI 11.81 to 15.04; $I^2 = 90\%$; 1008 participants).

Laboratory exercise test: VO_{2peak}

We combined five studies showing no additional effect of the intervention on VO_{2peak} (MD -0.01 L/min, 95% CI -0.05 to 0.03; $I^2 = 0\%$; 313 participants; [Analysis 1.15](#)). [Sykes 2005](#) reported no difference between the two groups, but without numerical data.

Respiratory muscle endurance pressure (P_{thmax}) (cmH₂O)

Two studies reported P_{thmax}, and we pooled them in an SMD meta-analysis because they used different techniques to measure the outcome. We got an overall effect estimate (SMD 1.22 cmH₂O, 95% CI -0.18 to 2.66; $I^2 = 80\%$; 52 participants; [Analysis 1.16](#)), which suggests a large effect according to Cohen's d effect size.

Respiratory muscle endurance time: T_{lim} (seconds)

We performed separate analyses according to the nature of the endurance test. We pooled three studies that measured the outcome through sustained ventilation according to %P_{lmax} (MD 84.62, 95% CI -50.77 to 220.02; 236 participants; [Analysis 1.17](#)), and two studies that measured it according to %MVV (MD 477.69, 95% CI 215.43 to 739.94; 51 participants; [Analysis 1.18](#)). For both methods of measurement, we found a larger effect in the intervention group.

Maximal voluntary ventilation (MVV)

We pooled in an SMD meta-analysis two studies that reported MVV in L/min (Berry 1996; Wang 2017), and one study in %Pred (Paneroni 2018). We got an overall effect estimate (SMD 0.40, 95% CI -0.02 to 0.83; $I^2 = 4\%$; 93 participants; Analysis 1.19), which suggests a moderate effect according to Cohen's d effect size.

Respiratory function: forced expiratory volume at 1 second (FEV1)

We combined the studies according to whether they reported FEV1 in %Pred and in liters. We did not find a better effect of PR+IMT, showing an overall effect estimate (MD 0.77, 95% CI -1.72 to 3.26; $I^2 = 0\%$; 6 studies, 173 participants; Analysis 1.20) and (MD 0.04, 95% CI -0.04 to 0.13; $I^2 = 56\%$; 6 studies, 889 participants; Analysis 1.21) respectively.

Respiratory function: residual volume

One study reported this outcome (Charususin 2018) and did not show a difference between the two groups.

Adverse events

One abstract (Masanga 2011) reported some adverse effects that were considered "minor and self-limited": headache (six participants), jaw pain (six participants), neck pain (six participants), back pain (four participants), abdominal pain (two participants), cough (one participant), blood-streaked sputum (one participant), shoulder pain (one participant) and chest pain (one participant).

Comparison 2: inspiratory muscle training versus control/sham

See: Summary of findings 2

Primary outcomes

Dyspnea

Borg scale

Six studies used the Borg scale to assess dyspnea at isotime, and we entered them into a meta-analysis. Breathlessness was lower with the intervention and the mean difference was close to the MCID of -1 unit (the lower limit of the 95% CI exceeded it), though results are imprecise (MD -0.94, 95% CI -1.36 to -0.51; $I^2 = 0\%$; 6 studies, 144 participants; Analysis 2.1). Only one study was at low risk of bias.

We downgraded the certainty by three levels due to very serious concerns with risk of bias and serious concerns with imprecision.

BDI-TDI

Eight studies (nine arms) used BDI-TDI to measure dyspnea (Analysis 2.2). Three studies (four arms) reported the TDI 'Change in Functional impairment' (MD 0.88, 95% CI 0.51 to 1.25), TDI 'Change in Magnitude of task' (MD 0.73, 95% CI 0.35 to 1.12), and TDI 'Change in Magnitude of effort' (MD 0.86, 95% CI 0.42 to 1.30), showing no potential improvement in dyspnea, according to an MCID of +1 unit (Mahler 2005). Eight studies (nine arms) reported the TDI 'Focal score' and revealed a greater effect with IMT (MD 2.98, 95% CI 2.07 to 3.89; $I^2 = 65\%$; 238 participants).

We created a subgroup analysis of the TDI 'Focal score' according to studies judged with or without respiratory muscle weakness. The

test for subgroup differences was not significant ($\text{Chi}^2 = 2.55$, $\text{df} = 1$ ($P = 0.11$), $I^2 = 60.8\%$; Analysis 2.3).

In sensitivity analysis, the overall effect estimates remained unchanged when switching to the fixed-effect model in the first three subgroups, while it increased to (MD 4.04, 95% CI 3.70 to 4.39, $I^2 = 65\%$) with 'Focal score'. This effect mainly resulted from (Sanchez Riera 2001), which got the greatest weight.

Only one study was at low risk of bias (Langer 2018), and we considered the certainty of evidence to be very low due to very serious concerns with risk of bias and serious concerns with imprecision.

mMRC

Four studies reported the mMRC score, and the overall effect estimate revealed a possible improvement in dyspnea in the IMT group (MD -0.59, 95% CI -0.76 to -0.43; $I^2 = 17\%$; 150 participants; Analysis 2.4), considering an MCID between -0.5 and -1 unit. We considered the certainty of evidence to be low due to serious concerns with risk of bias and imprecision.

Functional exercise capacity

6MWD

We combined 16 studies (17 arms) that reported the 6MWD, showing a better effect with the IMT compared to control/sham (MD 35.71, 95% CI 25.68 to 45.74; $I^2 = 16\%$; 501 participants; Analysis 2.5).

We judged the certainty of evidence to be moderate due to serious concerns with risk of bias.

In the first subgroup analysis, we divided the studies according to the duration of the intervention. Only one study (Saher 2021), had a follow-up of fewer than four weeks. The test for subgroup differences was not significant ($\text{Chi}^2 = 0.37$, $\text{df} = 2$ ($P = 0.83$), $I^2 = 0\%$; Analysis 2.6).

In the following subgroup analysis, we divided the studies according to the mean baseline P_{lmax}. The test for subgroup differences was not significant ($\text{Chi}^2 = 0.17$, $\text{df} = 1$ ($P = 0.68$), $I^2 = 0\%$; Analysis 2.7). One trial (Xu 2018), performed a subgroup analysis within study data using the same cut-off.

In sensitivity analysis, keeping just the studies at low risk of bias (Cutrim 2019; Xu 2018), revealed a larger effect, standing at around twice our MCID (MD 49.13, 95% CI -27.62 to 125.88, $I^2 = 83\%$). However, we should note that its confidence interval exceeded the line of no effect. The overall effect estimate remained nearly the same when switching to the fixed-effect model.

12MWD and W_{max}

We did not find an additional effect of IMT when we pooled the studies that used the 12MWD and W_{max}, revealing, respectively, an overall effect estimate (MD -33.31, 95% CI -158.10 to 91.48; $I^2 = 59\%$; 3 studies, 101 participants = 101; Analysis 2.8) and (MD 0.66, 95% CI -6.44 to 7.76; $I^2 = 42\%$; 7 studies, 206 participants; Analysis 2.9). However, for the 12MWD, we noticed large differences in baseline data between the two groups of Preusser 1994, which appeared significant when testing it with the RevMan calculator ($P = 0.03$). We removed that study in a sensitivity analysis, and we got a positive overall effect (MD 12.76, 95% CI -65.71 to 91.23, $I^2 = 0\%$).

Exercise time

Five studies (six arms) reported exercise time, but we did not pool the studies because they used different measurement methods (Analysis 2.10). Globally, their results were consistent and showed a trend of a greater effect in the IMT group compared to the control/sham group.

SWT

Three studies used the SWT, and we included two trials in our primary analysis. We did not find an additional effect in the IMT group (MD -7.45 meters, 95% CI -92.74 to 77.83; Analysis 2.11).

Health-related quality of life (HRQoL)

SGRQ

Two studies (Berton 2015; Saka 2021), reported the items 'Symptoms' (MD -2.10, 95% CI -3.50 to -0.71), 'Activity' (MD -9.86, 95% CI -15.08 to -4.63) and 'Impact' (MD -6.06, 95% CI -13.76 to 1.65). Six studies reported the 'total score' of the SGRQ, showing a larger effect in the IMT group (MD -3.85, 95% CI -8.18 to 0.48; $I^2 = 66%$; 182 participants; Analysis 2.12). The lower limit of the 95% CI exceeded the MCID of -4 units. The statistical heterogeneity is due to the effect of Saka 2021. Only one study was at low risk of bias (Xu 2018), and it has a similar effect estimate to the overall. The overall effect estimate increased to MD -5.11 (95% CI -6.81 to -3.40, $I^2 = 66%$), when switching to the fixed-effect model.

We judged the certainty of the evidence as 'very low' due to very serious concerns with risk of bias and imprecision.

CRQ

Four studies (5 arms) reported the four items of the CRQ, while one trial (Larson 1999), reported only dyspnea and fatigue. We divided the items into subgroups. The RCTs showed consistent results across the items, and there was an improvement (considering an MCID of +0.5 unit) in 'Dyspnea' (MD 1.63, 95% CI 0.23 to 3.03; Analysis 2.13), 'Fatigue' (MD 1.32, 95% CI 0.08 to 2.55), 'Emotion' (MD 2.64, 95% CI 0.82 to 4.46), and 'Mastery' (MD 1.57, 95% CI 0.07 to 3.06). None of the included studies was at low risk of bias.

CAT

Two trials reported CAT, revealing an overall effect estimate (MD -2.97, 95% CI -3.85 to -2.10; participants = 86; $I^2 = 0%$; Analysis 2.14). We downgraded the certainty of evidence by one level due to serious concerns with imprecision.

SF-36 and CCQ

In the studies that used SF-36 (Chuang 2017; Nikolettou 2016), and CCQ (Leelarungrayub 2017), there was a better effect in the IMT group in some domains of the scales.

Secondary outcomes

Inspiratory muscle strength: P_Imax (cmH₂O)

P_Imax

32 RCTs (34 arms) measured P_Imax. The upper limit of the 95% CI of summary effect between IMT and control/sham exceeded the MCID of 17.2 cmH₂O (MD 14.57 cmH₂O, 95% CI 9.85 to 19.29; $I^2 = 89%$; 916 participants; Analysis 2.15).

We downgraded the certainty of evidence by two levels due to serious concerns with risk of bias and a strongly suspected publication bias.

We divided the studies according to the training duration (Analysis 2.16). The test for subgroup differences was not significant ($\text{Chi}^2 = 0.67$, $\text{df} = 2$ ($P = 0.72$), $I^2 = 0%$).

We carried out subgroup analysis between studies with participants with and without respiratory muscle weakness. The test for subgroup differences was not significant ($\text{Chi}^2 = 0.34$, $\text{df} = 1$ ($P = 0.56$), $I^2 = 0%$; Analysis 2.17).

We also conducted a subgroup analysis of the measurement method. The test for subgroup differences was not significant ($\text{Chi}^2 = 0.93$, $\text{df} = 2$ ($P = 0.63$), $I^2 = 0%$; Analysis 2.18) between studies that measured P_Imax at residual volume, at functional residual capacity, and the studies that did not report the level of measurement.

In sensitivity analysis, keeping just the studies at low risk of bias (Cutrim 2019; Langer 2018; Xu 2018), showed a similar result (MD 12.79 cmH₂O, 95% CI 3.63 to 21.95, $I^2 = 55%$). Similarly, the overall effect estimate of the random-effects model did not differ from the fixed-effect model.

Laboratory exercise test (VO₂peak)

We combined 11 studies (12 arms) in an SMD meta-analysis, since they were reported in different units (L/min, mL/min, and mL/kg/min). We got an overall effect estimate suggesting a low to moderate effect according to Cohen's rule of thumb (SMD 0.31, 95% CI 0.05 to 0.57; $I^2 = 14%$; 286 participants; Analysis 2.19).

Respiratory muscle endurance pressure (P_{thmax}) (cmH₂O)

Eight RCTs used P_{thmax} in their respiratory endurance assessment, and they showed a larger overall effect in the IMT group (MD 9.71, 95% CI 4.93 to 14.50; $I^2 = 53%$; 179 participants; Analysis 2.20).

Respiratory muscle endurance time T_{lim} (seconds)

We pooled 10 studies (11 arms) that reported T_{lim}, and they showed a better improvement in the outcome in the IMT group (MD 270.57, 95% CI 182.44 to 358.71; $I^2 = 63%$; 260 participants; Analysis 2.21). One trial (Heijdra 1996) considered it as the time to reach P_{thmax} (not included in the analysis).

Maximal voluntary ventilation (MVV)

Two studies reported MVV. We pooled them in an SMD meta-analysis because Belman 1988 did not report the unit. We got a summary effect suggesting a large effect according to Cohen's d effect size (SMD 0.99, 95% CI 0.28 to 1.69; $I^2 = 0%$; 36 participants; Analysis 2.22).

Respiratory function: forced expiratory volume at 1 second

We ran two analyses according to FEV₁ unit without finding a significant difference between IMT and control/sham in either unit. Ten studies (11 arms) reported FEV₁ in %Pred showing a difference of (MD 2.62 %predicted, 95% CI 0.20 to 5.04 Analysis 2.23), while 12 studies (13 arms) reported it in liters (MD 0.04 L, 95% CI -0.06 to 0.14 Analysis 2.24).

Respiratory function: residual volume

Hill 2006 reported residual volume in %Pred and liters, showing a better improvement in the IMT group, while Ramirez Sarmiento 2002 reported it in liters and did not show a difference.

Adverse events

None of the included studies reported a side effect of the intervention.

DISCUSSION

This review has summarised the available evidence of the effect of inspiratory muscle training, used alone or in combination with pulmonary rehabilitation, in people with COPD. We combined the results of the studies in a logical way, so that no data were lost, taking into account the variety of scales, respiratory measurements and training protocols.

Summary of main results

We included 55 RCTs in this review (including trials with more than two arms). Twenty-two studies (1446 participants) investigated the effect of PR with IMT compared to PR, while 37 studies (1021 participants) focused on the effect of IMT compared to control/sham. Three trials (Abedi Yekta 2019; Larson 1999; Majewska-Pulsakowska 2016), explored both comparisons (four arms).

Comparison 1: pulmonary rehabilitation plus inspiratory muscle training versus pulmonary rehabilitation

In our first comparison (PR+IMT vs PR), we did not find a significant improvement in the intervention group in dyspnea measured with the Borg scale at isotime, referring to an MCID of -1 unit (moderate-certainty evidence). We also did not find a potential effect with the mMRC scale, based on an MCID between -0.5 to -1 unit (very low-certainty evidence).

Studies assessed functional exercise capacity with four measurements: 6MWD, 12MWD, Wmax, and exercise time. We did not find an additional effect of combining PR and IMT with the 6MWD, which we considered our main measurement for this outcome. The overall effect estimate did not reach the MCID of 26 meters (low-certainty evidence). In subgroup analysis, we divided the studies according to the training duration and mean baseline PImax. The test for subgroup differences was not significant. In sensitivity analysis, three studies at low risk of bias showed a relatively greater treatment effect.

The overall effect estimate of the 12MWD and exercise time showed a larger effect in the intervention group. However, it remains unclear to what extent this difference is clinically relevant because we did not find an MCID for this outcome. The studies that measured Wmax did not reveal a difference between the two groups.

Seven studies investigated the effect of the intervention on HRQoL through the SGRQ. The overall effect estimate showed a positive effect in two domains ('Symptoms' and 'Impact'), without exceeding the MCID of -4 units. There was no difference between the two groups with 'Impact' and 'Total score' (low-certainty evidence).

In the CRQ, we only found a clinically relevant difference (-0.5 units) in the 'Emotion' domain. The other items ('Dyspnea', 'Fatigue' and

'Mastery') did not show a difference, noting that only three studies reported this scale.

Two studies reported the CAT scale, and the overall effect estimate failed to reveal a benefit based on an MCID of about -1.6 units (very low-certainty evidence).

The combination of PR and IMT might have a greater effect on inspiratory muscle strength (PImax) than PR alone, but our treatment effect failed to reach the MCID (moderate-certainty evidence). In subgroup analysis, we did not find a difference between different training durations nor between studies with participants with or without respiratory muscle weakness.

The overall effect estimate of VO₂peak, MVV, FEV1 (%Pred and L), and residual volume did not reveal a difference between the two interventions. For respiratory muscle endurance tests (T_{lim} and P_{thmax}), we discovered a more significant effect in the PR plus IMT. Nonetheless, the clinical relevance of these estimations remained doubtful since we did not find an MCID.

Comparison 2: inspiratory muscle training versus control/sham

In our second comparison, IMT versus control/sham, taking account of the Borg MCID (-1 unit), we found a trend of an improvement in dyspnea in the IMT group (only the lower limit of the 95% CI exceeded the MCID) when the scale was measured at submaximal exercise capacity (very low-certainty evidence).

The studies that used BDI-TDI did not reveal a clinically meaningful change (MCID +1 unit) after IMT in the three items of the scale (Functional impairment, Magnitude of task, Magnitude of effort). On the other hand, we found a larger effect with 'Focal score' of the TDI (very low-certainty evidence). The difference between the items might be explained by the fact that more studies reported Focal score (N = 8) than the other items (N = 3). In subgroup analysis, we did not find a difference between studies with participants judged with and without respiratory muscle weakness.

The overall effect estimate of the mMRC showed a possible effect with IMT that exceeded the MCID of -0.5 units (low-certainty evidence).

Turning to functional exercise capacity, we combined 16 studies (17 arms) that used the 6MWD, showing a larger effect exceeding the MCID: 26 meters (moderate-certainty evidence). Following the same subgroup analysis of our first comparison, we did not find a difference between different training durations and between studies with participants judged with and without respiratory muscle weakness. In sensitivity analysis, studies at low risk of bias (N = 2) showed a greater treatment effect.

Studies also used other measurements to assess exercise capacity, and the overall estimation of the 12MWD and Wmax did not show an advantageous effect of IMT. We did not combine the studies that reported exercise time because the methods of measurement were inconsistent. Still, all the effect estimates were on the right-hand side of the line of no effect with different degrees of positive effect.

We pooled separately three scales (SGRQ, CRQ and CAT) that assessed the HRQoL in a meta-analysis. All the scales revealed a better improvement in life quality with IMT, but unlike CRQ and

CAT, only the lower limit of the 95% CI of SGRQ exceeded the MCID (-4 units; very low-certainty evidence). We noticed a larger effect estimate favouring the intervention in some items when other scales were used (CAT, SF-36, CCQ).

Plmax was the main secondary outcome in our second comparison. There was a trend to a greater effect with IMT, considering an MCID of 17.2 cmH₂O (low-certainty evidence). We did not find a difference between medium-term and long-term training nor between studies with participants with or without respiratory muscle weakness. Measuring the outcome at residual volume or functional residual capacity showed similar results. We got similar results with studies at low risk of bias ($N = 3$).

We combined the studies that reported VO₂peak in an SMD meta-analysis. According to Cohen's rule of thumb, we considered the overall effect estimate between low to moderate.

The studies that reported respiratory muscle endurance tests (P_{thmax} , T_{lim} , MVV) showed a beneficial effect favouring IMT. However, we could not judge the clinical relevance of this effect since we did not find the MCID of these outcomes. The overall effect estimate of FEV1 (%Pred, L) did not reveal a potential improvement with IMT compared to control/sham.

Overall completeness and applicability of evidence

This review was conducted on patients with stable COPD, aged 44 years and over. It explored a wide range of participants and included all RCTs regardless of COPD stages and training protocols (duration, load, supervision, devices, follow-up, measurements). Most studies, except [Ahmad 2013](#) excluded participants who were free from exacerbation for a couple of weeks preceding the trial and were not hospitalised. Therefore, the results of this review may not be applicable to IMT conducted just after an acute exacerbation. We also noticed a shift over the years from using resistive devices to threshold devices. We excluded RCTs that used resistive trainers without controlling the breathing pattern because participants could adapt a non-fatiguing respiratory pattern without exposing the respiratory muscles to the workload. Currently, with technological development, there has been increasing use of electronic devices that allow remote monitoring and accurate adjustment of the training load.

Studies that looked at the combined effect of PR plus IMT included people with mild to very severe COPD. One study ([Dellweg 2017](#)), performed the intervention on hypercapnic patients who remained dependent on non-invasive ventilation after prolonged weaning. Two studies ([Beaumont 2015](#); [Charususin 2018](#)), specifically trained participants with reduced Plmax, and two trials did subgroup analysis within study data according to the state of respiratory muscles ($Plmax \leq$ or > 60 cmH₂O) ([Beaumont 2018](#); [Wang 2017](#)). Participants in [Magadle 2007](#) and [Weiner 2000](#) received respectively three months and six weeks of PR before the start of IMT, while most of the other studies excluded this category of patients. These two studies had no impact on the overall effect size when we removed them in a sensitivity analysis. Apart from [De Farias 2019](#), [Mador 2005](#) and [Paneroni 2018](#), which conducted endurance training, all the studies performed strength training with various loads and different training durations.

Furthermore, there was a wide range of PR protocols, starting from just breathing exercises and respiratory drainage to a mix

of interventions (e.g. cycling, treadmill, muscle strengthening, therapeutic patient education). To the best of our knowledge, the current guidelines have not yet established a consensus for an optimal duration or components of a rehabilitation program. This might be explained by the multiple factors that could interfere with it, such as participants' motivation and financial and logistic resources.

In our second comparison, we were also exposed to a diversity of training protocols in the studies that compared IMT with control/sham. More than half of the included RCTs used a sham IMT, and only two ([Bavarsad 2015](#); [Dacha 2019](#)), had some participants with mild COPD. One trial ([Saher 2021](#)), focused on patients receiving non-invasive ventilation as part of COPD management. Almost all the trials excluded participants who had exacerbations before starting the trial, and two RCTs worked on patients with limited functional performance ([Covey 2001](#); [ZhouL 2016](#)). Two studies had a long intervention duration ([Beckerman 2005](#); [Kim 1993](#)), equal to one year and six months respectively; two studies trained their participants with a load up to 100% of their Plmax ([Chuang 2017](#); [Hill 2006](#)); two trials conducted endurance IMT ([Koppers 2006](#); [Scherer 2000](#)), and one trial performed both endurance and strength exercises ([Petrovic 2012](#)). The remaining studies had close characteristics (i.e. strength training, similar loads).

This review did not compare threshold IMT with normocapnic hyperpnea training (endurance IMT) nor compare training loads. We thought it would be best to work on these objectives in a separate review.

Overall, this review included all kinds of participants and interventions without preferences. However, our results might not be applicable for hospitalised patients following an exacerbation or patients with severe COPD who require long-term oxygen therapy.

Quality of the evidence

Using the GRADE approach, we assessed the certainty of the evidence of five outcomes: dyspnea (Borg, mMRC, BDI-TDI), functional exercise capacity (6MWD), HRQoL (SGRQ, CAT), and Plmax. When the number of RCTs exceeded 10, we created a funnel plot to investigate publication bias.

For both comparisons, we considered the certainty of evidence to be from very low to moderate across all the outcomes (see [Summary of findings 1](#); [Summary of findings 2](#)).

Comparison 1: pulmonary rehabilitation plus inspiratory muscle training versus pulmonary rehabilitation

In our first comparison (PR+IMT versus PR), we downgraded the certainty of evidence by one level for the Borg scale measured at submaximal exercise capacity because of serious concerns about risk of bias related to blinding participants.


We downgraded the certainty of evidence of the mMRC to very low due to very serious concerns with risk of bias (all the trials are at high risk of bias) and serious concerns with imprecision (the sample size is less than 400, rule of thumb).

We considered the certainty of evidence of the 6MWD to be very low due to serious concerns with risk of bias (studies at low risk of bias had different effect estimate from the overall effect estimate) and very serious concerns with inconsistency, although

we explained part of the heterogeneity by the difference in training durations. Indeed, there was substantial statistical heterogeneity, and the effect estimates were wide on both sides of the line of

no effect, making the benefit of the intervention doubtful. The funnel plot was symmetrical (Figure 2), so publication bias was improbable.

Figure 2. Funnel plot of comparison 7, PR+IMT vs PR, outcome: 7.3 functional exercise capacity: 6-minute walk distance (6MWD)




We judged the quality of evidence of the SGRQ (total score) as low because of very serious concerns with risk of bias. In fact, none of the studies was at low risk of bias; the main issue was the lack of blinding of participants (outcome measurement).

We considered the certainty of evidence of CAT to be very low due to very serious concerns with risk of bias (most of the evidence is from studies at high risk of bias and some concerns, lack of blinding) and serious concerns with inconsistency (considerable statistical heterogeneity, and confidence intervals do not overlap).

We downgraded the certainty of evidence of PImax by one level due to serious concerns with risk of bias (most of the evidence is from studies at high risk of bias and some concerns). We did not consider statistical heterogeneity because all the effect estimates were on one side showing a benefit). The funnel plot did not raise suspicions of publication bias (Figure 3). As a result, we considered the certainty of the evidence as moderate.

Figure 3. Funnel plot of comparison 7, PR+IMT vs PR, outcome: 7.12 respiratory muscle strength: PImax



Comparison 2: inspiratory muscle training versus control/sham


In our second comparison (IMT versus control/sham), we downgraded the certainty of evidence of Borg measured at submaximal exercise capacity to very low due to very serious concerns with risk of bias (most of the studies were at high risk or some concerns, with different effect estimates compared to low risk of bias studies, issues with blinding) and serious concerns with imprecision (small sample size less than 400 participants, rule of thumb).

For the same reasons explained above, we downgraded the certainty of evidence for dyspnea assessed with BDI-TDI by three levels.

We considered the certainty of evidence of mMRC to be low due to serious concerns with risk of bias (most of the evidence is from studies at high risk of bias and some concerns) and imprecision (sample size less than 400 participants, rule of thumb).

We judged the quality of evidence of the 6MWD as moderate due to serious concerns with risk of bias (most of the evidence is from studies at high risk of bias and with some concerns). We did not downgrade inconsistency, despite the substantial statistical heterogeneity, because the effect estimates were on one side of the line of no effect. So we were more confident about the direction of the effect. The funnel plot raised some concerns about publication bias (Figure 4), without downgrading the evidence.

Figure 4. Funnel plot of comparison 8, IMT vs control/sham, outcome: 8.8 functional exercise capacity: 6-minute walk distance (6MWD)




We downgraded the certainty of evidence of the SGRQ by three levels because of very serious concerns with risk of bias (studies with greater weight were at high risk of bias and some concerns, showing different effect estimates from studies at low risk of bias) and very serious concerns with imprecision (sample size less than 400 participants, rule of thumb; the 95% CI includes benefit and harm).

We considered the certainty of evidence of CAT to be moderate due to serious concerns with imprecision (sample size less than 400, rule of thumb).

We judged the certainty of evidence of P1max as low due to serious concerns with risk of bias (most of the evidence is from studies at high risk of bias and with some concerns) and a strongly suspected publication bias (Figure 5).

Figure 5.



Potential biases in the review process

The review is based on a published protocol (Ammous 2020). We discuss deviations from the protocol in the Differences between protocol and review section.

One of the main issues with the included studies in this review is the risk of bias, using RoB 2, which is outcome-dependent. Only six out of 22 studies were at low risk of bias for the first comparison, while only three out of 37 studies were at low risk for the second comparison. Across all the outcomes, the major problems were lack of detail about allocation concealment, some issues with missing data, participants being aware of their intervention (for the participant-reported outcome that involves judgments), and not publishing a protocol or listing the study in a trial register before launching the studies. Sometimes studies at high risk of bias tended to show a larger effect than the others, which reduced the confidence in our results.

One trial provided IMT for a short period of time (two weeks; Saher 2021), and participants received non-invasive ventilation throughout the trial.

From all our included studies, only one abstract (Masanga 2011), reported adverse events. Although it is unlikely that IMT may be associated with harm, we have some concerns due to the variety of side effects reported by Masanga 2011, and because the included studies did not discuss them. That is to say, one of the reasons for discontinuing the trial was the inability of participants to continue

the intervention. But the studies did not discuss the reasons that made patients take that decision.

Moreover, when dividing the studies according to mean baseline PImax to classify participants with or without respiratory muscle weakness, we assumed that the proportion of participants not belonging to their assigned subgroup was balanced across the studies. In other words, getting individual data and conducting a subgroup analysis within study data was impossible. So, we assumed this variation was balanced between the studies, considering that PImax follows a normal distribution with fewer outliers.

Agreements and disagreements with other studies or reviews

Ten published reviews that explored the effect of IMT on COPD have been published over the years. The first one was published in the 90s (Smith 1992). This review included 17 trials in which the participants had chronic airflow limitation without specifying the types of the diseases. They summarised the effect estimate with SMD meta-analysis and interpreted it according to Cohen’s d effect size. They found a small to moderate effect favouring IMT for MVV, PImax and laboratory exercise tests. However, this review included trials that run resistance training without controlling the breathing pattern.

Ten years later, another systematic review was published (Lötters 2002), including 15 RCTs. Unlike our review, this study pooled altogether in an SMD meta-analysis RCTs that looked at the effect

of IMT as a stand-alone intervention and when associated with PR. Then, they conducted a subgroup analysis to investigate the additional effect of PR. They found that IMT alone might improve dyspnea and respiratory muscle strength and endurance. When combined with PR, a beneficial effect was seen only in participants with respiratory muscle weakness.

The following review (16 studies) was conducted by [Crowe 2005](#), followed by an update with 18 studies ([O'Brien 2008](#)), in which they compared IMT separately with different types of interventions (exercises, breathing techniques, education, PR). The authors concluded, based on a study-level analysis, that IMT might improve respiratory muscle strength. However, there was less confidence with dyspnea and HRQoL. In our review, we considered the participants who underwent therapeutic patient education or one type of breathing exercises as control, because the aim was not to compare IMT with another intervention but to create a psychological effect for patients not receiving IMT. We excluded the studies that focused on comparing IMT with breathing exercises. Similarly, the same authors worked on another systematic review ([Geddes 2005](#)), and then they updated it ([Geddes 2008](#)). These reviews compared IMT with no intervention, low versus high IMT, and two different modes of IMT, which is not the purpose of our review.

A narrative review ([Shoemaker 2009](#)), reported that IMT, in comparison with sham IMT or no intervention, might improve dyspnea, HRQoL and PImax. Following that, [Gosselink 2011](#) showed similar results to [Lötters 2002](#). The authors included 32 RCTs, and found a possible effect of IMT in improving dyspnea, functional exercise capacity, HRQoL, PImax and respiratory muscle endurance. In subgroup analysis, they reported a benefit of IMT when associated with PR only in participants with respiratory muscle weakness. They also found a larger effect of strength training compared to endurance training.

In a subsequent study, [Beaumont 2018a](#) included 37 trials in a meta-analysis showing an improvement in dyspnea, HRQoL, functional exercise capacity and PImax after threshold IMT. However, they did not find a difference when combining IMT with PR. The last meta-analysis ([Figueiredo 2020](#)), compared IMT alone or associated with other interventions with control, sham, or other interventions. The authors did not find an improvement in dyspnea, HRQoL, or a larger effect in participants with respiratory muscle weakness. They showed a trend of a larger effect with higher training loads.

Our systematic review is the first to have excluded trials that used resistive trainers without controlling the breathing pattern. We are also the first that have excluded the trials that did not measure Borg at isotime. Generally, our results were consistent with past systematic reviews regarding no effect when adding IMT to PR and a possible benefit with IMT as a stand-alone intervention. However, we could not conclude that there is a better effect in participants with respiratory muscle weakness. A possible explanation for this finding includes the fact that most studies worked on participants without respiratory muscle weakness.

AUTHORS' CONCLUSIONS

Implications for practice

When associated with pulmonary rehabilitation (PR), inspiratory muscle training (IMT) may not have an additional benefit on dyspnea, functional exercise capacity and health-related quality of life. There was an increase in inspiratory muscle strength and endurance, but this difference was not clinically meaningful.

IMT may improve dyspnea, functional exercise capacity (mainly the 6MWD) and health-related quality of life, compared to sham or no intervention. There was also an increase in inspiratory muscle strength and endurance, but judging the clinical significance of these outcomes was challenging due to a non-standardised minimal clinically important difference.

For both interventions, we could not conclude that there was a possible larger effect in participants with respiratory muscle weakness and with longer durations of training.

Implications for research

Overall, we did not detect any potential advantage of combining PR and IMT, which was consistent with past systematic reviews. However, it is still unclear if this intervention is more beneficial in participants with respiratory muscle weakness, and future research may focus more on this group of participants.

We think there is enough evidence for the effect of IMT alone since our results are consistent with past systematic reviews. However, all published trials had a small sample size. So we believe future trials should increase the number of participants, investigate the possible larger effect on participants with respiratory muscle weakness and compare different IMT protocols.

IMT may be a starter intervention for patients unable to undergo PR (e.g. severe COPD, logistical or financial issues). Still, this should be explored further in clinical trials.

For both comparisons, we highly suggest providing a sham IMT for the control group. Although a sham IMT might not influence the results of an observer-reported outcome such as the 6-minute walk distance, we strongly believe that participants who know they are undergoing an intervention may overestimate the effect measured with participant-reported outcomes that involve judgments. Furthermore, future research may consider training with higher inspiratory flow rate and training at high lung volume (closer to total lung capacity) to improve performance in hyperinflated COPD patients.

ACKNOWLEDGEMENTS

We acknowledge Dr. Rujan Shrestha, Medical Officer, Nepal and Dr Karla Elizabeth Duque Jacome, Ecuador for their translating support.

We thank the staff of Cochrane Airways, especially Emma Jackson (Managing Editor), Emma Dennett (Deputy Co-ordinating Editor) and Elizabeth Stovold (Information Specialist) for their support.

The authors and Cochrane Airways Editorial Team are grateful to the following peer and consumer reviewers for their time and comments: Julia Robertson (Australia), Lissa M Spencer (Australia) and Elena Gimeno-Santos (Spain).

The [Background](#) and [Methods](#) sections of this review are based on a standard template used by Cochrane Airways.

This project was supported by the National Institute for Health and Care Research (NIHR), via Cochrane Infrastructure funding to

Cochrane Airways. The views and opinions expressed here are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health and Social Care.

REFERENCES
References to studies included in this review
Abedi Yekta 2019 {published data only}

Abedi Yekta AH, Poursaeed Esfahani M, Salehi S, Hassabi M, Khosravi S, Kharabian S, et al. Assessment of the effects of inspiratory muscle training (IMT) and aerobic training on the quality of life of patients with chronic obstructive pulmonary disease. *Tanaffus* 2019;**18**(3):223-9.

Bavarsad 2015 {published data only}

Bavarsad MB, Shariati A, Eidani E, Latifi M. The effect of home-based inspiratory muscle training on exercise capacity, exertional dyspnea and pulmonary function in COPD patients. *Iranian Journal of Nursing and Midwifery Research* 2015;**20**(5):613-18. [DOI: [10.4103/1735-9066.164588](https://doi.org/10.4103/1735-9066.164588)]

Beaumont 2015 {published and unpublished data}

* Beaumont M, Mialon P, Le Ber-Moy C, Lochon C, Peran L, Pichon R, et al. Inspiratory muscle training during pulmonary rehabilitation in chronic obstructive pulmonary disease: a randomized trial. *Chronic Respiratory Disease* 2015;**12**(4):305-12. [DOI: [10.1177/1479972315594625](https://doi.org/10.1177/1479972315594625)]

NCT01545011. Effects of inspiratory muscle training on dyspnea in subjects with chronic obstructive pulmonary disease (IMT) [Effects of inspiratory muscle training combined with a pulmonary rehabilitation program versus a program of pulmonary rehabilitation alone on dyspnea: a randomized trial]. clinicaltrials.gov/show/study/NCT01545011 (first received 6 March 2012).

Beaumont 2018 {published data only}

* Beaumont M, Mialon P, Le Ber C, Le Mevel P, Peran L, Meurisse O, et al. Effects of inspiratory muscle training on dyspnoea in severe COPD patients during pulmonary rehabilitation: controlled randomised trial. *European Respiratory Journal* 2018;**51**(1):01. [DOI: [10.1183/13993003.01107-2017](https://doi.org/10.1183/13993003.01107-2017)]

Beaumont M, Mialon P, Le Ber C, Le Mevel P, Peran L, Meurisse O, et al. Effects of inspiratory muscle training on dyspnea in severe COPD patients during pulmonary rehabilitation: controlled randomized trial. *European Respiratory Journal* 2017;**50**(suppl 61):OA2924. [DOI: [10.1183/1393003.congress-2017.OA2924](https://doi.org/10.1183/1393003.congress-2017.OA2924)]

NCT02074813. Inspiratory muscle training during pulmonary rehabilitation in COPD (EMI II) [Effects of inspiratory muscle training (IMT) on dyspnea in COPD during pulmonary rehabilitation: randomized controlled trial]. clinicaltrials.gov/show/study/NCT02074813 (first received 28 February 2014).

Beckerman 2005 {published data only}

* Beckerman M, Magadle R, Weiner M, Weiner P. The effects of 1 year of specific inspiratory muscle training in patients with COPD. *Chest* 2005;**128**(5):3177-82. [DOI: [10.1378/chest.128.5.3177](https://doi.org/10.1378/chest.128.5.3177)]

Beckermann M, Weiner M, Weiner P. The effects of one year specific inspiratory muscle training in patients with COPD. In:

European Respiratory Journal. Vol. 26 Suppl 49. 2005:Abstract No. 539.

Weiner P, Beckerman M, Magadle R, Berar-Yanay N. Maintenance of inspiratory muscle training in COPD patients: one year follow up [Abstract]. In: *European Respiratory Journal*. Vol. 22. 2003:P3438.

Weiner P, Magadle R, Beckerman M, Weiner M, Berar-Yanay N. Maintenance of inspiratory muscle training in COPD patients: one year follow-up. *European Respiratory Journal* 2004;**23**(1):61-5. [DOI: [10.1183/09031936.03.00059503](https://doi.org/10.1183/09031936.03.00059503)]

Belman 1988 {published data only}

Belman MJ, Shadmehr R. Targeted resistive ventilatory muscle training in chronic obstructive pulmonary disease. *Journal of Applied Physiology* 1988;**65**(6):2726-35. [DOI: [10.1152/jappl.1988.65.6.2726](https://doi.org/10.1152/jappl.1988.65.6.2726)]

Berry 1996 {published data only}

Berry MJ, Adair NE, Sevinsky KS, Quinby A, Lever, HM. Inspiratory muscle training and whole-body reconditioning in chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine* 1996;**153**(6):1812-16. [DOI: [10.1164/ajrccm.153.6.8665039](https://doi.org/10.1164/ajrccm.153.6.8665039)]

Berton 2015 {published data only}

Axmann de Castro M. Efeito do treinamento muscular inspiratório sobre a dispnéia, tolerância ao exercício e metaborreflexo da musculatura ventilatória em pacientes com doença pulmonar obstrutiva crônica [Masters thesis]. Porto Alegre: Federal University of Rio Grande do Sul (UFRGS), 2015. [<https://www.lume.ufrgs.br/bitstream/handle/10183/119402/000969567.pdf?sequence=1>]

Berton DC, Castro M, Frohlich LF, Castilho M, Dorneles R, Chiappa G, et al. Effects of inspiratory muscle training on leg blood flow and exercise tolerance in COPD. *European Respiratory Journal* 2015;**46** Suppl 59:PA2250.

* Castro MA, Fröhlich LF, Chiappa GR, Knorst MM, Neder JA, Berton DC, et al. Improvement in exercise capacity after inspiratory muscle training is related to increased calf blood flow during inspiratory load in COPD. *SM Journal of Pulmonary Medicine* 2016;**2**(1):1013.

Bustamante 2007 {published data only}

Bustamante V, Gáldiz Iturri JB, Gorostiza Manterola A, Camino Buey J, Talayero Sebastián N, Peña Víctor S. Comparison of 2 methods for inspiratory muscle training in patients with chronic obstructive pulmonary disease [Comparación de 2 métodos de entrenamiento muscular inspiratorio en pacientes con EPOC]. *Archivos de Bronconeumología* 2007;**43**(8):431-8. [DOI: [10.1157/13108782](https://doi.org/10.1157/13108782)]

Charususin 2018 {published and unpublished data}

* Charususin N, Gosselink R, Decramer M, Demeyer H, McConnell A, Saey D, et al. Randomised controlled trial of adjunctive inspiratory muscle training for patients

with COPD. *Thorax* 2018;**73**(10):942-50. [DOI: [10.1136/thoraxjnl-2017-211417](https://doi.org/10.1136/thoraxjnl-2017-211417)]

Charususin N, Gosselink R, Decramer M, Demeyer H, McConnell A, Saey D, et al. Relation between training quality, improvements in inspiratory muscle function, and changes in exercise capacity following an inspiratory muscle training intervention (IMTCO study). *European Respiratory Journal* 2017;**50 Suppl 61**:PA3278.

Charususin N, Gosselink R, Decramer M, McConnell A, Demeyer H, Saey D, et al. A multicentre randomised controlled trial of inspiratory muscle training for patients with chronic obstructive pulmonary disease (IMTCO study). *European Respiratory Journal* 2017;**50 Suppl 61**:OA2925. [DOI: [10.1183/1393003.congress-2017.OA2925](https://doi.org/10.1183/1393003.congress-2017.OA2925)]

Charususin N, Gosselink R, Decramer M, McConnell A, Saey D, Maltais F, et al. Inspiratory muscle training protocol for patients with chronic obstructive pulmonary disease (IMTCO study): a multicentre randomised controlled trial. *BMJ Open* 2013;**3**(8):e003101. [DOI: [10.1136/bmjopen-2013-003101](https://doi.org/10.1136/bmjopen-2013-003101)]

Charususin NA, Gosselink R, Decramer M, McConnell A, Saey D, Maltais F, et al. Inspiratory muscle training for patients with chronic obstructive pulmonary disease (IMTCO study): a multicentre randomised controlled trial. *European Respiratory Journal* 2016;**48 Suppl 60**:OA1519. [DOI: [10.1183/13993003.congress-2016.OA1519](https://doi.org/10.1183/13993003.congress-2016.OA1519)]

Gloeckl R, Schneeberger T, Jarosch I, Charususin N, Langer D, Kenn K, et al. Complementary inspiratory muscle training during pulmonary rehabilitation in COPD patients with inspiratory muscle weakness – a subgroup analysis of a randomized, controlled trial (IMTCO study). *European Respiratory Journal* 2017;**50 Suppl 61**:PA3277. [DOI: [10.1183/1393003.congress-2017.PA3277](https://doi.org/10.1183/1393003.congress-2017.PA3277)]

NCT01397396. Effects of inspiratory muscle training in COPD (IMTCO) [Effects of inspiratory muscle training in patients with chronic obstructive pulmonary disease - a randomized controlled trial]. clinicaltrials.gov/show/NCT01397396 (first received 19 July 2011).

Chuang 2017 {published data only}

Chuang H-Y, Chang H-Y, Fang Y-Y, Guo S-E. The effects of threshold inspiratory muscle training in patients with chronic obstructive pulmonary disease: a randomised experimental study. *Journal of Clinical Nursing* 2017;**26**(23-4):4830-8. [DOI: [10.1111/jocn.13841](https://doi.org/10.1111/jocn.13841)]

Covey 2001 {published data only}

Covey MK, Larson JL, Wirtz SE, Berry JK, Pogue NJ, Alex CG, et al. High-intensity inspiratory muscle training in patients with chronic obstructive pulmonary disease and severely reduced function. *Journal of Cardiopulmonary Rehabilitation* 2001;**21**(4):231-40. [PMID: 11508185]

Cutrim 2019 {published data only}

Cutrim AL, Duarte AA, Silva-Filho AC, Dias CJ, Urtado CB, Ribeiro RM, et al. Inspiratory muscle training improves autonomic modulation and exercise tolerance in chronic obstructive pulmonary disease subjects: a randomized-

controlled trial. *Respiratory Physiology and Neurobiology* 2019;**263**:31-7. [DOI: [10.1016/j.resp.2019.03.003](https://doi.org/10.1016/j.resp.2019.03.003)]

Dacha 2019 {published data only}

Dacha S, Rodrigues A, Louvaris Z, Janssens L, Janssens W, Gosselink R, et al. Effects of inspiratory muscle training (IMT) on dyspnea, respiratory muscle function and respiratory muscle activation in patients with COPD during endurance cycling. *European Respiratory Journal* 2019;**54 Suppl 63**:PA2199. [DOI: [10.1183/13993003.congress-2019.PA2199](https://doi.org/10.1183/13993003.congress-2019.PA2199)]

De Farias 2019 {published and unpublished data}

de Farias CA, Gualdi LP, da Silva SB, Parreira VF, Montemezzo D, Resqueti VR, et al. Effects of different modalities of inspiratory muscle training as an add-on to conventional treatment of patients with chronic obstructive pulmonary disease (COPD): study protocol for a randomized controlled trial. *Trials* 2019;**20**(1):231. [DOI: [10.1186/s13063-019-3271-1](https://doi.org/10.1186/s13063-019-3271-1)]

* Farias C, Fregonezi G, Batista IP, Castro L, Lopes N, Gualdi L, et al. Effects of respiratory muscle training with different modalities in COPD patients: a randomized controlled trial. *European Respiratory Journal* 2019;**54 Suppl 63**:PA2198. [DOI: [10.1183/13993003.congress-2019.PA2198](https://doi.org/10.1183/13993003.congress-2019.PA2198)]

RBR-94v6kd. Effects of exercise of the respiratory muscles with different types of devices in people with chronic obstructive pulmonary disease. <https://ensaiosclinicos.gov.br/rg/RBR-94v6kd/> (first received 23 November 2016).

Dekhuijzen 1991 {published data only}

Dekhuijzen PN, Beek MM, Folgering HT, Van Herwaarden CL. Psychological changes during pulmonary rehabilitation and target-flow inspiratory muscle training in COPD patients with a ventilatory limitation during exercise. *International Journal of Rehabilitation Research* 1990;**13**:109-17.

* Dekhuijzen PN, Folgering HT, Van Herwaarden CL. Target-flow inspiratory muscle training during pulmonary rehabilitation in patients with COPD. *Chest* 1991;**99**(1):128-33. [DOI: [10.1378/chest.99.1.128](https://doi.org/10.1378/chest.99.1.128)]

Dekhuijzen PNR, Van Herwaarden CLA, Folgering H. Target-flow inspiratory muscle training (IMT) increases inspiratory muscle strength and endurance. *European Respiratory Journal* 1989;**2**:389.

Dellweg 2017 {published data only}

* Dellweg D, Reissig K, Hoehn E, Siemon K, Haidl P. Inspiratory muscle training during rehabilitation in successfully weaned hypercapnic patients with COPD. *Respiratory Medicine* 2017;**123**:116-23. [DOI: [10.1016/j.rmed.2016.12.006](https://doi.org/10.1016/j.rmed.2016.12.006)]

NCT00291460. Inspiratory muscle training in hypercapnic COPD [Randomized controlled trial of inspiratory muscle training in patients with stable hypercapnic respiratory failure due to COPD]. clinicaltrials.gov/show/NCT00291460 (first received 14 February 2006).

Fanfa Bordin 2020 {published data only}

* Fanfa Bordin D, Cardoso DM, Wagner LE, Beckenkamp PR, Goncalves da Silva AL, Nunes Paiva D. Sternoleidomastoid

muscle activation following inspiratory muscle training in patients with chronic obstructive pulmonary disease: a randomized clinical trial. *Fisioterapia e Pesquisa* 2020;**27**(2):133-9. [DOI: [10.1590/1809-2950/19009727022020](https://doi.org/10.1590/1809-2950/19009727022020)]

NCT02014155. Effects of inspiratory muscle training on respiratory electromyographic activity in patients with COPD participants and non-participants of a pulmonary rehabilitation program. clinicaltrials.gov/show/nct02014155 (first received 18 December 2013).

Harver 1989 {published data only}

Harver, AA, Mahler D, Daubenspeck JA. Targeted inspiratory muscle training improves respiratory muscle function and reduces dyspnea in patients with chronic obstructive pulmonary disease. *Annals of Internal Medicine* 2020;**111**(2):117-24.

* Harver A, Mahler DA, Daubenspeck JA. Targeted inspiratory muscle training improves respiratory muscle function and reduces dyspnea in patients with chronic obstructive pulmonary disease. *Annals of Internal Medicine* 1989;**111**(2):117-24. [DOI: [10.7326/0003-4819-111-2-117](https://doi.org/10.7326/0003-4819-111-2-117)]

Heijdra 1996 {published data only}

* Heijdra YF, Dekhuijzen PN, Van Herwaarden CL, Folgering HT. Nocturnal saturation improves by target-flow inspiratory muscle training in patients with COPD. *American Journal Respiratory Critical Care Medicine* 1996;**153**(1):260-5.

Heijdra YF, Dekhuijzen PN, Van Herwaarden CL, Folgering HT. Target flow inspiratory muscle training (IMT) improves the nocturnal saturation in patients with COPD. *European Respiratory Journal* 1994;**7 Suppl 18**:237s.

Hill 2006 {published data only}

Hill K, Jenkins SC, Philippe DL, Cecins N, Shepherd KL, Green DJ, et al. High-intensity inspiratory muscle training in COPD. *European Respiratory Journal* 2006;**27**(6):1119-28. [PMID: 16772388]

Hill 2007 {published data only}

Hill K, Jenkins SC, Philippe DL, Shepherd KL, Hillman DR, Eastwood PR. Comparison of incremental and constant load tests of inspiratory muscle endurance in COPD. *European Respiratory Journal* 2007;**30**(3):479-86. [DOI: [10.1183/09031936.00095406](https://doi.org/10.1183/09031936.00095406)]

Hsiao 2003 {published data only}

Hsiao SF, Wu YT, Wu HD, Wang TG, Wang TG. Comparison of effectiveness of pressure threshold and targeted resistance devices for inspiratory muscle training in patients with chronic obstructive pulmonary disease. *Journal of the Formosan Medical Association / Taiwan yi zhi* 2003;**102**(4):240-5.

Kim 1993 {published data only}

Kim MJ, Larson JL, Covey MK, Vitalo CA, Alex CG, Patel M. Inspiratory muscle training in patients with chronic obstructive pulmonary disease. *Nursing Research* 1993;**42**(6):356-62.

Koppers 2006 {published data only}

Koppers RJ, Vos PJ, Boot CR, Folgering HT. Exercise performance improves in patients with COPD due to respiratory muscle endurance training. *Chest* 2006;**129**(4):886-92. [DOI: [10.1378/chest.129.4.886](https://doi.org/10.1378/chest.129.4.886)]

Langer 2018 {published data only}

Dacha S, Langer D, Ciavaglia C, Webb K, Preston M, O'Donnell DE. Effect of inspiratory muscle training (IMT) on static and dynamic respiratory muscle function in patients with COPD. *European Respiratory Journal* 2017;**50 Suppl 61**:OA2923. [DOI: [10.1183/1393003.congress-2017.OA2923](https://doi.org/10.1183/1393003.congress-2017.OA2923)]

Dacha S, Langer D, Ciavaglia C, Webb K, Preston M, O'Donnell DE. Effect of inspiratory muscle training on respiratory muscle function and diaphragm activation in patients with COPD. *American Journal of Respiratory and Critical Care Medicine* 2017;**195**(Meeting Abstracts):A2860. [DOI: [10.1164/ajrccm-conference.2017.A109](https://doi.org/10.1164/ajrccm-conference.2017.A109)]

Langer D, Charususin N, Jacome C, Hoffman M, McConnell A, Decramer M, et al. Efficacy of a novel method for inspiratory muscle training in people with chronic obstructive pulmonary disease. *Physical Therapy* 2015;**95**(9):1264-73. [DOI: [10.2522/ptj.20140245](https://doi.org/10.2522/ptj.20140245)]

* Langer D, Ciavaglia C, Faisal A, Webb KA, Neder JA, Gosselink R, et al. Inspiratory muscle training reduces diaphragm activation and dyspnea during exercise in COPD. *Journal of Applied Physiology* 2018;**125**(2):381-92. [DOI: [10.1152/jappphysiol.01078.2017](https://doi.org/10.1152/jappphysiol.01078.2017)]

Langer D, Ciavaglia C, Webb K, Preston M, Neder JA, Gosselink R, et al. Inspiratory muscle training reduces respiratory neural drive (RND) during exercise in patients with COPD. *European Respiratory Journal* 2014;**44 Suppl 58**:1912.

NCT01900873. Effects of inspiratory muscle training on dyspnea perception during exercise in patients with COPD (IMTCOCOPD). clinicaltrials.gov/show/NCT01900873 (first received 17 July 2013).

Larson 1988 {published data only}

Larson JL, Kim MJ, Sharp JT, Larson DA. Inspiratory muscle training with a pressure threshold breathing device in patients with chronic obstructive pulmonary disease. *American Review of Respiratory Disease* 1988;**138**(3):689-96. [DOI: [10.1164/ajrccm/138.3.689](https://doi.org/10.1164/ajrccm/138.3.689)]

Larson 1999 {published data only}

Larson JL, Covey MK, Wirtz SE, Berry JK, Alex CG, Langbein WE, et al. Cycle ergometer and inspiratory muscle training in chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine* 1999;**160**(2):500-7. [DOI: [10.1164/ajrccm.160.2.9804067](https://doi.org/10.1164/ajrccm.160.2.9804067)]

Leelarungrayub 2017 {published data only}

Leelarungrayub J, Pinkaew D, Puntumetakul R, Klaphajone J. REffects of a simple prototype respiratory muscle trainer on respiratory muscle strength, quality of life and dyspnea, and oxidative stress in COPD patients: a preliminary study.

International Journal of Chronic Obstructive Pulmonary Disease 2017;**12**:1415-25. [DOI: [10.2147/COPD.S131062](https://doi.org/10.2147/COPD.S131062)]

Lisboa 1997 {published data only}

Lisboa C, Villafranca C, Leiva A, Cruz E, Pertuze J, Borzone G. Inspiratory muscle training in chronic airflow limitation: effect on exercise performance. *European Respiratory Journal* 1997;**10**(3):537-42.

Mador 2005 {published data only}

* Mador MJ, Deniz O, Aggarwal A, Shaffer M, Kufel TJ, Spengler CM, et al. Effect of respiratory muscle endurance training in patients with COPD undergoing pulmonary rehabilitation. *Chest* 2005;**128**(3):1216-24. [DOI: [10.1378/chest.128.3.1216](https://doi.org/10.1378/chest.128.3.1216)]

Mador MJ, Deniz O, Aggarwal A, Shaffer M, Kufel TJ, Spengler CM. Effect of respiratory muscle endurance training in patients with COPD undergoing pulmonary rehabilitation. In: American Thoracic Society 99th International Conference; 2003 May 16-21; Seattle. 2003:B046 Poster D8.

Magadle 2007 {published data only}

Magadle R, McConnell AK, Beckerman M, Weiner P. Inspiratory muscle training in pulmonary rehabilitation program in COPD patients. *Respiratory Medicine* 2007;**101**(7):1500-5. [DOI: [10.1016/j.rmed.2007.01.010](https://doi.org/10.1016/j.rmed.2007.01.010)]

Majewska-Pulsakowska 2016 {published data only}

Majewska-Pulsakowska M, Wytrychowski K, Rozek-Piechura K. The role of inspiratory muscle training in the process of rehabilitation of patients with chronic obstructive pulmonary disease. *Advances in Experimental Medicine and Biology* 2016;**885**:47-51. [DOI: [10.1007/5584_2015_194](https://doi.org/10.1007/5584_2015_194)]

Masanga 2011 {published data only}

Masanga L, Fernandez L. Additive effects of high-intensity inspiratory muscle training in pulmonary rehabilitation among COPD patients in the Philippines. *Respirology* 2011;**16** Suppl 2:119 [1130]. [DOI: [10.1111/j.1400-1843.2011.02071.x](https://doi.org/10.1111/j.1400-1843.2011.02071.x)]

Nikoleitou 2016 {published data only}

Nikoleitou D, Backley JA, Gearing J, Man WD, Mustafa N, Johnson LC, et al. Inspiratory muscle strength and endurance training in moderate to severe COPD patients [Abstract]. In: American Thoracic Society 100th International Conference; 2004 May 21-26; Orlando. 2004:D96 Poster 115.

Nikoleitou D, Backley JA, Gearing J, Man WD, Mustafa N, Johnson LC, et al. A double-blind, randomised controlled trial of inspiratory muscle training in COPD patients [Abstract]. *Thorax* 2003;**58** Suppl 3:iii77.

* Nikoleitou D, Man WD-C, Mustafa N, Moore J, Rafferty G, Grant RL, et al. Evaluation of the effectiveness of a home-based inspiratory muscle training programme in patients with chronic obstructive pulmonary disease using multiple inspiratory muscle tests. *Disability and Rehabilitation* 2016;**38**(3):250-9. [DOI: [10.3109/09638288.2015.1036171](https://doi.org/10.3109/09638288.2015.1036171)]

Paneroni 2018 {published data only}

Paneroni M, Simonelli C, Saleri M, Trainini D, Fokom G, Speltoni I, et al. Short-term effects of normocapnic hyperpnea and exercise training in patients with chronic obstructive pulmonary disease: a pilot study. *American Journal of Physical Medicine and Rehabilitation* 2018;**97**(12):866-72. [DOI: [10.1097/PHM.0000000000000988](https://doi.org/10.1097/PHM.0000000000000988)]

Petrovic 2012 {published data only}

NCT00469313. Efficacy of inspiratory muscle training on inspiratory capacity in patients with COPD. clinicaltrials.gov/show/NCT00469313 (first received 4 May 2007).

* Petrovic M, Reiter M, Zipko H, Pohl W, Wanke T. Effects of inspiratory muscle training on dynamic hyperinflation in patients with COPD. *International Journal of Chronic Obstructive Pulmonary Disease* 2012;**7**:797-805. [DOI: [10.2147/COPD.S23784](https://doi.org/10.2147/COPD.S23784)]

Preusser 1994 {published data only}

Preusser BA, Winningham ML, Clanton TL. The effects of high versus low-intensity inspiratory muscle interval training in patients with COPD. *Chest* 1994;**106**(1):110-17.

Preusser BA, Winningham ML, Clanton TL. High- vs low-intensity inspiratory muscle interval training in patients with COPD. *Chest* 1994;**106**(1):110-7. [DOI: [10.1378/chest.106.1.110](https://doi.org/10.1378/chest.106.1.110)]

Ramirez Sarmiento 2002 {published data only}

Ramirez-Sarmiento A, Orozco-Levi M, Guell R, Barreiro E, Hernandez N, Mota S, et al. Inspiratory muscle training in patients with chronic obstructive pulmonary disease: structural adaptation and physiologic outcomes. *American Journal of Respiratory and Critical Care Medicine* 2002;**166**(11):1491-7. [DOI: [10.1164/rccm.200202-075OC](https://doi.org/10.1164/rccm.200202-075OC)]

Saher 2021 {published data only}

Saher T, Moiz JA, Bhati P, Ali MS, Talwar D. Effect of inspiratory muscle training in hypercapnic chronic obstructive pulmonary disease patients during acute care: a randomised clinical trial. *Comparative Exercise Physiology* 2021;**17**(1):55-63.

Saka 2021 {published data only}

NCT03517839. Assessment of impact of inspiratory muscle training on movement fear due to dyspnea in chronic obstructive pulmonary disease. clinicaltrials.gov/show/nct03517839 (first received 7 May 2018).

* Saka S, Gurses HN, Bayram M. Effect of inspiratory muscle training on dyspnea-related kinesiophobia in chronic obstructive pulmonary disease: a randomized controlled trial. *Complementary Therapies in Clinical Practice* 2021;**44**:101418.

Sanchez Riera 2001 {published data only}

Sanchez Riera H, Montemayor Rubio T, Ortega Ruiz F, Cejudo Ramos P, Del Castillo Otero D, Elias Hernandez T, et al. Inspiratory muscle training in patients with COPD: effect on dyspnea, exercise performance, and quality of life. *Chest* 2001;**120**(3):748-56. [DOI: [10.1378/chest.120.3.748](https://doi.org/10.1378/chest.120.3.748)]

Scherer 2000 {published data only}

Scherer TA, Spengler CM, Owassapian D, Imhof E, Boutellier U. Respiratory muscle endurance training in chronic obstructive

pulmonary disease: impact on exercise capacity, dyspnea, and quality of life. *American Journal of Respiratory and Critical Care Medicine* 2000;**162**(5):1709-14. [DOI: [10.1164/ajrccm.162.5.9912026](https://doi.org/10.1164/ajrccm.162.5.9912026)]

Schultz 2018 {published data only}

* Schultz K, Jelusic D, Wittmann M, Kramer B, Huber V, Fuchs S, et al. Inspiratory muscle training does not improve clinical outcomes in 3-week COPD rehabilitation: results from a randomised controlled trial. *European Respiratory Journal* 2018;**51**(1):1702000. [DOI: [10.1183/13993003.02000-2017](https://doi.org/10.1183/13993003.02000-2017)]

Schultz K, Kramer B, Fuchs S, Wingart S, Lehbert N, Huber V, et al. Effects of routine inspiratory muscle training (IMT) as add-on to pulmonary rehabilitation (PR) in COPD. *European Respiratory Journal* 2015;**46** Suppl 59:PA544. [DOI: [10.1183/13993003.congress-2015.PA544](https://doi.org/10.1183/13993003.congress-2015.PA544)]

Sykes 2005 {published data only}

Sykes K, Hang HW. Inspiratory muscle training in the treatment of chronic obstructive pulmonary disease: randomized controlled trial. *American Journal of Recreation Therapy* 2005;**4**(2):39-48.

Tounsi 2021 {published data only}

Tounsi B, Acheche A, Lelard T, Tabka Z, Trabelsi Y, Ahmaidi S. Effects of specific inspiratory muscle training combined with whole-body endurance training program on balance in COPD patients: randomized controlled trial. *PLOS ONE* 23 September 2023;**16**(9):e0257595. [DOI: [10.1371/journal.pone.0257595](https://doi.org/10.1371/journal.pone.0257595)]

Tout 2013 {published data only}

Tout R, Tayara L, Halimi M. The effects of respiratory muscle training on improvement of the internal and external thoraco-pulmonary respiratory mechanism in COPD patients. *Annals of Physical and Rehabilitation Medicine* 2013;**56**(3):193-211. [DOI: [10.1016/j.rehab.2013.01.008](https://doi.org/10.1016/j.rehab.2013.01.008)]

Wang 2017 {published data only}

* Wang K, Zeng G-Q, Li R, Luo Y-W, Wang M, Hu Y-H, et al. Cycle ergometer and inspiratory muscle training offer modest benefit compared with cycle ergometer alone: a comprehensive assessment in stable COPD patients. *International Journal of Chronic Obstructive Pulmonary Disease* 2017;**12**:2655-68. [DOI: [10.2147/COPD.S140093](https://doi.org/10.2147/COPD.S140093)]

Yuwen L, Peng L, Yun L, Yitai C, Yuxia H, Kai W, et al. Effects of combined cycle ergometer and inspiratory muscle training in patients with stable chronic obstructive pulmonary disease. *Respirology* 2016;**21** Suppl 3:176 (APSR6-0158). [DOI: [10.1111/resp.12939-15](https://doi.org/10.1111/resp.12939-15)]

Weiner 1992 {published data only}

* Weiner P, Azgad Y, Ganam R, Weiner M. Inspiratory muscle training combined with general exercise reconditioning in patients with COPD. *Chest* 1992;**102**(5):1351-6. [DOI: [10.1378/chest.102.5.1351](https://doi.org/10.1378/chest.102.5.1351)]

Weiner P, Azgad Y, Weiner M, Ganam R. Inspiratory muscle training combined with general exercise reconditioning in chronic obstructive pulmonary disease. *Harefuah* 1993;**124**(7):396-400, 456.

Weiner 2000 {published data only}

Weiner P, Magadle R, Berar-Yanay N, Davidovich A, Weiner M. The cumulative effect of long-acting bronchodilators, exercise, and inspiratory muscle training on the perception of dyspnea in patients with advanced COPD. *Chest* 2000;**118**(3):672-8. [DOI: [10.1378/chest.118.3.672](https://doi.org/10.1378/chest.118.3.672)]

Weiner 2003 {published data only}

Weiner P, Magadle R, Beckerman M, Weiner M, Berar-Yanay N. Comparison of specific expiratory, inspiratory, and combined muscle training programs in COPD. *Chest* 2003;**124**(4):1357-64. [DOI: [10.1378/chest.124.4.1357](https://doi.org/10.1378/chest.124.4.1357)]

Weiner 2006 {published data only}

Weiner P, Weiner M. Inspiratory muscle training may increase peak inspiratory flow in chronic obstructive pulmonary disease. *Respiration* 2006;**73**(2):151-6. [DOI: [10.1159/000088095](https://doi.org/10.1159/000088095)]

Wu 2017 {published data only}

NCT03101774. Inspiratory muscle training on respiratory muscle function, quality of life and exercise capacity in stable COPD [Effects of respiratory physiology-oriented inspiratory muscle training on respiratory muscle function, quality of life and exercise capacity in stable COPD with inspiratory muscle weakness: a randomised controlled trial]. clinicaltrials.gov/show/nct03101774 (first received 5 April 2017).

* Wu W, Guan L, Zhang X, Li X, Yang Y, Guo B, et al. Effects of two types of equal-intensity inspiratory muscle training in stable patients with chronic obstructive pulmonary disease: a randomised controlled trial. *Respiratory Medicine* 2017;**132**:84-91. [DOI: [10.1016/j.rmed.2017.10.001](https://doi.org/10.1016/j.rmed.2017.10.001)]

Xu 2018 {published data only}

Xu W, Li R, Guan L, Wang K, Hu Y, Xu L, et al. Combination of inspiratory and expiratory muscle training in same respiratory cycle versus different cycles in COPD patients: a randomized trial. *Respiratory Research* 2018;**19**(1):225. [DOI: [10.1186/s12931-018-0917-6](https://doi.org/10.1186/s12931-018-0917-6)]

ZhouL 2016 {published data only}

Zhou L-Q, Li X-Y, Li Y, Guo B-P, Guan L-L, Chen X, et al. Inspiratory muscle training followed by non-invasive positive pressure ventilation in patients with severe chronic obstructive pulmonary disease: a randomized controlled trial. *Nan fang yi ke da xue xue bao [Journal of Southern Medical University]* 2016;**36**(8):1069-74.

References to studies excluded from this review

Ahmad 2013 {published data only}

Ahmad H, Justine M, Othman Z, Mohan V, Mirza FT. The outcomes of short term inspiratory muscle training (IMT) combined with chest physiotherapy in hospitalized COPD patients. *Bangladesh Journal of Medical Science* 2013;**12**(4):398-404. [DOI: [10.3329/bjms.v12i4.13302](https://doi.org/10.3329/bjms.v12i4.13302)]

Aldrich 1985 {published data only}

Aldrich TK. The application of muscle endurance training techniques to the respiratory muscles in COPD. *Lung* 1985;**163**(1):15-22. [DOI: [10.1007/BF02713802](https://doi.org/10.1007/BF02713802)]

Anand 2013 {published data only}

Anand A, Narwal R, Sindhvani G. Accessory inspiratory muscles energy technique effect on pulmonary function in COPD subjects. *Indian Journal of Physiotherapy and Occupational Therapy* 2013;**7**(3):192-7. [DOI: [10.5958/j.0973-5674.7.3.091](https://doi.org/10.5958/j.0973-5674.7.3.091)]

Baines 2005 {published data only}

Baines SJ, Enright SJ. Does a threshold loading device maintain functional improvements following a programme of high intensity inspiratory muscle training in patients with chronic obstructive pulmonary disease? [Abstract]. *European Respiratory Journal* 2005;**26** Suppl 49:4583.

Basso Vanelli 2016 {published data only}

Basso-Vanelli R, Pantoni C, Borghi-Silva A, Labadessa I, Regueiro E, Di Lorenzo V, et al. Inspiratory muscle training improves thoracoabdominal asynchronism during unsupported upper limb exercise in COPD patients-pilot study. *European Respiratory Journal* 2016;**48** Suppl 60:PA1373. [DOI: [10.1183/13993003.congress-2016.PA1373](https://doi.org/10.1183/13993003.congress-2016.PA1373)]

* Basso-Vanelli RP, Pires Di Lorenzo VA, Labadessa IG, Regueiro Eloisa MG, Jamami Mauricio. Effects of inspiratory muscle training and calisthenics-and-breathing exercises in COPD with and without respiratory muscle weakness. *Respiratory Care* 2016;**61**(1):50-60. [DOI: [10.4187/respcare.03947](https://doi.org/10.4187/respcare.03947)]

NCT01510041. Effects of respiratory muscle training and respiratory exercise in exercise tolerance, performing daily life activities and quality of life of patients with chronic obstructive pulmonary disease. clinicaltrials.gov/show/NCT01510041 (first received 13 January 2012).

Battaglia 2009 {published data only}

Battaglia E, Fulgenzi A, Ferrero ME. Rationale of the combined use of inspiratory and expiratory devices in improving maximal inspiratory pressure and maximal expiratory pressure of patients with chronic obstructive pulmonary disease. *Archives of Physical Medicine and Rehabilitation* 2009;**90**(6):913-18. [DOI: [10.1016/j.apmr.2008.12.019](https://doi.org/10.1016/j.apmr.2008.12.019)]

Belman 1994 {published data only}

Belman MJ, Botnick WC, Nathan SD, Chon KH. Ventilatory load characteristics during ventilatory muscle training. *Journal of Cardiopulmonary Rehabilitation and Prevention* 1994;**14**(6):409.

Bgin 1991 {published data only}

Bégin P, Grassino A. Inspiratory muscle dysfunction and chronic hypercapnia in chronic obstructive pulmonary disease. *American Review of Respiratory Disease* 1991;**143**(5 Pt 1):905-12. [DOI: [10.1164/ajrccm/143.5_Pt_1.905](https://doi.org/10.1164/ajrccm/143.5_Pt_1.905)]

Bissett 2016 {published data only}

Bissett B, Leditschke IA, Neeman T, Boots R, Paratz J. Inspiratory muscle training to enhance recovery from prolonged mechanical ventilation: a randomized trial. *American Journal of Respiratory and Critical Care Medicine* 2016;**193**(Meeting Abstracts):A2613.

Bjerre Jepsen 1981 {published data only}

Bjerre-Jepsen K, Secher NH, Kok-Jensen A. Inspiratory resistance training in severe chronic obstructive pulmonary disease. *European Journal of Respiratory Diseases* 1981;**62**(6):405-11.

Cader 2010 {published data only}

Cader SA, Vale RG, Castro JC, Bacelar SC, Biehl C, Gomes MC, et al. Inspiratory muscle training improves maximal inspiratory pressure and may assist weaning in older intubated patients: a randomised trial. *Journal of Physiotherapy* 2010;**56**(3):171-7. [DOI: [10.1016/s1836-9553\(10\)70022-9](https://doi.org/10.1016/s1836-9553(10)70022-9)]

Chen 1985 {published data only}

Chen H, Dukes R, Martin BJ. Inspiratory muscle training in patients with chronic obstructive pulmonary disease. *American Review of Respiratory Disease* 1985;**131**(2):251-5. [DOI: [10.1164/arrd.1985.131.2.251](https://doi.org/10.1164/arrd.1985.131.2.251)]

Daynes 2018 {published data only}

Daynes E, Greening NJ, Harvey-Dunstan TC, Singh SJ. High-frequency airway oscillating device for respiratory muscle training in subjects with COPD. *Respiratory Care* 2018;**63**(5):584-90. [DOI: [10.4187/respcare.05837](https://doi.org/10.4187/respcare.05837)]

de Andrade 2005 {published data only}

de Andrade AD, Silva TN, Vasconcelos H, Marcelino M, Rodrigues-Machado MG, Filho VC, et al. Inspiratory muscular activation during threshold therapy in elderly healthy and patients with COPD. *Journal of Electromyography and Kinesiology* 2005;**15**(6):631-9. [DOI: [10.1016/j.jelekin.2005.06.002](https://doi.org/10.1016/j.jelekin.2005.06.002)]

de Lucas Ramos 1998 {published data only}

de Lucas Ramos P, Rodriguez Gonzalez-Moro JM, Garcia de Pedro J, Santacruz Siminiani A, Tatay Marti E, Cubillo Marcos JM. Training of inspiratory muscles in chronic obstructive lung disease. Its impact on functional changes and exercise tolerance. *Archivos de Bronconeumologia* 1998;**34**(2):64-70.

Di Mambro 2007 {published data only}

Di Mambro TR, Figueiredo PH, Wanderley TR, Kristki AL, Guimaraes FS. Inspiratory muscle training in chronic obstructive pulmonary disease: impact on quality of life, exercise intolerance, and dyspnea. *Fisioterapia e Pesquisa* 2007;**14**(2):65-71.

DRKS00005637 {published data only}

DRKS00005637. Respiratory muscle activation by respiratory muscle training in patients with advanced COPD [Respiratory muscle activation by respiratory muscle training in patients with advanced COPD - EMG-Train COPD]. ictrptest.azurewebsites.net/Trial2.aspx?TrialID=DRKS00005637 (first received 4 February 2014).

DRKS00006021 {published data only}

DRKS00006021. Influence of a controlled inspiratory muscle training on the course of COPD after acute exacerbation (AECOPD): a controlled study. www.drks.de/DRKS00006021 (first received 9 May 2014).

Elbouhy 2014 {published data only}

Elbouhy MS, AbdelHalim HA, Hashem AM. Effect of respiratory muscles training in weaning of mechanically ventilated COPD patients. *Egyptian Journal of Chest Diseases and Tuberculosis* 2014;**63**(3):679-87. [DOI: <http://dx.doi.org/10.1016/j.ejcdt.2014.03.008>]

Elmorsi 2016 {published data only}

Elmorsi AS, Eldesoky ME, Mona AA, Shalaby NM, Abdalla Dina A, et al. Effect of inspiratory muscle training on exercise performance and quality of life in patients with chronic obstructive pulmonary disease. *Egyptian Journal of Chest Diseases and Tuberculosis* 2016;**65**(1):41-6. [DOI: [10.1016/j.ejcdt.2015.10.006](https://doi.org/10.1016/j.ejcdt.2015.10.006)]

Enright 2005 {published data only}

Enright SJ, Baines SJ. Evaluation of a new computerised method of high intensity inspiratory muscle training in patients with chronic airways obstruction [Abstract]. *European Respiratory Journal* 2005;**26 Suppl 49**:Abstract No. 4584.

Garcia 2008 {published data only}

* Garcia S, Rocha M, Pinto P, Lopes AM, Bárbara C. Treino de músculos inspiratórios em doentes com DPOC. *Revista Portuguesa de Pneumologia* 2008;**14**(2):177-94.

Garcia S, Rocha M, Pinto P, Lopes AM, Barbara C. Inspiratory muscle training in COPD patients. *Revista Portuguesa de Pneumologia* 2008;**14**(2):177-94.

Goldstein 1989 {published data only}

Goldstein R, De Rosie J, Long S, Dolmage T, Avendano MA. Applicability of a threshold loading device for inspiratory muscle testing and training in patients with COPD. *Chest* 1989;**96**(3):564-71.

Gregg 1989 {published data only}

Gregg BL. Inspiratory muscle training with a weighted incentive spirometer in subjects with chronic airways obstruction. *Respiratory Care* 1989;**34**(10):860-7.

Guyatt 1992 {published data only}

Guyatt G, Keller J, Singer J, Halcrow S, Newhouse M. Controlled trial of respiratory muscle training in chronic airflow limitation. *Thorax* 1992;**47**(8):598-602. [DOI: [10.1136/thx.47.8.598](https://doi.org/10.1136/thx.47.8.598)]

Hart 2000 {published data only}

Hart N, Cramer D, Sylvester K, Nickol AH, Ward SP, Moxham J, et al. Effect of using powerbreathe™ on inspiratory muscle strength. *Thorax* 2000;**55 Suppl 3**:A51.

Heydari 2015 {published data only}

Heydari A, Farzad M, Ahmadi Hosseini SH. Comparing inspiratory resistive muscle training with incentive spirometry on rehabilitation of COPD patients. *Rehabilitation Nursing* 2015;**40**(4):243-8. [DOI: [10.1002/rnj.136](https://doi.org/10.1002/rnj.136)]

Hopp 1996 {published data only}

* Hopp LJ, Kim MJ, Larson JL, Sharp JT. Incremental threshold loading in patients with chronic obstructive pulmonary disease. *Nursing Research* 1996;**45**(4):196-202.

Hopp LJ. Incremental threshold loading in chronic obstructive pulmonary disease patients. *Nursing Research* 1992;**45**:147p.

Ibakordor 2013 {published data only}

Ibakordor S, Dhillon DK, Saravanan S. A safe inspiratory threshold load in chronic obstructive pulmonary disease patients. *Indian Journal of Physiotherapy and Occupational Therapy* 2013;**7**(2):48-52. [DOI: [10.5958/j.0973-5674.7.2.011](https://doi.org/10.5958/j.0973-5674.7.2.011)]

Ionescu 2005 {published data only}

Ionescu AA, Mickleborough TD, Bolton CE, Lindley MR, Chatham K, Nixon LS, et al. Limb and inspiratory muscle dynamic strength in patients with moderate chronic obstructive pulmonary disease [Abstract]. *Thorax* 2005;**2 Suppl II**:ii50.

Izumizaki 2008 {published data only}

Izumizaki M, Satake M, Takahashi H, Sugawara K, Shioya T, Homma I. Effects of inspiratory muscle thixotropy on the 6-min walk distance in COPD. *Respiratory Medicine* 2008;**102**(7):970-7. [DOI: [10.1016/j.rmed.2008.02.007](https://doi.org/10.1016/j.rmed.2008.02.007)]

Johnson 1996 {published data only}

Johnson PH, Cowley AJ, Kinnear WJ. Evaluation of the Threshold trainer for inspiratory muscle endurance training: comparison with the weighted plunger method. *European Respiratory Journal* 1996;**9**(12):2681-4. [DOI: [10.1183/09031936.96.09122681](https://doi.org/10.1183/09031936.96.09122681)]

Kivastik 2015 {published data only}

Kivastik J, Arend M, Maestu J. Comparison of different inspiratory muscle warm-up protocols. *European Respiratory Journal* 2015;**46 Suppl 59**:PA950.

Koch 2020 {published data only}

Koch R, Augusto TR, Ramos AG, Muller PT. Inspiratory muscle training potentiates the beneficial effects of proportional assisted ventilation on exertional dyspnea and exercise tolerance in COPD: a proof-of-concept randomized and controlled trial. *COPD* 2020;**17**(4):1-8. [DOI: [10.1080/15412555.2020.1789085](https://doi.org/10.1080/15412555.2020.1789085)]

Kolesnikova 2016 {published data only}

Kolesnikova E, Arutyunov G, Kostyukevich O, Rylova A. Respiratory muscles trainings are effective in lowering the pneumonia frequency in patients with heart failure and chronic obstructive pulmonary disease. *European Respiratory Journal* 2016;**48 Suppl 60**:PA3773. [DOI: [10.1183/13993003.congress-2016.PA3773](https://doi.org/10.1183/13993003.congress-2016.PA3773)]

Levine 1986 {published data only}

Levine S, Weiser P, Gillen J. Evaluation of a ventilatory muscle endurance training program in the rehabilitation of patients with chronic obstructive pulmonary disease. *American Review of Respiratory Disease* 1986;**133**(3):400-6. [DOI: [10.1164/arrd.1986.133.3.400](https://doi.org/10.1164/arrd.1986.133.3.400)]

Liao 2015 {published data only}

Liao LY, Chen KM, Chung WS, Chien JY. Efficacy of a respiratory rehabilitation exercise training package in hospitalized elderly patients with acute exacerbation of COPD: a randomized control

trial. *International Journal of Chronic Obstructive Pulmonary Disease* 2015;**10**(1):1703-9. [DOI: [10.2147/COPD.S90673](https://doi.org/10.2147/COPD.S90673)]

Lin 2012 {published data only}

Lin W-C, Yuan S-C, Chien J-Y, Weng S-C, Chou M-C, Kuo H-W. The effects of respiratory training for chronic obstructive pulmonary disease patients: a randomised clinical trial. *Journal of Clinical Nursing* 2012;**21**:2870-8. [DOI: [10.1111/j.1365-2702.2012.04124.x](https://doi.org/10.1111/j.1365-2702.2012.04124.x)]

Lisboa 1994 {published data only}

Lisboa C, Munoz V, Beroiza T, Leiva A, Cruz E. Inspiratory muscle training in chronic airflow limitation: comparison of two different training loads with a threshold device. *European Respiratory Journal* 1994;**7**(7):1266-74.

Lisboa 1995a {published data only}

Lisboa C, Villafranca C, Pertuze J, Leiva A, Repetto P. Clinical effects of inspiratory muscle training in patients with chronic airflow limitation. *Revista Medica de Chile* 1995;**123**(9):1108-15.

Lisboa 1995b {published data only}

Lisboa C, Villafranca C, Leiva A, Repetto L, Cruz E, Pertuze R. Inspiratory muscle training in patients with chronic bronchial obstruction: factors related to the improvement of dyspnoea. *Enfermedades respir. Cir. Torac* 1995;**11**(1):7-15.

Lisboa 1998 {published data only}

Lisboa C, Borzone G, Cruz E. Inspiratory muscle training in patients with chronic obstructive pulmonary disease. *Revista Medica de Chile* 1998;**126**(5):563-8.

Madariaga 2007 {published data only}

Madariaga VB, Iturri JB, Manterola AG, Buey JC, Sebastian NT, Pena VS. Comparison of 2 methods for inspiratory muscle training in patients with chronic obstructive pulmonary disease. *Archivos de Bronconeumologia* 2007;**43**(8):431-8. [DOI: [10.1157/13108782](https://doi.org/10.1157/13108782)]

Madsen 1985 {published data only}

Madsen F, Secher NH, Kay L, Rube N. Inspiratory resistance versus general physical training in patients with chronic obstructive pulmonary disease. *European Journal of Respiratory Diseases* 1985;**67**(3):167-76.

Martin 2006 {published data only}

Martin D, Davenport W, Gonzalez-Rothi J, Baz M, Banner J, Caruso L, et al. Inspiratory muscle strength training improves outcome in failure to wean patients [Abstract]. *European Respiratory Journal* 2006;**28** Suppl 50:369s [E2204].

McKeon 1986 {published data only}

McKeon JL, Turner J, Kelly C, Dent A, Zimmerman PV. The effect of inspiratory resistive training on exercise capacity in optimally treated patients with severe chronic airflow limitation. *Australian and New Zealand Journal of Medicine* 1986;**16**(5):648-52. [DOI: [10.1111/j.1445-5994.1986.tb00005.x](https://doi.org/10.1111/j.1445-5994.1986.tb00005.x)]

Meshcheriakova 2006 {published data only}

Meshcheriakova NN, Belevskiy AS, Cherniak AV, Nekludova GV, Appaeva AA. Threshold PEP and IMT devices (PID) for COPD

patient respiratory training [Abstract]. *European Respiratory Journal* 2006;**28** Suppl 50:553s [P3187].

Minoguchi 2002 {published data only}

Minoguchi H, Hibuya M, Miyagawa T, Kokubu F, Yamada M, Tanaka H, et al. Cross-over comparison between respiratory muscle stretch gymnastics and inspiratory muscle training. *Internal Medicine* 2002;**41**(10):805-12. [DOI: [10.2169/internalmedicine.41.805](https://doi.org/10.2169/internalmedicine.41.805)]

NCT01218295 {published data only}

NCT01218295. Inspiratory muscle training with normocapnic hyperpnea in chronic obstructive pulmonary disease (COPD) [Efficacy of inspiratory muscle training by means of Spirotiger® in COPD patients]. clinicaltrials.gov/show/nct01218295 (first received 11 October 2010).

NCT01556139 {published data only}

NCT01556139. Effectiveness of respiratory muscle training by Spirotiger in chronic patients [Evaluation of the effectiveness of respiratory muscle training by the technique of hypocapnia hyperpnea (Spirotiger) in COPD and CHF patients]. clinicaltrials.gov/show/NCT01556139 (first received 16 March 2012).

NCT01747694 {published data only}

NCT01747694. Respiratory muscle exercise training in COPD patients. clinicaltrials.gov/show/nct01747694 (first received 12 December 2012).

NCT01945398 {published data only}

NCT01945398. IMT in ventilatory muscle metaboreflex in COPD [Effect of inspiratory muscle training in the ventilatory muscle metaboreflex in chronic obstructive pulmonary disease patients]. clinicaltrials.gov/show/nct01945398 (first received 18 September 2013).

NCT01956565 {published data only}

NCT01956565. Feasibility of inspiratory muscle training in people with chronic obstructive pulmonary disease (COPD) who decline pulmonary rehabilitation. clinicaltrials.gov/show/NCT01956565 (first received 8 October 2013).

NCT02186340 {published data only}

NCT02186340. Effects of inspiratory muscle training on breathing pattern in patients with chronic obstructive pulmonary disease. clinicaltrials.gov/show/NCT02186340 (first received 10 July 2014).

NCT02278523 {published data only}

NCT02278523. The respiratory physiology variation of COPD patients in inspiratory muscle training. clinicaltrials.gov/show/nct02278523 (first received 30 October 2014).

NCT02579200 {published data only}

NCT02579200. Inspiratory muscle training for dyspneic patients with COPD-HF overlap: a multicenter, randomized controlled trial. clinicaltrials.gov/show/nct02579200 (first received 19 October 2015).

NCT02914093 {published data only}

NCT02914093. IMT in hypercapnic patients with COPD (THYPISK-f) [Feasibility of inspiratory muscle training in patients with chronic hypercapnia and severe chronic obstructive pulmonary disease]. [ClinicalTrials.gov/show/NCT02914093](https://clinicaltrials.gov/show/NCT02914093) (first received 26 September 2016).

NCT02935166 {published data only}

NCT02935166. Innovations in respiratory muscles training in patients with chronic obstructive pulmonary disease (INNOTORIO) [INNOTORIO: Innovations in the training of the inspiratory muscles on patients with chronic obstructive pulmonary disease: design of a digital dual valve and evaluation of a new high-intensity training scheme of short duration]. clinicaltrials.gov/show/nct02935166 (first received 17 October 2016).

NCT03186092 {published data only}

NCT03186092. Effects of respiratory muscle training on respiratory muscle strength, functional capacity and quality of life in pulmonary hypertension. clinicaltrials.gov/show/nct03186092 (first received 14 June 2017).

NCT03500042 {published data only}

NCT03500042. Effects of different modes of respiratory muscle training on respiratory mechanics and NRD in patient with stable COPD [Effects of different modes of respiratory muscular threshold loads training on respiratory mechanics and neural respiratory drive (NRD) in patient with stable chronic obstructive pulmonary disease (COPD)]. clinicaltrials.gov/show/NCT03500042 (first received 17 April 2018).

NCT03739879 {published data only}

NCT03739879. Effectivity of inspiratory muscle trainer in the chronic obstructive pulmonary disease rehabilitation program. clinicaltrials.gov/show/nct03739879 (first received 14 November 2018).

NCT03844711 {published data only}

NCT03844711. Transcutaneous electrical diaphragmatic stimulation and inspiratory muscle training in patients with COPD exacerbated. clinicaltrials.gov/show/nct03844711 (first received 18 February 2019).

NCT03880630 {published data only}

Inspiratory muscle activation pattern analysis in assisting precision in inspiratory muscle training in patients with chronic respiratory disease. clinicaltrials.gov/ct2/show/NCT03880630 (first received 19 March 2019).

NCT04084405 {published data only}

NCT04084405. Effect of inspiratory muscle training on balance in chronic obstructive pulmonary disease patients. clinicaltrials.gov/show/nct04084405 (first received 10 September 2019).

NCT04117399 {published data only}

NCT04117399. Effect of inspiratory muscle training on posture in chronic obstructive pulmonary disease patients. clinicaltrials.gov/show/nct04117399 (first received 7 October 2019).

NCT04460261 {published data only}

NCT04460261. The effects of 'functional' inspiratory muscle training [The effects of a new integrated exercise program called 'functional' inspiratory muscle training in geriatric individuals with and without chronic obstructive pulmonary disease]. [ClinicalTrials.gov/show/NCT04460261](https://clinicaltrials.gov/show/NCT04460261) (first received 7 July 2020).

Neves 2014a {published data only}

Neves LF, Reis MH, Plentz RD, Matte DL, Coronel CC, Sbruzzi G. Expiratory and expiratory plus inspiratory muscle training improves respiratory muscle strength in subjects with COPD: systematic review. *Respiratory Care* 2014;**59**(9):1381-8. [DOI: [10.4187/respcare.02793](https://doi.org/10.4187/respcare.02793)]

Neves 2014b {published data only}

Neves L, Chiappa A, Da Silva V, Vieira P, Cipriano G Jr, Arena R, et al. Comparative effects of inspiratory muscle training and resistance training on respiratory and skeletal muscle strength in COPD: responses of the a pulmonary rehabilitation program. *European Respiratory Journal* 2014;**44**:P598.

Nield 2007 {published data only}

Nield MA, Soo Hoo GW, Roper JM, Santiago S. Efficacy of pursed-lips breathing: a breathing pattern retraining strategy for dyspnea reduction. *Journal of Cardiopulmonary Rehabilitation and Prevention* 2007;**27**(4):237-44. [DOI: [10.1097/01.HCR.0000281770.82652.cb](https://doi.org/10.1097/01.HCR.0000281770.82652.cb)]

Noseda 1987 {published data only}

Noseda A, Carpiaux JP, Vandeput W, Prigogine T, Schmerber J. Resistive inspiratory muscle training and exercise performance in COPD patients. A comparative study with conventional breathing retraining. *Bulletin Europeen de Physiopathologie Respiratoire* 1987;**23**(5):457-63.

O'Connor 2019 {published data only}

O'Connor C, Lawson R, Waterhouse J, Mills GH. Is inspiratory muscle training (IMT) an acceptable treatment option for people with chronic obstructive pulmonary disease (COPD) who have declined pulmonary rehabilitation (PR) and can IMT enhance PR uptake? A single-group prepost feasibility study in a home-based setting. *BMJ Open* 2019;**9**(8):e028507. [DOI: [10.1136/bmjopen-2018-028507](https://doi.org/10.1136/bmjopen-2018-028507)]

Okura 2019 {published data only}

Okura K, Kawagoshi A, Iwakura M, Shibata K, Furukawa Y, Sugawara K, et al. The effectiveness of inspiratory muscle training in patients with COPD. *Journal of the Japan Society for Respiratory Care and Rehabilitation* 2019;**28**(2):274-8. [DOI: [10.15032/jsrcr.28.2_274](https://doi.org/10.15032/jsrcr.28.2_274)]

Okura 2020 {published data only}

Okura K, Takahashi H, Shiotani T, Iida Y, Inagaki T, Ogawa T, et al. Inspiratory muscle training for patients with chronic obstructive pulmonary disease Effect on physical activity — Multicenter, randomized controlled trial — [一多施設による無作為化比較対照試験]. ##### 2020;**47**(6):551-9.

PACTR201703002095224 {published data only}

PACTR201703002095224. Respiratory muscle training in chronic obstructive pulmonary disease [Comparative study between two different respiratory training protocols in patients with chronic obstructive pulmonary disease]. www.who.int/trialsearch/Trial2.aspx?TrialID=PACTR201703002095224 (first received 12 March 2017).

Padula 2001 {published data only}

Padula CA, Yeaw E. Inspiratory muscle training: an exploration of a home-based intervention. *Journal of Applied Research* 2001;**1**(2):85-94.

Perez 2010 {published data only}

Perez ME, Guirao L, Martinez A, Ortega P, Pleguezuelos E, Samitier B. Effect of a specific respiratory exercise program on respiratory muscle strength. In: 71st Annual Assembly of the American Academy of Physical Medicine and Rehabilitation; 2010 Nov 4-7; Seattle. Vol. 2. 2010:S25.

Pescaru 2016 {published data only}

Pescaru C, Oancea C, Tudorache E. New device used in COPD rehabilitation. TrainAir system. *European Respiratory Journal* 2016;**48**:PA695. [DOI: [10.1183/13993003.congress-2016.PA695](https://doi.org/10.1183/13993003.congress-2016.PA695)]

Quintero 1999 {published data only}

Quintero JI, Borzone G, Leiva A, Villafranca C, Lisboa C. Effects of inspiratory muscle training on the oxygen cost of breathing in patients with chronic obstructive pulmonary disease. *Revista Medica De Chile* 1999;**127**(4):421-8.

Richardson 1989 {published data only}

Richardson J, Dunn L, Pardy R. Inspiratory resistive endurance training in patients with chronic obstructive pulmonary disease: a pilot study. *Physiotherapie Canada* 1989;**41**(2):85-92.

Rocha 2015 {published data only}

Rocha T, Souza H, Brandao DC, Rattes C, Ribeiro L, Campos SL, et al. The manual diaphragm release technique improves diaphragmatic mobility, inspiratory capacity and exercise capacity in people with chronic obstructive pulmonary disease: a randomised trial. *Journal of Physiotherapy* 2015;**61**(4):182-9. [DOI: [10.1016/j.jphys.2015.08.009](https://doi.org/10.1016/j.jphys.2015.08.009)]

Sassoon 1992 {published data only}

Sassoon CS, Lodia R, Rheeman CH, Kuei JH, Light RW, Mahutte CK. Inspiratory muscle work of breathing during flow-by, demand-flow, and continuous-flow systems in patients with chronic obstructive pulmonary disease. *American Review of Respiratory Disease* 1992;**145**(5):1219-22. [DOI: [10.1164/ajrccm/145.5.1219](https://doi.org/10.1164/ajrccm/145.5.1219)]

Serón 2005 {published data only}

Serón P, Riedemann P, Muñoz S, Doussoulin A, Villarroel P, Cea X. Effect of inspiratory muscle training on muscle strength and quality of life in patients with chronic airflow limitation: a randomized controlled trial [Efecto del entrenamiento muscular inspiratorio sobre la fuerza muscular y la calidad de vida en pacientes con limitación crónica del flujo aéreo. Ensayo clínico aleatorizado]. *Archivos de Bronconeumología* 2005;**41**(11):601-6. [DOI: [10.1157/13081248](https://doi.org/10.1157/13081248)]

Shahin 2008 {published data only}

Shahin B, Germain M, Kazem A, Annat G. Benefits of short inspiratory muscle training on exercise capacity, dyspnea, and inspiratory fraction in COPD patients. *International Journal of Chronic Obstructive Pulmonary Disease* 2008;**3**(3):423-7. [DOI: [10.2147/copd.s1822](https://doi.org/10.2147/copd.s1822)]

Shioya 2007 {published data only}

Shioya T, Satake M, Takahashi H, Sugawara K, Kasai C, Kiyokawa N, et al. Combination of chest wall mobilization and respiratory muscle training in comprehensive outpatient pulmonary rehabilitation improves pulmonary function in patients with COPD [Abstract]. In: *Respirology*. Vol. 14 Suppl 3. 2009:A191 [PD 05-05].

* Shioya T, Satake M, Takahashi H, Sugawara K, Kasai C, Kiyokawa N, et al. Combination of chest wall mobilization and respiratory muscle training in comprehensive outpatient pulmonary rehabilitation improves pulmonary function in patients with COPD [Abstract]. In: *European Respiratory Journal*. Vol. 30 Suppl 51. 2007:188s [P1160].

Similowski 1994 {published data only}

Similowski T, Derenne JP. Inspiratory muscle testing in stable COPD patients. *European Respiratory Journal* 1994;**7**(10):1871-6.

Sivashanmugam 2019 {published data only}

Sivashanmugam M, Vinod Kumar V, Sridhar R, Kumar R, Stanley G. Comparison of short term inspiratory muscle training and peripheral muscle training on lung function and quality of life in chronic obstructive pulmonary disease patients - a prospective observational study. *Lung India* 2019;**9** Suppl:S107-8. [DOI: [10.4103/0970-2113.271103](https://doi.org/10.4103/0970-2113.271103)]

Soicher 1998 {published data only}

Soicher J, Dechman G. Inspiratory muscle function in chronic obstructive pulmonary disease (COPD). *Physical Therapy Reviews* 1998;**3**(1):31-9. [DOI: [10.1179/ptr.1998.3.1.31](https://doi.org/10.1179/ptr.1998.3.1.31)]

Sonne 1982 {published data only}

Sonne LJ, Davis JA. Increased exercise performance in patients with severe COPD following inspiratory resistive training. *Chest* 1982;**81**(4):436-9. [DOI: [10.1378/chest.81.4.436](https://doi.org/10.1378/chest.81.4.436)]

Sudo 1997 {published data only}

Sudo E, Ohga E, Matsuse T, Teramoto S, Nagase T, Katayama H, et al. The effects of pulmonary rehabilitation combined with inspiratory muscle training on pulmonary function and inspiratory muscle strength in elderly patients with chronic obstructive pulmonary disease. *Nihon Ronen Igakkai zasshi [Japanese Journal of Geriatrics]* 1997;**34**(11):929-34. [DOI: [10.3143/geriatrics.34.929](https://doi.org/10.3143/geriatrics.34.929)]

Sugiyama 2010 {published data only}

Sugiyama N, Yokoyama T, Abe T, Kono Y, Sugiyama S, Setoguchi Y. Combined effects of short-term lower and upper extremity training plus inspiratory muscle training in patients with chronic obstructive pulmonary disease. *Journal of Tokyo Medical University* 2010;**68**(3):322-9.

Sun 2003 {published data only}

Sun JX, Yin MX, Shao H, Li ZS, Li SW. Effect of respiratory muscle gymnastics on lung function and quality of life in the old patients with chronic obstructive pulmonary disease. *Zhonghua Linchuang Kangfu Zazhi* 2003;**7**(27):3698-9.

TCTR20191009004 {published data only}

TCTR20191009004. Physiological change in lung function and respiratory muscle strength in chronic obstructive pulmonary disease patients; cause by pursed-lip breathing exercise using windmill toy model. ictrptest.azurewebsites.net/Trial2.aspx?TrialID=TCTR20191009004 (first received 9 October 2019).

UMIN000030937 {published data only}

UMIN000030937. Efficacy and applicability of respiratory muscle training on the balance of inspiratory-to-expiratory muscle strength and breathing timing in patients with chronic obstructive pulmonary disease: a preparatory research study. ictrptest.azurewebsites.net/Trial2.aspx?TrialID=JPRN-UMIN000030937 1 February 2018.

Van't Hul 2006 {published data only}

Van 't Hul A, Gosselink R, Hollander P, Postmus P, Kwakkel G. Training with inspiratory pressure support in patients with severe COPD. *European Respiratory Journal* 2006;**27**(1):65-72. [DOI: [10.1183/09031936.06.00036505](https://doi.org/10.1183/09031936.06.00036505)]

Villafranca 1998 {published data only}

Villafranca C, Borzone G, Leiva A, Lisboa C. Effect of inspiratory muscle training with an intermediate load on inspiratory power output in COPD. *European Respiratory Journal* 1998;**11**(1):28-33. [DOI: [10.1183/09031936.98.11010028](https://doi.org/10.1183/09031936.98.11010028)]

Wada 2016 {published data only}

Wada JT, Borges-Santos E, Porras DC, Paisani DM, Cukier A, Lunardi AC, et al. Effects of aerobic training combined with respiratory muscle stretching on the functional exercise capacity and thoracoabdominal kinematics in patients with COPD: a randomized and controlled trial. *International Journal of Chronic Obstructive Pulmonary Disease* 2016;**11**:2691-700. [DOI: [10.2147/COPD.S114548](https://doi.org/10.2147/COPD.S114548)]

Wu 2006 {published data only}

Wu X, Hou L, Bai W. Effects of breathing training on quality of life and activities of daily living in elderly patients with stable severe chronic obstructive pulmonary disease. *Chinese Journal of Rehabilitation Medicine* 2006;**21**(3-4):307-10.

Xi 2015 {published data only}

Xi F, Wang Z, Qi Y, Brightwell R, Roberts P, Stewart A, et al. Long-term effect of respiratory training for chronic obstructive pulmonary disease patients at an outpatient clinic: a randomised controlled trial. *Clinical and Translational Medicine* 2015;**4**(1):31. [DOI: <http://dx.doi.org/10.1186/s40169-015-0073-2>]

Yamaguti 2012 {published data only}

Yamaguti WP, Claudino RC, Neto AP, Chammass MC, Gomes AC, Salge JM, et al. Diaphragmatic breathing training program improves abdominal motion during natural breathing in patients with chronic obstructive pulmonary disease: a

randomized controlled trial. *Archives of Physical Medicine and Rehabilitation* 2012;**93**(4):571-7. [DOI: [10.1016/j.apmr.2011.11.026](https://doi.org/10.1016/j.apmr.2011.11.026)]

Yan 1996 {published data only}

Yan Q, Sun Y. Quantitative research for improving respiratory muscle contraction by breathing exercise. *Chinese Medical Journal* 1996;**109**(10):771-5.

Yang 2005 {published data only}

Yang S-Y, Feng E-Z, Shen J-L, Zhang Y, Zhao L-H, Wu X-M, et al. Effect of suitable electrostimulation on phrenic nerve in ameliorating diaphragmatic fatigue and pulmonary function in patients with stable chronic cor pulmonale at high altitude area. *Chinese Journal of Clinical Rehabilitation* 2005;**9**(23):28-30.

Zhang 2008 {published data only}

Zhang Z-Q, Chen R-C, Yang Q-K, Li P, Wang C-Z, Zhang Z-H, et al. Effects of respiratory training in relation to respiratory pathophysiology on respiratory muscle function and exercise tolerance in chronic obstructive pulmonary disease patients. [Chinese]. *Journal of Clinical Rehabilitative Tissue Engineering Research* 2008;**12**(20):3966-71.

References to studies awaiting assessment
Barter 1987 {published data only}

Barter C, Nosworthy J, Flynn M, Pretto JE. Effect of leg and inspiratory muscle training on exercise performance and minute ventilation in chronic airflow obstruction. [Abstract]. *Australian and New Zealand Journal of Medicine* 1987;**17**:511.

Bustamante 1997 {published data only}

Bustamante V, Galdiz JB, Ruiz L, Cabriada V, Serrano L, Sobradillo V. Benefits evolution of inspiratory muscle training after 6 months discontinuation [Abstract]. *European Respiratory Journal* 1997;**10** Suppl 25:7S.

Cassidy 2009 {published data only}

Cassidy C. A study investigating the effects of inspiratory muscle training (IMT) in subjects with chronic obstructive pulmonary disease (COPD) following an acute exacerbation [Abstract]. *Irish Journal of Medical Science* 2009;**178** Suppl 11:S435-6.

Cejudo 1998 {published data only}

Cejudo P, Sanchez H, Ortega F, Toral J, Villagomez R, Montemayor T. Home ventilatory muscle training in patients with chronic obstructive pulmonary disease (COPD). *European Respiratory Journal* 1998;**12** Suppl 28:3S.

Chen 2017 {published data only}

* Chen X, Luo P, Chen Y, Huang Y, Wang K, Hu Y, et al. Effects of combined cycle ergometer and inspiratory muscle training in patients with stable chronic obstructive pulmonary disease. In: *American Journal of Respiratory and Critical Care Medicine*. Vol. 195. 2017:A2861. [DOI: [10.1164/ajrccm-conference.2017.A109](https://doi.org/10.1164/ajrccm-conference.2017.A109)]

Chen X, Xu W-H. Combined use of inspiratory and expiratory threshold pressure training with data monitor in COPD. In: *European Respiratory Journal*. Vol. 52 Suppl 62. 2018:PA4046.

NCT02200549. Effects of combined cycle training and inspiratory muscle training in patients with COPD [Effects of combined cycle training and inspiratory muscle training on exercise performance, health-related quality, dyspnoea, body composition, depressive symptomatology in patients with chronic obstructive pulmonary disease (COPD)]. clinicaltrials.gov/show/nct02200549 (first received 25 July 2014).

Croitoru 2013 {published data only}

Croitoru A, Ionita D, Pele I, Marinescu L, Gologanu D, Dumitrescu A, et al. Inspiratory muscle training with threshold loading in a rehabilitation program of COPD patients. *European Respiratory Journal* 2013;**42 Suppl 57**:774s [P3732].

Del Castillo Otero 1998 {published data only}

Del Castillo-Otero D, Sanchez-Riera H, Ortega-Ruiz F, Cejudo-Ramos P, Toral-Marin J, Elias-Hernandez T, et al. Home respiratory muscle training in COPD. *Archivos de Bronconeumologia* 1998;**34 Suppl 1**:33.

Di Marzo 2000 {published data only}

Di Marzo A, Torrice M, Ciappi G. Inspiratory muscle training and relaxation therapy in advanced COPD patients. *European Respiratory Journal* 2000;**16 Suppl 31**:46s.

Di Marzo 2002 {published data only}

Di Marzo A, Mancuso A, Bevignani G. Breathing reeducation with a pressure threshold device in advanced COPD patients. *American Journal of Respiratory and Critical Care Medicine* 2002;**165 Suppl 8**:A506.

Downes Vogel 2002 {published data only}

Downes Vogel PJ. Effect of adding inspiratory muscle training to a pulmonary rehabilitation program for patients with COPD which includes upper extremity exercises. *American Journal of Respiratory and Critical Care Medicine* 2002;**165 Suppl 8**:A737.

Eastwood 2005 {published data only}

Eastwood P, Hill K, Jenkins S, Philippe D, Shepherd K, Cecins N, et al. High intensity inspiratory muscle training (HIMT) improves dyspnea and health-related quality of life (QoL) in COPD [Abstract]. In: American Thoracic Society International Conference; 2005 May 20-25; San Diego. 2005:[C63] [Poster: H96].

Gething 2001 {published data only}

Gething AD, Davies B, Williams EM. Improvements in cycling endurance following nine weeks of inspiratory muscle training. *Thorax* 2001;**56 Suppl 3**:iii50.

Göhl 2006 {published data only}

Gohl O, Schacher C, Grensemann S, Worth H. Effects of inspiratory muscle training with the RESPIFIT S in addition to an outpatient exercise training program for patients with COPD [Abstract]. *European Respiratory Journal* 2006;**28 Suppl 50**:556s [P3202].

IRCT201104266299N1 {published data only}

IRCT201104266299N1. The effect of inspiratory muscle training in COPD patients. ictrptest.azurewebsites.net/Trial2.aspx?

TrialID=IRCT201104266299N1 (first received 22 September 2011).

IRCT20180205038633N1 {published data only}

IRCT20180205038633N1. The assessment of effectiveness of respiratory muscle training (IMT) and aerobic training in patients with pulmonary disease [The assessment of effectiveness of respiratory muscle training (IMT) and aerobic training on quality of life in patients with chronic obstructive pulmonary disease (COPD)]. ictrptest.azurewebsites.net/Trial2.aspx? TrialID=IRCT20180205038633N1 (first received 21 March 2017).

ISRCTN19258620 {published data only}

ISRCTN19258620. Inspiratory muscle training in patients with chronic obstructive pulmonary disease (COPD). ictrptest.azurewebsites.net/Trial2.aspx? TrialID=ISRCTN19258620 1 March 2000.

Jones 1985 {published data only}

Jones DT, Thomson RJ, Sears MR. Physical exercise and resistive breathing training in severe chronic airways obstruction--are they effective? *European Journal of Respiratory Diseases* 1985;**67**(3):159-66.

Koppers 2004 {published data only}

Koppers R, Vos P, Folgering H. Endurance respiratory muscle training improves outcomes of pulmonary rehabilitation in COPD-patients [Abstract]. In: American Thoracic Society 100th International Conference; 2004 May 21-26; Orlando. 2004:B60 Poster K8.

Liu 1989 {published data only}

Liu Y, Zhao ZS, Yang DC. Inspiratory muscle loading training compared with the breathing gymnastics in patients with COPD. *Chinese Journal of Geriatrics* 1989;**8**(2):112-14.

Manuel Vargas 1995 {published data only}

Manuel Vargas D, Alonso Puig X, Pia de la Maza CM, Morales XP, Vargas BD, Bunout BD, et al. Effects of an inspiratory muscle training program and nutritional support in patients with chronic obstructive lung disease. *Revista Medica de Chile* 1995;**123**:1225-34.

Mendoza 2007 {published data only}

Mendoza L, Levia A, Jover E, Cavada G, Lisboa C. Effect of inspiratory muscle training on lung function, dyspnea, exercise tolerance, and quality of life in COPD patients [Abstract]. *European Respiratory Journal* 2007;**30 Suppl 51**:514s [E3083].

Meshcherykova 2018 {published data only}

Meshcherykova N, Belevsky A, Cherniak A. The effect of inspiratory muscle training on lung hyperinflation in COPD patients. *European Respiratory Journal* 2018;**52 Suppl 62**:PA4139. [DOI: [10.1183/13993003.congress-2018.PA4139](https://doi.org/10.1183/13993003.congress-2018.PA4139)]

NCT01056081 {published data only}

NCT01056081. Inspiratory muscle training in chronic obstructive pulmonary disease [I]. clinicaltrials.gov/show/nct01056081 26 January 2010.

NCT01903772 {published data only}

NCT01903772. Effects of inspiratory muscle training in chronic obstructive pulmonary disease (COPD) (IMTCO). clinicaltrials.gov/show/NCT01903772 19 July 2013.

NCT02392715 {published data only}

NCT02392715. Inspiratory muscle training combined with general exercise training in COPD (IMTGET) [Inspiratory muscle training combined with general exercise training, compared to general exercise training alone in patients with COPD: randomized controlled trial]. clinicaltrials.gov/show/nct02392715 19 March 2015.

NCT02673242 {published data only}

NCT02673242. Inspiratory muscle training for patients with chronic obstructive pulmonary disorder. clinicaltrials.gov/show/NCT02673242 3 February 2016.

NCT03080662 {published data only}

NCT03080662. Impact of inspiratory muscle training on daily physical activity (INAF). clinicaltrials.gov/show/nct03080662 15 March 2017.

NCT03438019 {published data only}

NCT03438019. Inspiratory muscle training in COPD [Inspiratory muscle strength and endurance training in veterans with COPD]. clinicaltrials.gov/show/nct03438019 19 February 2018.

NCT03790410 {published data only}

NCT03790410. The effect of inspiratory muscle training on postural control in chronic obstructive pulmonary disease. clinicaltrials.gov/show/nct03790410 31 December 2018.

Newall 1998 {published data only}

Newall C, Caine MP, Stockley RA, McConnel AK, Hill SL. Inspiratory muscle training using a pressure threshold device in patients with COPD. *European Respiratory Journal* 1998;**12** Suppl 28:267S.

Newall 2000 {published data only}

Newall C, Richardson B, McConnell AK, Stockley RA, Hill SL. Inspiratory muscle training (IMT) as an adjunct to a pulmonary rehabilitation programme in COPD. *American Journal of Respiratory and Critical Care Medicine* 2000;**161**(3 Suppl):A753.

NTR2990 {published data only}

NTR2990. Effects of inspiratory muscle training in patients with chronic obstructive pulmonary disease. ictrptest.azurewebsites.net/Trial2.aspx?TrialID=NTR2990 (first received 18 July 2011).

Pertuze 1994 {published data only}

Pertuze J, Villafranca C, Leiva A, Reppeto P, Cruz E, Lisboa C. Improved quality of life of patients with chronic bronchial obstruction with inspiratory muscle training. *Enfermedades Respir. Cir. Torac* 1994;**10**(4):299.

Ramirez Sarmiento 2000 {published data only}

Ramirez Sarmiento A, Orozo Levi M, Belalcazar V, Mendez R, Barreiro E, Guell R, et al. Structural basis of improved muscle

function after inspiratory training in severe COPD patients. *European Respiratory Journal* 2000;**16** Suppl 31:37s.

Reidi 2005 {published data only}

Reidi C, Toledo A, Ribeiro KP, Silva MI, Costa D. Effects of a respiratory muscle training with or without load in patients with COPD. *Rehabilitar* 2005;**7**(27):4-10.

Valderramas 2009 {published data only}

Valderramas S, Souza MC. Inspiratory muscle training in patients with chronic obstructive pulmonary disease submitted to a pulmonary rehabilitation program [Abstract]. In: European Respiratory Society Annual Congress; 2009 Sep 12-16; Vienna. 2009:[E4293].

Vargas 1995 {published data only}

Vargas M, Puig A, de la Maza M, Morales P, Vargas D, Bunout B, et al. Patients with chronic airflow limitation: effects of the inspiratory muscle training with threshold load valve, built with appropriate technology, associated to nutritional support. *Revista Medica de Chile* 1995;**123**(10):1225-34.

Vargas 1998 {published data only}

Vargas R, Sanchez H, Del Castillo D, Cejudo P, Ortega F, Montemayor T. Impact of ventilatory muscle training on dyspnea, exercise and quality of life in COPD. *Neumosur* 1998;**10**(1):17.

Wang 2004 {published data only}

Wang H, Wan S-Y. Effect of respiratory muscle training on pulmonary function and quality of life in patients with chronic obstructive pulmonary diseases in remission period. *Chinese Journal of Clinical Rehabilitation* 2004;**8**(18):3436-7.

Wanke 1994 {published data only}

Wanke T, Formanek D, Lahrman H, Brath H, Wild M, Wagner C, et al. Effects of combined inspiratory muscle and cycle ergometer training on exercise performance in patients with COPD. *European Respiratory Journal* 1994;**7**(12):2205-11. [DOI: [10.1183/09031936.94.07122205](https://doi.org/10.1183/09031936.94.07122205)]

Weiner 2006a {published data only}

Weiner P, McConnell AK, Magadle R, Beckerman M. The addition of inspiratory muscle training to pulmonary rehabilitation program in patients with significant COPD [Abstract]. *European Respiratory Journal* 2006;**28** Suppl 50:554s [P3194].

Wolstenholme 1998 {published data only}

Wolstenholme RJ, Thomas E, Enright S. An evaluation of the clinical effectiveness of two respiratory muscle training devices in the rehabilitation of patients with chronic airflow obstruction. [Abstract]. *European Respiratory Journal. Supplement* 1998;**12** Suppl 28:353S.

References to ongoing studies
CTRI/2020/11/029226 {published data only}

CTRI/2020/11/029226. Effects of inspiratory muscle training v/s autogenic drainage in hospitalised COPD patient. trialsearch.who.int/Trial2.aspx?TrialID=CTRI/2020/11/029226 19 November 2020.

CTRI/2021/05/033469 {published data only}

CTRI/2021/05/033469. To test the efficacy of inspiratory muscle training device Airofit in reducing breathlessness in COPD patients. www.ctri.nic.in/Clinicaltrials/pmaindet2.php?trialid=54407 1 June 2021.

CTRI201712010952 {published data only}

CTRI201712010952. Effect of breathing exercises to improve the strength of respiratory muscles in COPD population. trialsearch.who.int/Trial2.aspx?TrialID=CTRI/2017/12/010952 21 December 2017.

De Souza 2019 {published data only}

De Souza Y, Medeiros S, Da Silva KM, Macedo J, Condesso D, Figueira B, et al. Does inspiratory muscle training (IMT) reduce depression in patients with COPD? *European Respiratory Journal* 2019;**54 Suppl 63**:PA2197. [DOI: [10.1183/13993003.congress-2019.PA2197](https://doi.org/10.1183/13993003.congress-2019.PA2197)]

Formiga 2020 {published data only}

* Formiga MF, Dosbaba F, Hartman M, Batalik L, Plutinsky M, Brat K, et al. Novel versus traditional inspiratory muscle training regimens as home-based, stand-alone therapies in COPD: protocol for a randomized controlled trial. *International Journal of Chronic Obstructive Pulmonary Disease* 2020;**15**:2147-55. [DOI: [10.2147/COPD.S266234](https://doi.org/10.2147/COPD.S266234)]

NCT04415788. Inspiratory muscle training and COPD [Novel versus traditional inspiratory muscle training regimens as home-based, stand-alone therapies in COPD]. clinicaltrials.gov/show/NCT04415788 4 June 2020.

JPRN-UMIN000039893 {published data only}

JPRN-UMIN000039893. The effectiveness of inspiratory muscle training in patients with chronic obstructive pulmonary disease. trialsearch.who.int/Trial2.aspx?TrialID=JPRN-UMIN000039893 (first received 22 March 2020).

JPRN-UMIN000043099 {published data only}

JPRN-UMIN000043099. Effect of inspiratory muscle Training on diaphragm and exercise tolerance in patients with COPD. trialsearch.who.int/Trial2.aspx?TrialID=JPRN-UMIN000043099 (first received 22 January 2021).

NCT04120142 {published data only}

NCT04120142. Effect of inspiratory muscle training on dyspnoea and exercise tolerance in COPD patients [Effect of inspiratory muscle training during PR on dyspnoea and exercise tolerance in chronic obstructive (COPD)]. clinicaltrials.gov/show/nct04120142 (first received 9 October 2019).

NCT04201522 {published data only}

NCT04201522. The effect of respiratory training on exercise tolerance in COPD (ERTET) [The effect of respiratory training with normocapnic hyperpnea on exercise tolerance in COPD]. clinicaltrials.gov/show/NCT04201522 17 December 2019.

NCT04387318 {published data only}

NCT04387318. Inspiratory muscle training and neuromuscular electrical stimulation in chronic obstructive pulmonary disease [Effects of inspiratory muscle training and neuromuscular

electrical stimulation in patients with chronic obstructive pulmonary disease]. clinicaltrials.gov/show/NCT04387318 13 May 2020.

NCT04802096 {published data only}

NCT04802096. Effects of inspiratory muscle training in addition to pulmonary rehabilitation in patients with moderate to severe COPD exacerbation. clinicaltrials.gov/ct2/show/NCT04802096 17 March 2021.

RBR-10nyzcqc {published data only}

Effects of Inspiratory Muscle Training on Breathlessness, Exercise Capacity and Postural Control in Patients with COPD.. <https://ensaiosclinicos.gov.br/rg/RBR-10nyzcqc> 2021.

RBR 42rmqy {published data only}

RBR 42rmqy. Inspiratory muscle training in chronic obstructive pulmonary disease oxygen-dependent patients: a randomized controlled trial [Effectiveness of a respiratory muscle training program in the ability to exercise exercise capacide, subjective perception of dyspnea, respiratory and peripheral muscle strength and quality of life in patients participating in the home oxygen therapy program of the health state department of the Distrito Federal -: inspiratory muscle training]. ictrptest.azurewebsites.net/Trial2.aspx?TrialID=RBR-42rmqy 29 September 2017.

TCTR20210604001 {published data only}

TCTR20210604001. Is single bout of inspiratory muscle training alter blood pressure and cardio autonomic modulation in COPD patients? : a pilot study. trialsearch.who.int/Trial2.aspx?TrialID=TCTR20210604001 11 January 2021.

Tctr 2022 {published data only}

Effect of inspiratory muscle training on cardiovascular autonomic functions in chronic obstructive pulmonary disease patients. <https://trialsearch.who.int/Trial2.aspx?TrialID=TCTR20220513001> 2022.

Additional references

Agustí 2005

Agustí G, Alvar N. Systemic effects of chronic obstructive pulmonary disease. *Proceedings of the American Thoracic Society* 2005;**2**(4):367-70.

Alma 2018

Alma H, De Jong C, Tsiligianni I, Sanderman R, Kocks J, Van Der Molen T. Clinically relevant differences in COPD health status: systematic review and triangulation. *European Respiratory Journal* 2018;**52 Suppl 62**:PA1168.

American Thoracic Society

COPD Assessment Test (CAT). <https://www.thoracic.org/members/assemblies/assemblies/srn/questionnaires/copd.php>.

Ammous 2022

Ammous O, Feki W, Lotfi T, Khamis AM, Gosselink R, Rebai A, et al. ROB2_IRPG_beta_v7_IMT_COP.xlsm. 10.6084/m9.figshare.20066087 (accessed 14 June 2022) 15 November 2022. [DOI: <https://doi.org/10.6084/m9.figshare.20066087.v1>]

Andersen 1979

Andersen JB, Dragsted L, Kann T, Johansen SH, Nielsen KB, Karbo E, et al. Resistive breathing training in severe chronic obstructive pulmonary disease. A pilot study. *Scandinavian Journal of Respiratory Diseases* 1979;**60**(3):151-6.

Anzueto 2017

Anzueto A, Miravittles M. Pathophysiology of dyspnea in COPD. *Postgraduate Medicine* 2017;**129**(3):366-74.

Araújo 2017

Araújo Oliveira AL Andrade L, Marques A. Minimal clinically important difference and predictive validity of the mMRC and mBorg in acute exacerbations of COPD. *European Respiratory Journal* 2017/09/01;**50** Suppl **61**:PA4705.

Beaumont 2015

Beaumont M, Mialon P, Le Ber-Moy C, Lochon C, Péran L, Pichon R, et al. Inspiratory muscle training during pulmonary rehabilitation in chronic obstructive pulmonary disease: a randomized trial. *Chronic Respiratory Disease* 2015;**12**(4):305-12.

Beaumont 2018a

Beaumont M, Forget P, Couturaud F, Reychler G. Effects of inspiratory muscle training in COPD patients: a systematic review and meta-analysis. *Clinical Respiratory Journal* 2018;**12**(7):2178-88. [PMID: 29665262]

Bégin 1991

Bégin P, Grassino A. Inspiratory muscle dysfunction and chronic hypercapnia in chronic obstructive pulmonary disease. *American Review of Respiratory Disease* 1991;**143**(5 Pt 1):905-12.

Belman 1980

Belman M-J, Mittman C. Ventilatory muscle training improves exercise capacity in chronic obstructive pulmonary disease patients. *American Review of Respiratory Disease* 1980;**121**(2):273-80.

Belman 1994

Belman MJ, Botnick WC, Nathan SD, Chon KH. Ventilatory load characteristics during ventilatory muscle training. *American Journal of Respiratory and Critical Care Medicine* 1994;**149**(4 Pt 1):925-9.

Berg 2016

Berg K, Wright JL. The pathology of chronic obstructive pulmonary disease: progress in the 20th and 21st centuries. *Archives of Pathology and Laboratory Medicine* 2016;**140**(12):1423-8. [PMID: 27922768]

Bestall 1999

Bestall JC, Paul EA, Garrod R, Garnham R, Jones PW, Wedzicha JA. Usefulness of the Medical Research Council (MRC) dyspnoea scale as a measure of disability in patients with chronic obstructive pulmonary disease. *Thorax* 1999;**54**(7):581-586.

Boles 2007

Boles J-M, Bion J, Connors A, Herridge M, Marsh B, Melot C, et al. Weaning from mechanical ventilation. *European Respiratory Journal* 2007;**29**(5):1033.

Borg 1982

Borg Gunnar aV. Psychophysical bases of perceived exertion. *Medicine & Science in Sports & Exercise* 1982;**14**(5):377-381.

Caron 2011

Caron MA, Debigaré R, Dekhuijzen PN, Maltais F. Diaphragm and skeletal muscle dysfunction in COPD [L'atteinte du diaphragme et du quadriceps dans la BPCO: une manifestation systémique de cette maladie?]. *Revue des Maladies Respiratoires* 2011;**28**(10):1250-64.

Caruso 2015

Caruso P, Albuquerque AL, Santana PV, Cardenas LZ, Ferreira JG, Prina E, et al. Diagnostic methods to assess inspiratory and expiratory muscle strength. *Jornal Brasileiro de Pneumologia* 2015;**41**(2):110-23.

Cazzola 2015

Cazzola M, Hanania NA, MacNee W, Rudell K, Hackford C, Tamimi N. A review of the most common patient-reported outcomes in COPD--revisiting current knowledge and estimating future challenges. *International Journal of Chronic Obstructive Pulmonary Disease* 2015;**10**:725-38.

Charususin 2016

Charususin N, Gosselink R, McConnell A, Demeyer H, Topalovic M, Decramer M, et al. Inspiratory muscle training improves breathing pattern during exercise in COPD patients. *European Respiratory Journal* 2016;**47**(4):1261-4.

Clanton 2009

Clanton TL, Levine S. Respiratory muscle fiber remodeling in chronic hyperinflation: dysfunction or adaptation? *Journal of Applied Physiology* 2009;**107**(1):324-35.

Clark 1996

Clark CJ, Cochrane L, Mackay E. Low intensity peripheral muscle conditioning improves exercise tolerance and breathlessness in COPD. *European Respiratory Journal* 1996;**9**(12):2590.

Cochrane Airways 2022

Cochrane Airways Trials Register. airways.cochrane.org/trials-register (accessed 19 October 2022).

Cohen 1988

Jacob Cohen. Statistical power analysis for the behavioral sciences. Lawrence Erlbaum Associates, 1988.

Costanzo 2019

Costanzo LS. Physiology. 7th edition. Wolters Kluwer, 2019. [LCCN 2017054852 | ISBN 9781496367617]

Covidence [Computer program]

Veritas Health Innovation Covidence. Melbourne, Australia: Veritas Health Innovation, accessed 19 October 2020. Available at covidence.org.

Crowe 2005

Crowe J, Reid WD, Geddes EL, O'Brien K, Brooks D. Inspiratory muscle training compared with other rehabilitation interventions in adults with chronic obstructive pulmonary disease: a systematic literature review and meta-analysis. *COPD* 2005;**3**:319-29.

Deeks 2021

Deeks JJ, Higgins JP, Altman DG editor(s). Chapter 10: Analysing data and undertaking meta-analyses. In: Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.

Doucet 2004

Doucet M, Debigaré R, Joanisse DR, Côté C, Leblanc P, Grégoire J, et al. Adaptation of the diaphragm and the vastus lateralis in mild-to-moderate COPD. *European Respiratory Journal* 2004;**24**(6):971-9.

Figueiredo 2020

Figueiredo RI, Azambuja AM, Cureau FV, Sbruzzi G. Inspiratory muscle training in COPD. *Respiratory Care* 2020;**65**(8):1189.

Geddes 2005

Geddes EL, Reid DW, Crowe J, O'Brien K, Brooks D. Inspiratory muscle training in adults with chronic obstructive pulmonary disease: a systematic review. *Respiratory Medicine* 2005;**99**(11):1440-58.

Geddes 2008

Geddes EL, O'Brien K, Reid WD, Brooks D, Crowe J. Inspiratory muscle training in adults with chronic obstructive pulmonary disease: an update of a systematic review. *Respiratory Medicine* 2008;**102**(12):1715-29.

GOLD 2022

Global Initiative for Chronic Obstructive Lung Disease. Global Initiative for Chronic Obstructive Lung Disease. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease (2022 report). <https://goldcopd.org/2022-gold-reports-2/> (accessed prior to 1 November 2022).

Gosselink 2011

Gosselink R, De Vos J, Van den Heuvel SP, Segers J, Decramer M, Kwakkel G. Impact of inspiratory muscle training in patients with COPD: what is the evidence? *European Respiratory Journal* 2011;**37**(2):416-25.

GRADE Handbook

Schünemann H, Brożek J, Gguyatt G, Oxman A, editor(s). Handbook for grading the quality of evidence and the strength of recommendations using the GRADE approach (updated October 2013). GRADE Working Group, 2013. Available from gdt.guidelinedevelopment.org/app/handbook/handbook.html.

GRADEpro GDT [Computer program]

McMaster University and Evidence Prime GRADEpro Guideline Development Tool [Software]. Version accessed October 2022.

Hamilton (ON): McMaster University and Evidence Prime, 2022. Available at gradepro.org.

Higgins 2003

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**:557-60.

Higgins 2021

Higgins JP, Savović J, Page MJ, Elbers RG, Sterne JA, editor(s). Chapter 8: Assessing risk of bias in a randomized trial. In: Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.

Hill 2010

Hill K, Cecins NM, Eastwood PR, Jenkins SC. Inspiratory muscle training for patients with chronic obstructive pulmonary disease: a practical guide for clinicians. *Archives of Physical Medicine and Rehabilitation* 2010;**91**(9):1466-70. [PMID: 20801269]

Holland 2014

Holland Anne E, Spruit Martijn A, Troosters Thierry, Puhan Milo A, Pepin Véronique, Saey Didier, McCormack Meredith C, Carlin Brian W, Sciurba Frank C, Pitta Fabio, Wanger Jack, MacIntyre Neil, Kaminsky David A, Culver Bruce H, Revill Susan M, Hernandez Nidia A, Andrianopoulos Vasileios, Camillo Carlos Augusto, Mitchell Katy E, Lee Annemarie L, Hill Catherine J, Singh Sally J. An official European Respiratory Society/American Thoracic Society technical standard: field walking tests in chronic respiratory disease. *Eur Respir J* 2014/12/01;**44**(6):1428.

Iwakura 2020

Iwakura M, Okura K, Kubota M, Sugawara K, Kawagoshi A, Takahashi H, et al. Estimation of minimal clinically important difference for quadriceps and inspiratory muscle strength in older outpatients with chronic obstructive pulmonary disease: a prospective cohort study. *Physical Therapy Research* 2020;**24**(1):35-42.

Jones 1992

Jones PW, Quirk FH, Baveystock CM, Littlejohns P. A self-complete measure of health status for chronic airflow limitation. The St. George's Respiratory Questionnaire. *The American Review of Respiratory Disease* 1992;**145**(6):1321-1327.

Jones 2009

Jones PW, Harding G, Berry P, Wiklund I, Chen W-H, Kline Leidy N. Development and first validation of the COPD Assessment Test. *The European Respiratory Journal* 2009;**34**(3):648-654.

Langer 2015

Langer D, Charusisin N, Jácome C, Hoffman M, McConnell A, Decramer M, et al. Efficacy of a novel method for inspiratory muscle training in people with chronic obstructive pulmonary disease. *Physical Therapy* 2015;**95**(9):1264-73.

Laveneziana 2019

Laveneziana P, Albuquerque A, Aliverti A, Babb T, Barreiro E, Dres M, et al. ERS statement on respiratory muscle testing at rest and during exercise. *European Respiratory Journal* 2019;**53**(6):1801214.

Leith 1976

Leith DE, Bradley M. Ventilatory muscle strength and endurance training. *Journal of Applied Physiology* 1976;**41**(4):508-16.

Lötters 2002

Lötters F, Van Tol B, Kwakkel G, Gosselink R. Effects of controlled inspiratory muscle training in patients with COPD: a meta-analysis. *European Respiratory Journal* 2002;**20**(3):570-7.

Mahler 2005

Mahler DA, Witek TJ Jr. The MCID of the transition dyspnea index is a total score of one unit. *COPD* 2005;**2**(1):99-103.

Maillard 1998

Maillard JO, Burdet L, Van Melle G, Fitting JW. Reproducibility of twitch mouth pressure, sniff nasal inspiratory pressure, and maximal inspiratory pressure. *European Respiratory Journal* 1998;**11**(4):901-5.

Maltais 1997

Maltais F, LeBlanc P, Jobin J, Bérubé C, Bruneau J, Carrier L. Intensity of training and physiologic adaptation in patients with chronic obstructive pulmonary disease. *American Journal of Respiratory and Critical Care Medicine* 1997;**155**(2):555-61.

McCarthy 2015

McCarthy B, Casey D, Devane D, Murphy K, Murphy E, Lacasse Y. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2015, Issue 2. Art. No: CD003793. [DOI: [10.1002/14651858.CD003793.pub3](https://doi.org/10.1002/14651858.CD003793.pub3)]

McConnell 2004

McConnell AK, Romer LM. Respiratory muscle training in healthy humans: resolving the controversy. *International Journal of Sports Medicine* 2004;**25**(04):284-93.

McGowan 2016

McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS peer review of electronic search strategies: 2015 guideline statement. *Journal of Clinical Epidemiology* 2016;**75**:40-6. [DOI: [10.1016/j.jclinepi.2016.01.021](https://doi.org/10.1016/j.jclinepi.2016.01.021)]

McGrath 2020

McGrath S, Zhao X-F, Steele R, Thombs BD, Benedetti A, DEPRESSion Screening Data (DEPRESSD) Collaboration. Estimating the sample mean and standard deviation from commonly reported quantiles in meta-analysis. *Statistical Methods in Medical Research* 2020;**29**(9):2520-37.

Menzes 2018

Menzes KK, Nascimento LR, Avelino PR, Polese JC, Salmela LF. A review on respiratory muscle training devices. *Journal of Pulmonary and Respiratory Medicine* 2018;**08**(02):451.

Meyer 2013

Meyer FJ, Borst MM, Buschmann HC, Ewert R, Friedmann-Bette B, Ochmann U, et al. Exercise testing in respiratory medicine. *Pneumologie* 2013;**67**:16-34. [DOI: [10.1055/s-0032-1325901](https://doi.org/10.1055/s-0032-1325901)]

Moher 2009

Moher D, Liberati A, Tetzlaff J, Altman D. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine* 2009;**6**(7):e1000097. [DOI: [10.1371/journal.pmed.1000097](https://doi.org/10.1371/journal.pmed.1000097)]

Nickerson 1982

Nickerson BG, Keens TG. Measuring ventilatory muscle endurance in humans as sustainable inspiratory pressure. *Journal of Applied Physiology* 1982;**52**(3):768-72. [DOI: [10.1152/jappl.1982.52.3.768](https://doi.org/10.1152/jappl.1982.52.3.768)]

O'Brien 2008

O'Brien K, Geddes E, Reid W, Brooks D, Crowe J. Inspiratory muscle training compared with other rehabilitation interventions in chronic obstructive pulmonary disease: a systematic review update. *Journal of Cardiopulmonary Rehabilitation and Prevention* 2008;**28**(2):128-41.

O'Donnell 2006

O'Donnell DE. Hyperinflation, dyspnea, and exercise intolerance in chronic obstructive pulmonary disease. *Proceedings of the American Thoracic Society* 2006;**3**(2):180-4.

O'Donnell 2007

O'Donnell DE, Banzett RB, Carrieri-Kohlman V, Casaburi R, Davenport PW, Gandevia SC, et al. Pathophysiology of dyspnea in chronic obstructive pulmonary disease. *Proceedings of the American Thoracic Society* 2007;**4**(2):145-68.

Ottenheijm 2008

Ottenheijm CA, Heunks LM, Dekhuijzen RP. Diaphragm adaptations in patients with COPD. *Respiratory Research* 2008;**9**(1):12.

Padula 2006

Padula CA, Yeaw E. Inspiratory muscle training: integrative review. *Research and Theory for Nursing Practice* 2006;**20**(4):291-304.

Papandrinopoulou 2012

Papandrinopoulou D, Tzouda V, Tsoukalas G. Lung compliance and chronic obstructive pulmonary disease. *Pulmonary Medicine* 2012;**2012**:6. [Article ID 542769]

Pessoa 2014

Sclauser Pessoa IM, Franco Parreira V, Fregonezi GA, Sheel AW, Chung F, Reid WD. Reference values for maximal inspiratory pressure: a systematic review. *Canadian Respiratory Journal* 2014;**21**:43-50.

Puhan 2011

Puhan MA, Chandra D, Mosenifar Z, Ries A, Make B, Hansel NN, et al. The minimal important difference of exercise tests in severe COPD. *European Respiratory Journal* 2011;**37**(4):784.

Ramos 2014

Ramos FL, Krahnke JS, Kim V. Clinical issues of mucus accumulation in COPD. *International Journal of Chronic Obstructive Pulmonary Disease* 2014;**9**:139-50.

Review Manager 2020 [Computer program]

The Cochrane Collaboration Review Manager 5 (RevMan 5). Version 5.4. Copenhagen: The Cochrane Collaboration, 2020.

RevMan Web 2022 [Computer program]

The Cochrane Collaboration Review Manager Web (RevMan Web). Version 4.16.2. Copenhagen: The Cochrane Collaboration, 2022. Available at revman.cochrane.org.

Ries 2005

Ries AL. Minimally clinically important difference for the UCSD shortness of breath questionnaire, Borg Scale, and visual analog scale. *COPD* 2005;**2**(1):105-10.

Rochester 2015

Rochester CL, Vogiatzis I, Holland AE, Lareau SC, Marciniuk DD, Puhan MA, et al. An official American Thoracic Society/European Respiratory Society policy statement: enhancing implementation, use, and delivery of pulmonary rehabilitation. *American Journal of Respiratory and Critical Care Medicine* 2015;**192**(11):1373-86. [PMID: 26623686]

Salito 2015

Salito C, Luoni E, Aliverti A. Alterations of diaphragm and rib cage morphometry in severe COPD patients by CT analysis. In: 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society; 2015 Aug 25-29; Milan. 2015:6390-3.

Schultz 2018

Schultz K, Jelusic D, Wittmann M, Krämer B, Huber V, Fuchs S, et al. Inspiratory muscle training does not improve clinical outcomes in 3-week COPD rehabilitation: results from a randomised controlled trial. *European Respiratory Journal* 2018;**51**(1):1702000. [PMID: 29371382]

Schünemann 2021

Schünemann HJ, Higgins JP, Vist GE, Glasziou P, Akl EA, Skoetz N, et al. Chapter 14: Completing 'Summary of findings' tables and grading the certainty of the evidence. In: Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.

Shoemaker 2009

Shoemaker MJ, Donker S, Lapoe A. Inspiratory muscle training in patients with chronic obstructive pulmonary disease: the state of the evidence. *Cardiopulmonary Physical Therapy Journal* 2009;**20**(3):5-15.

Smith 1992

Smith K, Cook D, Guyatt GH, Madhavan J, Oxman AD. Respiratory muscle training in chronic airflow limitation: a meta-analysis. *American Review of Respiratory Disease* 1992;**145**(3):533-9.

Spruit 2013

Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C, et al. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *American Journal of Respiratory and Critical Care Medicine* 2013;**188**(8):e13-64.

Sterne 2019

Sterne JA, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;**366**:l4898. [PMID: 31462531]

WebPlotDigitizer [Computer program]

WebPlotDigitizer version 4.4. automeris.io/WebPlotDigitizer, 2010. [EMAIL ADDRESS: ankitrohatgi@hotmail.com]

Welling 2015

Welling Jorrit BA, Hartman Jorine E, Ten Hacken NH T, Klooster K, Slebos DJ. The minimal important difference for the St George's Respiratory Questionnaire in patients with severe COPD. *European Respiratory Journal* 2015;**46**(6):1598.

WHO 2013

World Health Organization. Projections of mortality and causes of death, 2015 and 2030. www.who.int/healthinfo/global_burden_disease/projections2015_2030/en/ (accessed prior to 14 May 2021).

WHO 2018

World Health Organization. The top 10 causes of death. www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death (accessed prior to 14 May 2021).

WHO 2020

World Health Organization. WHO burden of COPD. www.who.int/respiratory/copd/burden/en/ (accessed 8 March 2020). [<https://www.who.int/respiratory/copd/burden/en/>]

Wijkstra 1994

Wijkstra PJ, TenVergert EM, Van Altena R, Otten V, Postma DS, Kraan J, Koëter GH. Reliability and validity of the chronic respiratory questionnaire (CRQ).. *Thorax* 1994;**49**(5):465-467.

Wood 2017

Wood KL. Tests of respiratory muscle function - pulmonary disorders. www.msmanuals.com/professional/pulmonary-disorders/tests-of-pulmonary-function-pft/tests-of-respiratory-muscle-function (accessed prior to 6 September 2020).

Wu 2017

Wu W, Zhang X, Lin L, Ou Y, Li X, Guan Li, et al. Transdiaphragmatic pressure and neural respiratory drive measured during inspiratory muscle training in stable patients with chronic obstructive pulmonary disease. *International Journal of Chronic Obstructive Pulmonary Disease* 2017;**12**:773-81.

References to other published versions of this review
Ammous 2020

Ammous O, Feki W, Lotfi T, Khamis AM, Rebai A, Kammoun S.
 Inspiratory muscle training, with or without concomitant
 pulmonary rehabilitation, for chronic obstructive pulmonary

disease (COPD). *Cochrane Database of Systematic Reviews* 2020,
 Issue 11. Art. No: CD013778. [DOI: [10.1002/14651858.CD013778](https://doi.org/10.1002/14651858.CD013778)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]

Abedi Yekta 2019
Study characteristics

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics PR+IMT <ul style="list-style-type: none"> • N (randomized/analyzed): 17/15 • Loss to follow-up or excluded: 2 • Age, mean (SD) in years: 51.33 (10.4) • Gender (M/F): 9/6 • BMI, mean (SD), kg/m²: 27.6 (3.7) PR <ul style="list-style-type: none"> • N (randomized/analyzed): 17/14 • Loss to follow-up or excluded: 3 • Age, mean (SD) in years: 53.5 (10.37) • Gender (M/F): 8/6 • BMI, mean (SD), kg/m²: 27.39 (5.11) IMT <ul style="list-style-type: none"> • N (randomized/analyzed): 17/16 • Loss to follow-up or excluded: 1 • Age, mean (SD) in years: 51.88 (9.05) • Gender (M/F): 7/9 • BMI, mean (SD), kg/m²: 27.05 (4.53) Control/sham <ul style="list-style-type: none"> • N (randomized/analyzed): 17/15 • Loss to follow-up or excluded: 2 • Age, mean (SD) in years: 55.67 (11.08) • Gender (M/F): 7/8 • BMI, mean (SD), kg/m²: 25.98 (4.1) Overall <ul style="list-style-type: none"> • N (randomized/analyzed): 68/60 • Loss to follow-up or excluded: 8 • Age, mean (SD) in years: 53.07 (10.11) • Range age (min, max): 32,70 • Gender (M/F): 31/29

Abedi Yekta 2019 (Continued)

- COPD stage (GOLD): moderate to severe

Included criteria

- Stage 2 or 3 COPD according to the GOLD criteria;
- Age range 30-70 years;
- No previous known diseases, such as heart disease (e.g. congestive heart failure and coronary artery disease), renal disease (e.g. end-stage renal disease and chronic renal failure), or liver disease (e.g. hepatic cirrhosis and hepatic cancer); no history of known pulmonary diseases, such as lung cancer and pleural disease; no history of musculoskeletal diseases (e.g. myasthenia gravis) or restrictive deformities of the lungs;
- Lack of severe limitations in the limbs inhibiting aerobic exercise; no pulmonary surgery in the past 12 months;
- No recent fracture of the ribs in the past 6 months;
- No history of psychotropic diseases; and no use of drugs, alcohol, or psychiatric drugs.

Excluded criteria

- Exacerbation of the disease during the study;
- Need for long-term oxygen therapy for more than 15 h/d;
- The occurrence of complications, such as pneumothorax or diseases exacerbating and disrupting the treatment.

Interventions	<p>Intervention characteristics</p> <p>PR: consisted of aerobic exercise performed on a treadmill ergometer, at 40%-60% of the heart rate, 2 d/week, for 8 weeks and 40 min/session.</p> <p>IMT: conducted 2 d/week for 8 weeks, 15 min/session with Powerbreathe KH4. Each session lasted 15 min, all sessions were supervised, and the training load was set at 40%-60% of S-Index.</p> <p>PR+IMT: consisted of a combination of PR and IMT protocols.</p> <p>Control: no intervention received by this group.</p>
Outcomes	HRQoL: SGRQ (Total)
Identification	<p>Sponsorship source: Shahid Beheshti University of Medical Sciences</p> <p>Country: Iran</p> <p>Setting: Hospital Hussein Department of Physiotherapy</p> <p>Author's name: Saeed Rezaei</p> <p>Institution: Department of Sports Medicine, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran</p> <p>Email: saeedrezaee1394@yahoo.com</p> <p>Clinical trial register: https://www.irct.ir/trial/29724</p>
Notes	A change of 4 units in SGRQ score was considered significant.

Bavarsad 2015

Study characteristics

Methods	Study design: RCT
---------	--------------------------

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

Bavarsad 2015 (Continued)

Study grouping: parallel-group

Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 20/15 • Loss to follow-up or excluded: 5 • Age mean (SD) in years: 58.8 (6.82) • Gender (M/F): 13/2 • BMI mean (SD), kg/m²: 24.88 (5.02) <p>Control/Sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 20/15 • Loss to follow-up or excluded: 5 • Age mean (SD) in years: 54.2 (8.09) • Gender (M/F): 14/1 • BMI mean (SD), kg/m²: 24.97 (4.72) <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 40/30 • Loss to follow-up or excluded: 10 • Age range (min, max): 45, 65 • Gender (M/F): 27/3 • COPD stage (GOLD): mild to severe <p>Included criteria</p> <ul style="list-style-type: none"> • Patients with spirometric evidence of significant chronic airflow limitation (FEV₁/FVC < 70%pred) with mild to very severe COPD diagnosis according to the GOLD criteria; • Age range 45 - 65 years; • No history of PR; • Established treatment plan 1 month prior to the study. <p>Excluded criteria</p> <ul style="list-style-type: none"> • BMI > 35; • Comorbid conditions such as diabetes, musculoskeletal disorders, cardiovascular diseases; and neurological diseases that can affect the results of 6MWD; • Having exacerbation for 1 month prior to the study; • A history of long-term oxygen therapy; • A history of spontaneous pneumothorax and rib fracture.
Interventions	<p>Intervention characteristics</p> <p>IMT</p> <p>The training consisted of unsupervised sessions, 15 min/d, 6 d/week for 8 weeks. The device used was an incentive spirometer (Respivol), at a load equal to or more than the initial inspiratory volume. The researchers were informed of the training sessions through phone calls during the 8 weeks. A checklist, which was designed to be completed on weekdays, was prepared for the participants so that they could mark the relevant day after a training session</p> <p>Control</p> <p>No intervention was received by this group.</p>
Outcomes	Dyspnea: Borg

Bavarsad 2015 (Continued)

- Notes: dyspnea was assessed at the beginning and the end of the 6MWD

Functional exercise capacity: 6MWD

Respiratory function: FEV1 (%pred)

Respiratory function: FEV1 (L)

Identification

Sponsorship source: The Deputy of Research Affairs at the Ahvaz Jundishapur University of Medical Sciences

Country: Iran

Setting: Specialized Pulmonary Clinic of Ahvaz

Author's name: Esmaeil Eidani

Institution: Pulmonary Unit, Department of Medicine, Ahvaz Jundishapur University of Medical Sciences

Email: esmaileidani@gmail.com

Address: Ahzav, Iran

Notes

Beaumont 2015

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Subgroup analysis: FEV1 > or ≤ 50% of the predicted value

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 16/16
- Loss to follow-up or excluded: 0
- Age, mean (SD), in years: 62 (10)
- Gender (M/F): 4/12
- BMI mean (SD), kg/m²: 27.3 (4.2)

PR

- N (randomized/analyzed): 18/16
- Loss to follow-up or excluded: 0
- Age, mean (SD), in years: 61 (8)
- Gender (M/F): 7/11
- BMI mean (SD), kg/m²: 26.8 (6.0)

Overall

- N (randomized/analyzed): 34/32
- Loss to follow-up or excluded: 0
- Gender (M/F): 11/23

Beaumont 2015 (Continued)

- COPD stage (GOLD): moderate to very severe

Included criteria

COPD diagnosed according to [American Thoracic Society](#)/European Respiratory Society criteria;

Plmax > 60 cmH₂O at admission.

Excluded criteria

- Previous pneumonectomy or lobectomy in the past 6 months;
- Impossibility to measure IC at the end of the 6MWD;
- The incapacity to follow the standard rehabilitation program;
- The absence of written informed consent.

Pretreatment: FEV 1 was lower in the IMT group; the Borg scale was higher in the control group.

Interventions

Intervention characteristics

PR+IMT

- PR: conducted for 3 weeks, 5 d/week, and included aerobic exercise on a cycle ergometer and a treadmill (30 min/d each), strengthening of lower and upper limb muscle groups, therapeutic educational program, aerobic gymnastics in groups, smoking cessation program, and socio-psychological and dietary advice.
- IMT: performed daily in 2 sessions of 15 min each, 5 times/week, for 3 weeks, supervised by a physiotherapist. The participants had to breathe slowly with an increased tidal volume. A threshold inspiratory muscle trainer (Threshold IMT1) was used at 40% of Plmax. The intensity was not modified during the program.

PR

Participants in this group received only the standardised PR program.

Outcomes

Dyspnea: Borg

Dyspnea: MDP

- Unpleasantness
- Sensory intensity
- Muscle work/effort
- Not enough air/smother/air hunger
- Mental effort/concentrate
- Tight/constricted
- Breathing a lot (rapid, deep, and heavy)
- Crush
- Depression
- Satisfaction
- Anxiety
- Frustration
- Anger
- Happiness
- Fear

Functional exercise capacity: 6MWD

Respiratory muscle strength: Plmax

Identification

Sponsorship source: EA3878 (G.E.T.B.O.), CIC INSERM 0502, University Hospital of Brest

Beaumont 2015 (Continued)

Country: France

Setting: Pulmonary Rehabilitation Unit, Morlaix Hospital Centre, European University of Occidental Brittany

Author's name: Marc Beaumont

Institution: Pulmonary Rehabilitation Unit, Morlaix Hospital Centre, European University of Occidental Brittany

Email: marc.beaumont@univ-brest.fr

Address: Morlaix29672, Cedex, France

Clinical trial register: NCT01545011

Notes

P-value of the change from baseline is from the adjusted analysis.

Beaumont 2018

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Subgroup analysis: P_{lmax} (> or ≤ 60 cmH₂O)

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 74/74
- Loss to follow-up or excluded: 1
- Age, mean (SD) in years: 62.2 (8.0)
- Gender (M/F): 44/30
- BMI, mean (SD), kg/m²: 26.2 (5.9)

PR

- N (randomized/analyzed): 75/75
- Loss to follow-up or excluded: 1
- Age, mean (SD) in years: 65.9 (8.9)
- Gender (M/F): 50/25
- BMI, mean (SD), kg/m²: 24.7 (5.9)

Overall

- N (randomized/analyzed): 149
- Loss to follow-up excluded: 2
- Gender (M/F): 94/55
- COPD stage (GOLD): severe to very severe

Included criteria

- Severe or very severe COPD diagnosed according to ATS/ERS criteria at admission (FEV₁ < 50%pred)

Excluded criteria

Beaumont 2018 (Continued)

- Previous pneumonectomy or lobectomy in the past 6 months;
- Spontaneous risk of pneumothorax or rib fracture;
- Incapacity to follow a standard rehabilitation program (locomotor deficits, acute cardiac failure and acute exacerbation of COPD at the beginning of the program);
- The absence of written informed consent.

Pretreatment: 6 patients with lobectomy or pneumonectomy in the control group were included.

Interventions	<p>Intervention characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • PR: conducted over 4 weeks, 5 d/week and included aerobic exercise on a cycle ergometer and a treadmill (each for 30 min/day), strengthening of lower and upper limb muscle groups, a therapeutic, educational program, aerobic gymnastics in groups, a smoking cessation program and socio-psychological and dietary advice. • IMT: consisted of 2 sessions of 15 min each, supervised by a physiotherapist, 5 times/week, over 4 weeks. The patients had to breathe slowly with an increased tidal volume; after 10 inspirations, they could have a break by breathing at rest for a short time. The cycle of 10 inspirations was repeated 15 times. The inspiratory muscle training was performed using a threshold inspiratory muscle trainer (PowerBreathe Medic; PowerBreathe, Southam, UK) at a resistance of 50% of P_Imax. The intensity was increased (+10%) after 10 days of training to reach 60% of the initial P_Imax. <p>PR: participants in this group received only the standardised PR program.</p>
Outcomes	<p>Dyspnea: Borg</p> <p>Dyspnea: mMRC</p> <p>Dyspnea: MDP:</p> <ul style="list-style-type: none"> • Unpleasantness • Sensory intensity • Muscle work/effort • Not enough air/smother/air hunger • Tight/constricted • Mental effort/ concentrate • Breathing a lot (rapid, deep, and heavy) • Depression • Anxiety • Frustration • Anger • Fear <p>Functional exercise capacity: 6MWD</p> <p>HRQoL: SGRQ</p> <ul style="list-style-type: none"> • Symptoms • Activity • Impact • Total <p>Respiratory muscle strength: P_Imax</p>
Identification	<p>Country: France</p> <p>Setting: Rehabilitation programme unit of Centre Hospitalier des Pays de Morlaix</p> <p>Author's name: Marc Beaumont</p>

Beaumont 2018 (Continued)

Institution: Pulmonary Rehabilitation Unit, Morlaix HospitalCentre, European University of Occidental Brittany

Email: marc.beaumont@univ-brest.fr

Address: 29672 Morlaix CEDEX, France

Clinical trial register: NCT02074813

Notes

Borg and MDP scales were conducted at the end of 6MWD.
 Subgroup analysis according to P_{lmax} (> or ≤ 60 cmH₂O) was conducted. See supplementary table.

Beckerman 2005
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): 21/17
- Loss to follow-up or excluded: 4
- Age, mean (SD), in years: 67.7 (16.49)
- Gender (M/F): 17/4

Control/sham

- N (randomized/analyzed): 21/14
- Loss to follow-up or excluded: 7
- Age, mean (SD), in years: 66.9 (15.12)
- Gender (M/F): 15/6

Overall

- N (randomized/analyzed): 42/31
- Loss to follow-up or excluded: 11
- Gender (M/F): 32/10
- COPD stage: severe to very severe

Included criteria

- Spirometric evidence of significant chronic airflow limitation (FEV₁ < 50% of predicted, FEV₁/FVC < 70% of predicted) with a diagnosis of COPD according to the criteria of the ATS were recruited from the community.
- The patients were all new to an IMT program, and none were receiving additional regular exercise or dietary supplements.

Excluded criteria

- Patients with cardiac disease and poor compliance and needing supplemental oxygen

Interventions

Intervention characteristics

Beckerman 2005 (Continued)

IMT: participants trained daily in 2 sessions of 15 min each, 6 times/week for 12 months. The training was performed using a threshold inspiratory muscle trainer (POWERbreathe; Gaiam Ltd; Southam, Warwickshire, UK). The participants started breathing at a resistance that required generation of 15% of Pimax for 1 week. The load was then increased incrementally, 5% to 10% each session, to reach 60% of Pimax at the end of the first month. IMT was then continued at 60% of the Pimax adjusted monthly to the new Pimax achieved. The training was conducted in a rehabilitation center for 1 month under the supervision of a respiratory therapist followed by home training, verified by a respiratory therapist daily by phone and once weekly by a personal visit, for the next 11 months.

Control/sham: this group trained with the same protocol at a load equal to 7 cm H₂O.

Outcomes	Dyspnea: Borg Functional exercise capacity: 6MWD HRQoL: SGRQ Respiratory muscle strength: PImax
Identification	<p>Country: Israel</p> <p>Setting: Home</p> <p>Author's name: Paltiel Weiner</p> <p>Institution: Department of Medicine A, Hillel Yaffe Medical Center</p> <p>Email: weiner@hillel-yaffe.health.gov.il</p> <p>Address: Hadera, Israel 38100</p>
Notes	The study reported data as mean and SE, we computed SD for the baseline characteristics. The SE reported in the text of the trial is less than shown in the graph.

Belman 1988
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/8 • Age, mean (SD), in years: 64 (8.4) • Gender (M/F): 4/4 <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/9 • Age, mean (SD), in years: 64 (9.0) • Gender (M/F): 6/3 <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 20/17 • Loss to follow-up or excluded: 3

Belman 1988 (Continued)

- Gender (M/F): 10/7
- COPD stage: moderate to severe

Included criteria

- Presence of COPD as defined by the ATS, and improvement in FEV1 < 20% after inhaled isoproterenol

Excluded criteria

- Evidence of coronary artery disease, cardiac arrhythmias, congestive heart failure, and orthopedic problems such as shoulder girdle and spinal abnormalities, which would interfere with the performance of the breathing manoeuvres

Interventions	<p>Intervention characteristics</p> <p>IMT: the training protocol consisted of unsupervised sessions, 2 sessions/d, 7/week for 6 weeks. Each session lasted 15 min, using a Pflex device (the breathing pattern was controlled), and the training load was as tolerated.</p> <p>Control/sham: the control group received similar training at a load of around 7.5-10 cmH₂O.</p>
Outcomes	<p>Respiratory muscle strength: P_Imax (FRC)</p> <p>Respiratory muscle endurance:</p> <ul style="list-style-type: none"> • MVV • P_{th}max <p>Respiratory function: FEV1 (L)</p>
Identification	<p>Country: USA</p> <p>Setting: hospital</p> <p>Author's name: Michael J Belman</p> <p>Institution: Division of Pulmonary Medicine, Cedars-Sinai Medical Center, and The University of California</p> <p>Address: Los Angeles, California 90048</p>
Notes	

Berry 1996

Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 8/7 • Loss to follow-up or excluded: 1 • Age, mean (SD), in years: 67.0 (3.39) • Gender (M/F): 4/4

Berry 1996 (Continued)

PR

- N (randomized/analyzed): 9/9
- Loss to follow-up or excluded: 0
- Age, mean (SD), in years: 70.8 (4.8)
- Gender (M/F): 6/3

Overall

- N (randomized/analyzed): 17/16
- Loss to follow-up or excluded: 1
- Gender (M/F): 10/7
- COPD stage (GOLD): moderate to severe

Included criteria

- FEV1/FVC < 0.65;
- Dyspnea on exertion experienced during activities of daily living;
- A history of cigarette or tobacco smoke exposure > 20 pack-years;
- The ability to self ambulate;
- Age > 60 years;
- Willingness to undergo testing and intervention procedures;
- COPD under appropriate medical management.

Excluded criteria

- Significant cardiac disease;
- Orthopedic or neurologic impairment, serious renal, liver, or gastrointestinal disorders; current psychiatric illness or substance dependence; uncontrolled diabetes or hypertension;
- Current or previous (within 6 months) enrollment in a rehabilitation or exercise program;
- SaoO₂ < 90% during exercise at a heart rate > 50% of age-predicted maximum.

Interventions

Intervention characteristics

PR+IMT

- PR: the rehabilitation program involved walking, upper extremity strength training, and progressive IMT. Walking intensity was set at 50%-75% of the participant's heart rate reserve. This prescription was based on the heart rate response from the participant's initial graded exercise test. The duration of walking was increased progressively throughout the intervention to a maximum of 20 min. Participants were taught to monitor their heart rate during walking by palpating their radial artery and to adjust their exercise intensity as needed. Heart rates were taken midway through and at the end of the walking sessions. Upper extremity weight-training consisted of 5 different exercises. Participants performed 2 sets of 12 repetitions for each exercise with the weight being progressively increased as their strength increased. All walking and strength training was performed at the exercise center.
- IMT: was performed twice daily for a 15-min period, 7 d/week, for 12 weeks, using threshold IMT device (Healthscan Products, Cedar Grove, NJ). The initial pressure setting was set at 15% of the participant's P_Imax for 2 weeks. During Weeks 3 and 4, the threshold pressure was increased to 30% of the participant's P_Imax. During weeks 5 and 6, the threshold pressure was increased to 60% of the participant's P_Imax. During Weeks 7 through 12, the threshold pressure was set at 80% of the participants P_Imax.

PR + sham IMT: this group underwent PR as described above, and IMT at 15% of P_Imax for the duration of the 12-week study period.

Outcomes

Dyspnea: Borg

Functional exercise capacity:

- exercise time (treadmill)
- 12MWD

Berry 1996 (Continued)

Respiratory muscle strength: PImax

Laboratory exercise test: VO₂peak (mL/kg/min)

Respiratory muscle endurance: MVV

Respiratory function:

- FEV1 (%pred)
- FEV1 (L)

Identification
Sponsorship source:

Country: USA

Setting: Wake Forest University

Author's name: Michael J. Berry

Institution: Department of Health and Sport Science

Address: P.O. Box 7234, Wake Forest University, Winston-Salem, NC 27109

Notes

All values are adjusted means ± SEM. Values were adjusted using pre-intervention scores as the covariate.

Berton 2015
Study characteristics
Methods

Study design: RCT

Study grouping: parallel-group

Participants
Baseline characteristics

IMT

- N (randomized/analyzed): /7
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 65.3
- Gender (M/F): 3
- BMI, mean (SD), kg/m²: 25.8

Control/sham

- N (randomized/analyzed): /6
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 66.3 (9.2)
- Gender (M/F): 1/5
- BMI, mean (SD), kg/m²: 24.5 (5.0)

Overall

- N (randomized/analyzed): 24/13
- Loss to follow-up or excluded: 9
- Age, mean (SD) in years: 66.8 (9.1)
- Gender (M/F): 4/9
- COPD stage: moderate to very severe

Berton 2015 (Continued)

Included criteria

- Patients with spirometric evidence of significant chronic airflow limitation (FEV1 < 70%pred, FEV1/FVC < 0.7) according to the criteria of Global Strategy for Diagnosis, Management, and Prevention of COPD;
- Patients with a long history of smoking (> 20 pack-years) were invited to participate from the tertiary clinic care center.
- Participants were receiving continuously formoterol/budesonide (12/400 µg) twice a day (dry powder inhaler), short-acting bronchodilators as rescue medications, and had not participated in PR in the previous 24 months.

Excluded criteria

- Exacerbation of COPD in the previous 3 months or during the study;
- Cardiac disease (acute coronary syndrome in the previous 3 months or cardiac ejection fraction < 50%);
- long term oxygen therapy or arterial oxygen saturation < 90% at rest,
- Neuromuscular disease, peripheral arterial disease, cancer;
- Physically unable to move.

Interventions

Intervention characteristics

IMT: the training was unsupervised, 30 min/d, 7 d/week, for 8 weeks. The device used was Power-breathe(Southam, UK) set at 30% of PImax.

Control/sham: this group underwent similar training at no load.

Outcomes

Dyspnea: Borg

Functional exercise capacity: exercise time

HRQoL: SGRQ

- Symptoms
- Activity
- Impact
- Total

Inspiratory muscle strength (PImax)

Identification

Country: Brazil

Author's name: Danilo C. Berton

Institution: Graduation Program in Pulmonology, Federal University of Rio Grande do Sul (UFRGS), Brazil

Email: dberton@hcpa.edu.br

Address: Rua Ramiro Barcelos, 2350, Room 2050. Postal Code: 90035-003, Porto Alegre, RS, Brazil

Clinical trial register: NCT 01945398

Notes

Participants were instructed to maintain diaphragmatic breathing, with a breathing rate of 15-20 breaths/min.

Bustamante 2007

Study characteristics

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

Bustamante 2007 (Continued)

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 12/12 • Loss to follow-up or excluded: 0 • Age, mean (SD), in years: 62 (13.7) • BMI, mean (SD), kg/m²: 26.03 (3.46) <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 10/10 • Loss to follow-up or excluded: 0 • Age, mean (SD), in years: 61.5 (8.6) • BMI, mean (SD), kg/m²: 26.9 (4.41) <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 22/22 • Loss to follow-up excluded: 0 • COPD stage (GOLD): moderate to severe <p>Included criteria</p> <ul style="list-style-type: none"> • Patients with a clinically stable COPD for a month before selection. <p>Excluded criteria</p> <ul style="list-style-type: none"> • Hypoxemia (PaO₂ < 60 mmHg), asthma, coronary heart disease, metabolic diseases, orthopedic diseases, history of recent thoracic or abdominal surgery, treatment with corticosteroid, hormones and chemotherapy.
Interventions	<p>Intervention characteristics</p> <p>IMT: the training was unsupervised, conducted twice a day, 15 min/session, 7 d/week, for 6 weeks. Threshold IMT device was used at the maximum tolerated load.</p> <p>Control/sham: this group underwent a similar training at a load equal to 7cmH₂O.</p>
Outcomes	<p>HRQoL: CRQ</p> <ul style="list-style-type: none"> • Dyspnea • Fatigue • Emotion • Mastery <p>Respiratory muscle strength: P_{lmax}</p> <p>Respiratory muscle endurance time: T_{lim} (Threshold device)</p> <p>Notes: sustained time with a threshold of 66% of P_{lmax}</p>
Identification	<p>Country: Spain</p> <p>Author's name: Víctor Bustamante Madariaga</p> <p>Institution: Servicio de Neumología. Hospital de Basurto. Osakidetza. Vizcaya. España</p>

Bustamante 2007 (Continued)

Email: VICTOR.BUSTAMANTEMADARIAGA@hbas.osakidetza.net

Notes

Charusisin 2018
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 110/89
- Loss to follow-up or excluded: 26
- Age, mean (SD) in years: 66 (8)
- Gender (M/F): 52/58
- BMI, mean (SD), kg/m² : 25 (6)

PR

- N (randomized/analyzed): 109/85
- Loss to follow-up or excluded: 29
- Age, mean (SD) in years: 65 (7)
- Gender (M/F): 43/66
- BMI, mean (SD), kg/m² : 24 (6)

Overall

- N (randomized/analyzed): 219/174
- Loss to follow-up or excluded: 55
- Gender (M/F): 95/124
- COPD stage (GOLD): moderate to very severe

Included criteria

- Clinically stable COPD patients with reduced maximal inspiratory mouth pressure (P_Imax < 60 cm H₂O or < 50% of the predicted normal value) participated in the study between February 2012 and October 2016.

Excluded criteria

- Diagnosed psychiatric or cognitive disorders
- Progressive neurological or neuromuscular disorders
- Severe orthopaedic problems having a major impact on daily activities
- Previous inclusion in a rehabilitation program (< 1 year)

Pretreatment: "One of the centres offering a 36 session program (32% of total inclusions) consistently exceeded between-group differences in the centre offering 20 sessions (36% of total inclusions). In the other centres offering 36 sessions (combined 32% of total inclusions), between-group differences in these outcomes were consistently smaller than in the centre offering a lower training volume (see on-line supplementary table E4)"

Interventions

Intervention characteristics
Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

64

Charususin 2018 (Continued)

PR+IMT

- PR: consisted of cycling, treadmill walking, stair climbing, arm ergometry and resistance training of arm as well as leg muscles training volume, and ranged from 20 sessions (Germany) to 36 sessions (other centers). Training frequency ranged from 3-5 sessions/week. Duration of training sessions was around 60 min. Participants performed endurance training or interval training at moderate to high intensities. During PR, training intensities were progressively increased according to a Borg CR-10 scale ratings of 4–6 on dyspnea sensation.
- IMT: performed daily using the PowerBreathe KHP2 device (POWERbreatheKHP2, HaB International, Southam, UK) according to previously described methods: 2-3 sessions/d, 7 min each, 7 d/week, for 12 weeks. Training intensity in the intervention group was set initially at a load of approximately 50% of participants' maximal inspiratory mouth pressure (P_Imax). This initial load was then continuously and gradually increased to the highest tolerable intensity during each of the supervised sessions.

PR +sham IMT: this group underwent the same training as described above with IMT load at 10% of P_Imax. The load was not modified throughout the intervention period.

Outcomes	<p>Dyspnea: Borg</p> <ul style="list-style-type: none"> • post-6MWD • Incremental cycle ergometer test • Constant cycle ergometer test <p>Functional exercise capacity:</p> <ul style="list-style-type: none"> • 6MWD • Wmax • Exercise time <p>HRQoL: CRQ</p> <ul style="list-style-type: none"> • Dyspnea • Fatigue • Emotion • Mastery • Total <p>Respiratory muscle strength: P_Imax</p> <p>Laboratory exercise test: VO₂peak</p> <p>Respiratory muscle endurance: MVV</p> <p>Respiratory muscle endurance time: T_{lim}</p> <p>Respiratory function:</p> <ul style="list-style-type: none"> • FEV1 (L) • RV
Identification	<p>Sponsorship source: DL and HD are postdoctoral fellows of Research Foundation Flanders. HaB International (Southam, UK) and McRoberts (The Hague, The Netherlands) provided equipment for testing and training in this study on loan. This study was further supported by local funds throughout the participating centers. The following specific funding sources were reported: University Hospital Leuven, Belgium (FWO grant GOA4516N en KU Leuven grant C22/15/035); Ghent University Hospital, Belgium (UZ Gent grant FS/LGZ/994); Institut Universitaire de Cardiologie et de Pneumologie de Québec, Université Laval, Québec, Canada (Ordre professionnel de la physiothérapie du Québec).</p> <p>Country: Belgium, The Netherlands, Germany, Canada</p> <p>Setting: multicenter</p>

Charususin 2018 (Continued)

Author's name: Daniel Langer
Institution: Department of Rehabilitation Sciences, KU Leuven
Email: daniel.langer@kuleuven.be
Address: Leuven 3001, Belgium
Clinical trial register: NCT01397396

Notes Adjusted difference (95% CI) at post-training with its P value were reported.

Chuang 2017

Study characteristics

Methods **Study design:** RCT
Study grouping: parallel-group

Participants **Baseline characteristics**

IMT

- N (randomized/analyzed): 30/27
- Loss to follow-up or excluded: 3
- Age, mean (SD), in years: 66.22 (12.76)
- Gender (M/F): 17/10

Control/sham

- N (randomized/analyzed): 30/28
- Loss to follow-up or excluded: 2
- Age, mean (SD), in years: 66.04 (10.99)
- Gender (M/F): 19/9

Overall

- N (randomized/analyzed): 60/55
- Loss to follow-up or excluded: 5
- Gender (M/F): 36/19
- COPD stage (GOLD): moderate to very severe

Included criteria

- Outpatients with a stable condition
- Showed airflow limitation on pulmonary function test, that is FEV1 < 80% and FEV1/FVC < 70%
- Participants were conscious and able to express themselves orally, were self-walkers and co-operated with the intervention.

Excluded criteria

- Cardiovascular diseases
- Severe unstable diseases, such as pulmonary heart disease or cancer
- Ongoing oxygen treatment
- Muscle power rating < 5
- ADL scale score < 80

Chuang 2017 (Continued)

- Severe mental illness

Interventions	<p>Intervention characteristics</p> <p>IMT: the training was conducted 5 d/week, for 8 weeks. Each session lasted for 21–30 min. It consisted of a cycle of 2 min inspiratory training with a pressure threshold loading device and 1 min of rest, and then repeated 7 cycles. The training load ranged from 15 cmH₂O to 40 cmH₂O. Each participant was followed up once a day by phone from Monday–Friday, and visits were made by research assistants every 2 weeks to increase the pressure threshold loading progressively after assessing the participant’s condition.</p> <p>Control: no intervention was received by this group.</p>
Outcomes	<p>Dyspnea: BDI</p> <p>Functional exercise capacity: 6MWD</p> <p>HRQoL: SF-36 Questionnaire</p> <ul style="list-style-type: none"> • Physiological functions • Mental functions <p>Respiratory muscle strength: PImax</p>
Identification	<p>Country: Taiwan</p> <p>Author's name: Hsiao-Yun Chang</p> <p>Institution: Department of Nursing, Fooyin University, Kaohsiung City</p> <p>Email: chang369@gmail.com</p> <p>Address: Taiwan</p>
Notes	

Covey 2001
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 19/12 • Loss to follow-up or excluded: 10 • Age mean (SD) in years: 65 (6) • BMI, mean (SD), kg/m²: 26 (4.8) <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 18/15 • Loss to follow-up or excluded: 3 • Age mean (SD) in years: 67 (10) • BMI, mean (SD), kg/m²: 28.8 (6) <p>Overall</p>

Covey 2001 (Continued)

- N (randomized/analyzed): 37/27
- Loss to follow-up or excluded: 10
- Gender (M/F): 18/9
- COPD stage: severe to very severe

Included criteria

- Severe to very severe airflow obstruction (FEV1 < 50%pred)
- Age: 45-75
- Severely limited functional performance

Excluded criteria

- History of asthma or > 25% increase in FEV1 after bronchodilator
- History of a major exacerbation within the past 2 months
- Current oral corticosteroid use (> 10 mg prednisone per day)
- Other health problems that would inhibit their ability to participate

Interventions	<p>Intervention characteristics</p> <p>IMT: participants performed IMT at home 5 d/week, 30 min/d for 16 weeks (for a total of 80 training sessions) using Threshold IMT device. Starting training loads were 30% of P_{lmax} as tolerated. Participants were visited weekly at home by a nurse who supervised training, evaluated training loads, and progressively increased the training load as tolerated with a goal of achieving 60% of P_{lmax}. Home visits generally lasted approximately 30 min. An interval training protocol was used with participants performing 6 work sets of 5 mins' duration separated by rest intervals lasting 1-3 min.</p> <p>Control/sham: participants were visited by a nurse every 2 weeks for 16 weeks for a structured program of health education. Each home-based session lasted approximately 1-1.5 h and covered such topics as nutrition, relaxation techniques, pursed lip breathing, respiratory medications, respiratory infection, energy conservation techniques, oxygen therapy, and smoking cessation.</p>
Outcomes	<p>Dyspnea: Borg</p> <p>HRQoL: CRQ</p> <p>Respiratory muscle strength (P_{lmax})</p> <p>Respiratory muscle endurance pressure: P_{thmax}</p>
Identification	<p>Sponsorship: This study was supported by a grant from the National Institutes of Nursing Research, grant number NRO1428; and was conducted at the University of Illinois at Chicago, College of Nursing, Chicago, Ill and Hines VA Hospital, Section of Critical Care and Pulmonary Medicine, Hines, Ill.</p> <p>Country: USA</p> <p>Setting: home-based</p> <p>Author's name: Margaret K. Covey</p> <p>Institution: University of Illinois at Chicago, College of Nursing</p> <p>Email: mkcovey@uic.edu</p> <p>Address: 845 South Damen Avenue, Chicago, IL 60612</p>
Notes	

Cutrim 2019
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 11/11 • Loss to follow-up or excluded: 0 • Age, mean (SD), in years: 66 (8.5) • Gender (M/F): 8/3 • BMI, mean (SD), kg/m² : 25 (4) <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 11/11 • Loss to follow-up or excluded: 0 • Age, mean (SD), in years: 70 (8.0) • Gender (M/F): 9/2 • BMI, mean (SD), kg/m² : 25 (4) <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 22/22 • Gender (M/F): 17/5 <p>Included criteria</p> <ul style="list-style-type: none"> • Patients with COPD without heart failure or pulmonary hypertension • With inspiratory muscle weakness (P_Imax < 70% of predicted) and with stable pharmacological treatment, i.e. no drug change at least 1 month before the start of IMT <p>Excluded criteria</p> <ul style="list-style-type: none"> • Functional limiting factors that would interfere with the performance of IMT and/or the exercise capacity test, such as acute myocardial infarction 3 months before inclusion in the study, unstable angina or unstable ventricular arrhythmia or in the last 3 months prior to initiation, acute respiratory disease, rheumatic diseases, degenerative diseases, neurological sequelae, cognitive deficit etc
Interventions	<p>Intervention characteristics</p> <p>IMT: the session of IMT consisted of 30 min (in a clinical setting) 3 times/week, using the Threshold Inspiratory Muscle Training device (POWERbreathe Medic+Plus, NCS, Barueri, SP, Brazil). The inspiratory load was set at 30% of P_Imax, for 12 weeks. During exercise, participants were instructed to maintain diaphragmatic breathing at a rate of 15–20 breaths/min.</p> <p>Control/sham: no intervention received by this group (except diaphragmatic breathing).</p>
Outcomes	<p>Functional exercise capacity: 6MWD</p> <p>Respiratory muscle strength: P_Imax</p>
Identification	<p>Sponsorship source: Fundação de Amparo à Pesquisa do Estado do Maranhão (FAPEMA); Hospital Universitário Presidente Dutra; Cristiano Mostarda received grants from CNPq (Universal 442374/2014-3) and FAPEMA (Bolsa Produtividade and Universal 00358/15).</p> <p>Country: Brazil</p>

Cutrim 2019 (Continued)

Setting: Hospital Universitário Presidente Dutra

Author's name: Cristiano Teixeira Mostardaa

Institution: Universidade Federal do Maranhão, São Luís, Brazil

Email: cristiano.mostarda@gmail.com

Address: Av. dos Portugueses, 1966, Cidade Universitária Dom Delgado, São Luís, MA, Brazil

Clinical trial register: RBR-4mz6w9

Notes

Participants were instructed to maintain diaphragmatic breathing at a rate of 15–20 breaths/min. Adjusted analysis for age, weight and baseline were reported.

Dacha 2019
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): not reported/6
- N (analyzed): 6
- Loss to follow-up or excluded: 0
- Age, mean (SD), in years: 65 (4)
- Gender (M/F): 3/3
- BMI, mean (SD), kg/m²: 29 (8)
- COPD stage (GOLD): mild to very severe

Control/sham

- N (randomized/analyzed): not reported/4
- N (analyzed): 4
- Loss to follow-up excluded: 0
- Age, mean (SD), in years: 68 (9)
- Gender (M/F): 1/3
- BMI, mean (SD), kg/m²: 22 (2)
- COPD stage (GOLD): moderate to very severe

Overall

- N (randomized/analyzed): 24/10
- N (analyzed): 10
- Loss to follow-up excluded: 0
- Gender (M/F): 4/6
- COPD stage (GOLD): mild to very severe

Included criteria

- Clinical Diagnosis of COPD
- Inspiratory muscle weakness (P_{Imax} < 70%pred or < 60 cmH₂O)

Dacha 2019 (Continued)

- BDI < 7
- Peripheral muscle fatigue present after CPET

Excluded criteria

- Major cardiovascular limiting exercise capacity more than pulmonary function impairment
- Severe orthopedic with a major impact on ADL
- Psychiatric or cognitive disorders
- Progressive neurological or neuromuscular disorders
- Long term O2 therapy
- Previous inclusion in a rehabilitation program (< 1 year)

Interventions

Intervention characteristics

IMT: the training was conducted twice a day, 7 d/week, for 8 weeks. All the sessions were supervised, 5 min each, using Powerbreathe device set at a 50% of PImax.

Control/sham: participants in this group received a similar protocol with a training load at 10% of PImax.

Outcomes

Dyspnea: Borg

Respiratory muscle strength: PImax

Laboratory exercise test: VO2peak

Identification

Sponsorship source: KU Leuven

Country: Belgium

Author's name: Sauwaluk Dacha

Institution: Faculty of Kinesiology and Rehabilitation Sciences, Department of Rehabilitation Sciences, Research Group for Rehabilitation in Internal Disorders

Email: sauwaluk.dacha@kuleuven.be

Address: KU Leuven, Leuven, Belgium

Clinical trial register: NCT03240640

Notes

This study is still ongoing.

De Farias 2019

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Overall

- N (randomized/analyzed): 33/31
- Loss to follow-up or excluded: 2
- Age, mean (SD) in years: 66.2 (4.9)
- BMI: 28 (4.3)

De Farias 2019 (Continued)

Inclusion criteria

- Individuals with a clinical diagnosis of COPD according to GOLD
- Being treated at the Ambulatory Pulmonology department of University Hospital Onofre Lopes (HUOL)/Empresa Brasileira de Serviços Hospitalares (EBSERH)
- Aged 40-80 years
- Living in the city of Natal, RN/Brazil
- Not using oxygen therapy or presenting disease exacerbation in the last 3 months
- Not practising regular physical activity in the last 6 months

Exclusion criteria

- Musculoskeletal comorbidities that impair gait
- SpO₂ < 90% during 6MWD
- Hypertensive without control medication as well as those presenting with a hypertensive peak (> 140/90 mmHg) for > 3 consecutive days
- An intellectual understanding impairment that interferes with the evaluation tests
- Those who stop the therapeutic program, miss activity for > 1 week, or miss reevaluation

Interventions

Intervention characteristics

PR+ sham IMT

- PR: performed 3 times/week for 10 weeks. All individuals were instructed to perform the rehabilitation program 5 days/week — 3 days with supervision and 2 without supervision — for at least 1 h/d. The program consisted of health education, treadmill aerobic training with 70% of max incremental shuttle walk test speed and peripheral muscle strength training.
- Sham IMT: performed with Powerbreathe device at no load for 3 cycles of 12 repetitions.

PR+ threshold IMT

- PR: the same protocol as described above
- Threshold IMT: participants trained using POWERbreathe KH1 (POWERbreathe International Ltd.) at 35% of MIP), increasing 5% every week until reaching 80% of MIP at the 10th week, which was maintained until the end of the protocol. The MIP was assessed weekly to adjust the training load percentage. Participants performed 3 cycles of 12 repetitions.

PR + isocapnic hyperpnea

- PR: the same protocol as described above.
- Isocapnic hyperpnea: participants received training with a duration of up to 20 min, 1 min of training and 1 min of rest, where they were encouraged with instructions such as “breathe faster.” The training respiratory rate chosen was calculated as 35-fold FEV₁, so that ventilation corresponded to 50%–60% of MVV. The device used was STMedical device (SpiroTiger, Chamonix Mont Blanc, France).

Outcomes

Dyspnea: mMRC
Functional exercise capacity: 6MWD
Respiratory muscle strength: PImax

Identification

Sponsorship source: no funding

Country: Brazil

Author's name: Guilherme Augusto de Freitas

Institution: PneumoCardioVascular Lab/HUOL, Empresa Brasileira de Serviços Hospitalares - EBSERH), Universidade Federal do Rio Grande do Norte (UFRN),

Email: fregonezi.guilherme@gmail.com

De Farias 2019 (Continued)

Address: Natal, Rio Grande do Norte, Brazil

Notes

Dekhuijzen 1991
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0
- Age, mean (SD), in years: 58/8
- Gender (M/F): 14/6

PR

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0
- Age, mean (SD), in years: 60/7
- Gender (M/F): 16/4

Overall

- N (randomized/analyzed): 40/40
- Loss to follow-up or excluded: 0
- Gender (M/F): 30/10
- COPD stage (GOLD): moderate to severe

Included criteria: patients with functional limitations due to COPD

Excluded criteria: not reported

Interventions

Intervention characteristics
PR+IMT

- PR: consisted of exercise training, i.e. cycling, walking, and training of back, shoulder, and abdominal muscles. The intensity of the exercise training was determined by the symptoms of the patients. Moreover, the heart rate during these exercises did not exceed 80% of the maximal heart rate reached during the maximal bicycle ergometer test. Other parts of the PR program were callisthenics, conventional physiotherapy (breathing retraining, relaxation exercises), and education about pulmonary disease and the purpose and use of the medications. The training was conducted 2 h every day, 5 d/week, for 10 weeks.
- IMT: was conducted along with PR, 15 min twice a day, supervised by the physiotherapist, using incentive spirometer at 70% of P_Imax.

PR: this group received only the PR described above.

Outcomes

HRQoL: ADL

Functional exercise capacity:

Dekhuijzen 1991 (Continued)

- 12MWD
- Wmax (incremental cycle ergometer test)

Respiratory muscle strength: P_{lmax}

Laboratory exercise test: VO_{2peak}

Respiratory muscle endurance time: T_{lim}

Identification	<p>Country: The Netherlands</p> <p>Setting: Medical Centre Dekkerswald (outpatient clinic)</p> <p>Author's name: P. N. Richard Dekhuijzen</p> <p>Institution: the University of Nijmegen, Department of Pulmonary Diseases, Medical Centre Dekkerswald</p> <p>Address: Groesheek, the Netherlands</p>
Notes	We used convertunits.com to convert from Kpa to cmH2O.

Dellweg 2017
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 15/15 • Loss to follow-up or excluded: 0 • Age, mean (SD), in years: 66 (8) • Gender (M/F): 7/8 • BMI, mean (SD), kg/m²: 24.5 (6.4) <p>PR</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 14/14 • Loss to follow-up or excluded: 0 • Age, mean (SD), in years: 66 (7.5) • Gender (M/F): 9/5 • BMI, mean (SD), kg/m²: 23.9 (4) <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 29/29 • Loss to follow-up or excluded: 0 • Gender (M/F): 16/13 • COPD stage (GOLD): severe to very severe <p>Included criteria</p> <ul style="list-style-type: none"> • Patients with persistent hypercapnic respiratory failure who required non-invasive ventilation after prolonged weaning according to the criteria of Boles et al (Boles 2007)

Dellweg 2017 (Continued)

- Main diagnosis of COPD
- Participants had to be admitted to our in-patient post-weaning rehabilitation unit, had to be ambulatory, co-operative, and physically able to participate in twice-daily physiotherapeutic sessions
- Patients had to have confirmed COPD stage 3 or 4 by a lung function test (Fev1 < 50%, Fev1/FVC < 70%)
- Had to be free of exacerbation

Excluded criteria

- Renal impairment (serum creatinine levels < 2 mg/dL)
- Severe cardiac impairment (ejection fraction < 40%)

Interventions	Intervention characteristics PR+IMT: <ul style="list-style-type: none"> • PR: consisted of twice daily, hour-long physiotherapy group sessions in the rehabilitation gym. Physiotherapy sessions included training with arm and leg ergometers, as well as weight training. Additionally, every patient participated in daily ergotherapy group sessions lasting 1 h each to improve fine motor skills. • IMT: consisted of supervised sessions, once daily during weekdays, for 4 weeks. Participants underwent strength training at 80% of P_{lmax} and endurance training at 60% of P_{lmax} using Respifit S Trainer (Biegler, Mauerbach, Austria) PR+ (sham IMT): participants in this group received a similar protocol as described above with IMT training load fixed at 5 cm H ₂ O, and using Threshold IMT (Philips-Respironics, Pittsburgh, PA, USA).
Outcomes	Functional exercise capacity: 6MWD Respiratory muscle strength: P _{lmax} (RV) Respiratory function: FEV1
Identification	Sponsorship source: no funding Country: Germany Author's name: Dominic Dellweg Institution: Department for Pulmonology, Intensive Care and Rehabilitation Email: d.dellweg@fkkg.de Address: 1, 57392 Schmallenberg, Germany Clinical trial register: NCT00291460
Notes	

Fanfa Bordin 2020
Study characteristics

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics PR+IMT

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

75

Fanfa Bordin 2020 (Continued)

- N (randomized or analyzed): 10/10
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 63.2 (5.7)
- Gender (M/F): 5/5
- BMI, mean (SD), kg/m² : 25.3 (5.1)

PR

- N (randomized/analyzed): 12/10
- Loss to follow-up or excluded: 2
- Age, mean (SD) in years: 66.2 (9.2)
- Gender (M/F): 4/6
- BMI, mean (SD), kg/m² : 25.4 (5.8)

Overall

- N (randomized/analyzed): 22/20
- Loss to follow-up or excluded: 2
- Gender (M/F): 9/11
- COPD stage (GOLD): moderate to very severe

Included criteria

- COPD patients at the stages 2-4, according to the GOLD classification
- Participating in a lung rehabilitation program for at least 2 months
- Clinically stable
- Signed the informed consent

Excluded criteria

- Individuals with asthma, and/or cardiovascular disease and individuals with cognitive and/or behavioral impairments

Interventions	<p>Intervention characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • PR: the lung rehabilitation program included 30 min of cycling exercise on a vertical cycle ergometer for lower limbs (Movement, BM 2700, Brazil) set at 60% of the maximal heart rate determined using the Karvonen method modified for the reserve heart rate (Meyer 2013). Participants also performed strengthening exercises for the upper and lower limbs' major muscles, with the intensity of 50%-80% of the 1 repetition maximum test (1RM). • IMT: involved supervised sessions, 3 times/week for 20 min over 4 weeks, which was increased to 25 min on weeks 5 and 6, and to 30 min on weeks 7 and 8. Participants used Threshold IMT set at 50% of P_Imax. <p>PR: this group received only the PR described above.</p>
Outcomes	Respiratory muscle strength: P _I max
Identification	<p>Country: Brazil</p> <p>Setting: Santa Cruz do Sul Hospital, Brazil</p> <p>Author's name: Diogo Fanfa Bordin</p> <p>Institution: Universidade de Santa Cruz do Sul (Unisc) – Santa Cruz do Sul (RS), Brazil</p> <p>Email: diogo.fanfa@hotmail.com</p>

Fanfa Bordin 2020 (Continued)

Clinical trial register: NCT02014155

Notes

Harver 1989
Study characteristics

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics IMT <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/10 • Loss to follow-up or excluded: not reported • Age, mean (SD) in years: 61.1 (9.5) • Gender (M/F): 8/2 Control/sham <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/9 • Loss to follow-up or excluded: not reported • Age, mean (SD) in years: 64.8 (8.4) • Gender (M/F): 8/1 Overall <ul style="list-style-type: none"> • N (randomized/analyzed): 23/19 • Loss to follow-up or excluded: 4 • Gender (M/F): 16/3 • COPD stage (GOLD): moderate to severe Included criteria <ul style="list-style-type: none"> • Participants for the study were recruited from the clinical practice of pulmonary physicians at the institution • Participants were stable both clinically and functionally, and medications were not changed during the study. Excluded criteria: not reported
Interventions	Intervention characteristics IMT: participants trained twice a day, 15 min each session, 7 d/week, for 8 weeks. The training device was Pflex (with a controlled breathing pattern), and the training load ranged from 5 cmH ₂ O to 35 cmH ₂ O (≈30% P _I max FRC) Control/sham: this group received a similar IMT protocol with a training load set at 5cmH ₂ O
Outcomes	Dyspnea: BDI-TDI: <ul style="list-style-type: none"> • Functional impairment • Magnitude of task • Magnitude of effort

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

77

Harver 1989 (Continued)

- Focal score
- Respiratory muscle strength: PImax
- RV
 - FRC
- Respiratory function: FEV1
- L
- Respiratory muscle endurance: MVV

Identification

Sponsorship source: The American Lung Association of New Hampshire and by grants HL07449 and HL29068 from the National Heart, Lung, and Blood Institute

Country: USA

Setting: Outpatient pulmonary clinic and pulmonary function laboratory

Author's name: Andrew Harver

Institution: Department of Psychology, SUNY Stony Brook

Address: Stony Brook, NY 11794

Notes

Heijdra 1996

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): 10/10
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 62.4 (8.8)
- Gender (M/F): 7/3
- BMI, mean (SD), kg/m²: 23.7 (3.3)

Control/sham

- N (randomized/analyzed): 10/10
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 61.8 (7.3)
- Gender (M/F): 8/2
- BMI, mean (SD), kg/m²: 23 (3.2)

Overall

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0
- Gender (M/F): 15/5

Heijdra 1996 (Continued)

- COPD stage (GOLD): moderate to very severe

Included criteria

- Mean nocturnal arterial oxygen saturation was < 92%
- All participants were in a stable condition at the time of study as defined by a fluctuation in FEV1 of < 10% in the preceding 6 months

Excluded criteria

- Patients with other pulmonary diseases, chest wall deformations, a previous thoracotomy, diabetes mellitus, neuromuscular diseases, obstructive sleep apnea syndrome, or an overlap syndrome

Interventions

Intervention characteristics

IMT: participants trained at home, daily, 2 sessions of 15 min/day for 10 weeks. They used incentive flowmeter (INSPIRx; Resprecare Medical Inc., the Hague, the Netherlands) set at 60% of PImax

Control/sham: participants received a similar training protocol and the resistance was set at 10% of PImax

Outcomes

Inspiratory muscle strength (PImax)

Respiratory muscle endurance time (T_{lim})

Identification

Sponsorship source: Dutch Asthma Foundation (No. 90-27)

Country: The Netherlands

Author's name: Yvonne F. Heijdra, Department of Pulmonary Diseases

Institution: University of Nijmegen, Medical Centre Dekkerswald

Address: P.O. Box 9001, 6560 GB Groesbeek, the Netherlands

Notes

Hill 2006

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): 18/16
- Loss to follow-up or excluded: 2
- Age, mean (SD) in years: 69.4 (7.2)
- Gender (M/F): 11/5
- BMI, mean (SD), kg/m²: 24.9 (4.3)

Control/sham

- N (randomized/analyzed): 17/17
- Loss to follow-up or excluded: 0

Hill 2006 (Continued)

- Age, mean (SD) in years: 66.6 (9.8)
- Gender (M/F): 11/6
- BMI, mean (SD), kg/m²: 24.1 (3.7)

Overall

- N (randomized/analyzed): 35/33
- Loss to follow-up or excluded: 2
- Gender (M/F): 22/11
- COPD stage (GOLD): severe to very severe

Included criteria

- Participants who had a diagnosis of COPD
- Smoking history of 10 pack-years
- FEV1 ranging 15%–70% of predicted normal

Excluded criteria

- Comorbid conditions likely to reduce exercise capacity (e.g. symptomatic ischemic heart disease, BMI > 35 kg.m²)
- Previous lung surgery
- The use of long-term oxygen therapy
- Weaning doses of oral corticosteroids

Interventions	Intervention characteristics
	<p>IMT: participants attended supervised training sessions 3 times/week for 8 weeks. Each session lasted 21 min and comprised 7 cycles of 2 min of breathing on an inspiratory threshold loading device (Threshold IMT; Respiroics, Cedar Grove, NJ, USA) followed by 1 minute of rest. The training load was set at a range of around 45%-101% of P_Imax</p> <p>Control/sham: this group received a similar IMT protocol with a training load set at 10% of P_Imax</p>
Outcomes	<p>Dyspnea: Borg</p> <ul style="list-style-type: none"> • Post 6MWD • Wmax <p>Functional exercise capacity:</p> <ul style="list-style-type: none"> • 6MWD • Wmax <p>HRQoL: CRQ</p> <ul style="list-style-type: none"> • Dyspnea • Fatigue • Emotion • Control • Total <p>Respiratory muscle strength (P_Imax)</p> <p>Respiratory muscle endurance: Respiratory muscle endurance pressure (P_{th}max)</p> <p>Laboratory exercise test: VO₂peak (mL/kg/min)</p> <p>Respiratory function: FEV1</p> <ul style="list-style-type: none"> • %pred

Hill 2006 (Continued)

- L
- Respiratory function: RV
- %pred
 - L

Identification

Sponsorship source: The National Health and Medical Research Council (Canberra, Australia) grant number 212016

Country: Australia

Author's name: P.R. Eastwood

Institution: Dept of Pulmonary Physiology, Sir Charles Gairdner Hospital

Email: peter.eastwood@health.wa.gov.au

Address: Hospital Avenue Nedlands Western Australia, Australia 6009

Notes
Hill 2007
Study characteristics
Methods

Study design: RCT

Study grouping: parallel-group

Participants
Baseline characteristics
IMT

- N (randomized/analyzed): 18/16
- Loss to follow-up or excluded: 2
- Age, mean (SD) in years: 69.4 (7.2)
- Gender (M/F): 11/5
- BMI, mean (SD), kg/m²: 24.9 (4.3)

Control/sham

- N (randomized/analyzed): 17/17
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 66.6 (9.8)
- Gender (M/F): 11/6
- BMI, mean (SD), kg/m²: 24.1 (3.7)

Overall

- N (randomized/analyzed): 35/33
- Loss to follow-up or excluded: 2
- Gender (M/F): 22/11
- COPD stage (GOLD): severe to very severe

Included criteria

- Participants who had a diagnosis of COPD
- Smoking history of 10 pack-years

Hill 2007 (Continued)

- FEV1 ranging 15%–70% of predicted normal

Excluded criteria

- Comorbid conditions likely to reduce exercise capacity (e.g. symptomatic ischemic heart disease, BMI > 35 kg.m²)
- Previous lung surgery
- The use of long-term oxygen therapy
- Weaning doses of oral corticosteroids

Interventions	<p>Intervention characteristics</p> <p>IMT: participants attended supervised training sessions 3 times/week for 8 weeks. Each session lasted 21 min and comprised 7 cycles of 2 min of breathing on an inspiratory threshold loading device (Threshold IMT; Respironics, Cedar Grove, NJ, USA) followed by 1 minute of rest. The training load was set at a range of around 45%-101% of P_Imax</p> <p>Control/sham: this group received a similar IMT protocol with a training load set at 10% of P_Imax</p>
Outcomes	<p>Respiratory muscle endurance time: T_{lim}</p> <p>Respiratory muscle endurance pressure (P_{thmax})</p> <ul style="list-style-type: none"> • Note: the same as Hill 2006
Identification	<p>Sponsorship source: The National Health and Medical Research Council (Canberra, Australia) grant number 212016</p> <p>Country: Australia</p> <p>Author's name: P.R. Eastwood</p> <p>Institution: Dept of Pulmonary Physiology, Sir Charles Gairdner Hospital</p> <p>Email: peter.eastwood@health.wa.gov.au</p> <p>Address: Hospital Avenue Nedlands Western Australia, Australia 6009</p>
Notes	

Hsiao 2003
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT (Threshold device)</p> <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/10 • Loss to follow-up or excluded: not reported • Age, mean (SD) in years: 68.2 (6.5) • Gender (M/F): 10/0 <p>IMT (Incentive spirometer)</p>

Hsiao 2003 (Continued)

- N (randomized/analyzed): not reported/10
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 70.4 (5.3)
- Gender (M/F): 8/2

Control/sham

- N (randomized/analyzed): not reported/10
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 71.1 (3.9)
- Gender (M/F): 8/2

Overall

- N (randomized/analyzed): 42/30
- Loss to follow-up or excluded: 12
- Gender (M/F): 26/4
- COPD stage (GOLD): moderate to severe

Included criteria

- Patient with moderate to severe COPD
- FEV1 < 2L and FEV1/FVC < 60%

Excluded criteria

- Restrictive lung disease
- History of cardiovascular disease or musculoskeletal conditions that could interfere with the training or the testing maneuvers

Interventions

Intervention characteristics

IMT: participants trained twice a day, 15 min/session, 5days/week for 8 weeks. The sessions were unsupervised. Participants were divided into two groups using either Threshold IMT or Respirex, which were set at 50% of P_Imax

Control/sham: this group did not receive any intervention

Outcomes

Functional exercise capacity: 6MWD

Respiratory muscle strength: P_Imax (RV)

Respiratory muscle endurance time: T_{lim} (Threshold device)

- Notes: measured while breathing against 70% of P_Imax

Identification

Country: Taiwan

Setting: Pulmonary clinic of 1 university hospital

Author's name: Ying Tai Wu

Institution: School and graduate institute of physical therapy, college of medicine, National Taiwan University Hospital

Address: 7 Chung Shan South Road, Taipei 100, Taiwan

Notes

Kim 1993

Study characteristics

Methods **Study design:** RCT
Study grouping: parallel-group

Participants **Baseline characteristics**

IMT

- N (randomized/analyzed): not reported/41
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 66 (7)
- BMI, mean (SD), kg/m²: 25 (4)

Control/sham

- N (randomized/analyzed): not reported/26
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 63 (8)
- BMI, mean (SD), kg/m²: 25 (4)

Overall

- N (randomized/analyzed): 129/67
- Loss to follow-up or excluded: 62
- Gender (M/F): 51/16
- BMI, mean (SD), kg/m²:
- COPD stage (GOLD): severe to very severe

Included criteria

- No history of asthma or restrictive lung disease
- Free of respiratory tract infection for at least 2 months prior to enrollment and free of other chronic health problems that would interfere with their ability to participate in the study

Excluded criteria

- > 10 mg of prednisone daily

Interventions **Intervention characteristics**

IMT: participants trained 7 days/week, 30 min/session, for 6 months. Training sessions were unsupervised, and the device used was a Threshold IMT device set at 30% of P_Imax

Control/sham: this group received IMT at no load

Outcomes

Functional exercise capacity: 12MWD

Respiratory muscle endurance time: T_{lim}

- Notes: measured while breathing at 66% of P_Imax.

Respiratory muscle strength: P_Imax (RV)

Identification **Sponsorship source:** supported by a grant from the National Center for Nursing Research, grant number NRO1428

Country: USA

Author's name: Mija Kim

Kim 1993 (Continued)

Institution: College of Nursing, University of Illinois at Chicago

Notes

Koppers 2006
Study characteristics

Methods	Study design: RCT Study grouping: parallel-group Subgroup analysis: P _{lmax} < 75%pred
---------	--

Participants

IMT

- N (randomized/analyzed): 19/18
- Loss to follow-up or excluded: 1
- Age, mean (SD) in years: 54.4 (7.7)
- Gender (M/F): 8/10
- BMI, mean (SD), kg/m²: 26.7 (5.0)

Control/sham

- N (randomized/analyzed): 20/18
- Loss to follow-up or excluded: 2
- Age, mean (SD) in years: 57.0 (8.5)
- Gender (M/F): 9/9
- BMI, mean (SD), kg/m²: 27.5 (3.3)

Overall

- N (randomized/analyzed): 39/36
- Loss to follow-up or excluded: 3
- Range age (min, max): 38,73
- Gender (M/F): 17/19
- COPD stage (GOLD): moderate to severe

Included criteria

- Chronic airflow obstruction, defined as an FEV₁/FVC ratio 70%, FEV₁ of 30%-80%pred, after bronchodilation
- Stable clinical condition for at least 6 weeks

Excluded criteria

- Hypoxemia at rest or during exercise
- Cardiac orthopaedic disease
- BMI > 30 kg/m²

Interventions

Intervention characteristics

IMT: respiratory muscle endurance training was performed by means of tube breathing. A tube (internal diameter, 3 cm) connected to a mouthpiece was added to the respiratory system to rebreathe exhaled carbon dioxide (normocapnic hyperpnea). The maximum ventilatory capacity that can be sustained for 15 min is approximately 60% of MVV. Therefore, the aimed level of ventilation during training

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

85

Koppers 2006 (Continued)

was set at 60% of MVV, which was calculated from 35 times FEV1 ($60\% \text{ MVV} = 0.6 * 35 * \text{FEV1}$). The dead space was adjusted to 60% of the participant's IVC plus the resting tidal volume. Participants trained twice daily, for 15 min, 7 d/week for 5 weeks.

Control/sham: participant breathed 6-7 times/min through an incentive flowmeter (Inspirix; Respire-care Medical; the Hague, the Netherlands). Airflow resistance was set at 5% of P_{lmax}.

Outcomes	<p>Dyspnea: Borg</p> <ul style="list-style-type: none"> Notes: Borg was measured at isocapnic time during constant-load exercise training on a cycle ergometer <p>Functional exercise capacity:</p> <ul style="list-style-type: none"> 6MWD W_{max} Exercise time (Constant cycle ergometer test) <p>HRQoL: CRQ (Total)</p> <p>Respiratory muscle strength: P_{lmax} (RV)</p> <p>Laboratory exercise test: VO_{2peak} (mL/kg/min)</p> <p>Respiratory muscle endurance: Respiratory muscle endurance pressure (P_{thmax})</p> <p>Respiratory function: FEV1 (L)</p>
Identification	<p>Sponsorship source:</p> <p>Country: The Netherlands</p> <p>Setting: Department of Pulmonology Dekkerswald, University Medical Center Nijmegen</p> <p>Author's name: Ralph J. H. Koppers</p> <p>Institution: MedicalCenter Leeuwarden, Leeuwarden; and Department of Pulmonology (Drs. Vos, Boot, and Folgering), Dekkerswald, University Medical Center Nijmegen</p> <p>Email: R.J.H.Koppers@ZNB.nl</p> <p>Address: Nijmegen, the Netherlands</p>
Notes	

Langer 2018

Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> N (randomized/analyzed): 10/10 Loss to follow-up or excluded: 0 Age, mean (SD) in years: 73 (4)

Langer 2018 (Continued)

- Gender (M/F): 4/6
- BMI, mean (SD), kg/m²: 24.1 (4.6)

Control/sham

- N (randomized/analyzed): 10/10
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 67 (8)
- Gender (M/F): 3/7
- BMI, mean (SD), kg/m²: 25.1 (6.7)

Overall

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 70 (7)
- Gender (M/F): 7/13
- BMI, mean (SD), kg/m²: 24.6 (5.6)
- COPD stage (GOLD): moderate to very severe

Included criteria

- Participants were clinically stable COPD patients with reduced inspiratory muscle strength (P_{imax} < 70 cmH₂O measured at plethysmographic FRC)
- Persistent activity-related dyspnea (BDI < 9) despite optimal medical therapy

Excluded criteria

- Inability to perform physiological testing
- Active cardiovascular comorbidity (i.e. severe heart failure with reduced left ventricular ejection fraction, cardiomyopathy, recent acute myocardial infarction, cardiac arrhythmias, or stroke), or other conditions that could impact dyspnea or exercise capacity

Interventions

Intervention characteristics

IMT: the training was performed with a home-based protocol using an electronic device: Powerbreathe KH2 (HAB International, Southam, UK). Participants trained 2-3 daily sessions of 30 breaths (4-5 min/session) performed 7 days/week for 8 weeks. The training load started at around 40% of P_{imax} and it was increased weekly until the highest tolerable intensity

Control/sham: this group performed IMT at a load of < 10% of P_{imax}

Outcomes

Dyspnea: Borg

- isotime cycle ergometer test)
- peak exercise cycle ergometer test)
- Notes: Borg measured at isotime: cycle ergometer test

Dyspnea: BDI-TDI: Total

Dyspnea: mMRC

Functional exercise capacity: Exercise time (constant cycle ergometer test)

Respiratory muscle strength: P_{imax}

- RV
- FRC

Respiratory muscle endurance time (T_{lim})

Langer 2018 (Continued)

- Notes: measured through breathing against 50%-60% of PImax

Laboratory exercise test: VO₂peak (Constant cycle ergometer test) (L/min)

Respiratory function: FEV₁ (L)

Identification

Sponsorship source: this work was supported by the Ontario Thoracic Society, Spear/StartEndowment Fund, Queen's University. D. Langer received a postdoctoral fellowship and travel grant from the Research Foundation Flanders; thePowerBreathe devices used in the study were provided by HaB International

Country: Canada

Setting: Queen's University Health Sciences and Affiliated Teaching Hospitals

Author's name: D. E. O'Donnell

Institution: Respiratory Investigation Unit, Queen's University and Kingston Health Sciences Centre

Email: odonnell@queensu.ca

Address: 102 Stuart St., Kingston, ON, Canada, K7L 2V6

Clinical trial register: NCT01900873

Notes

webplotdigitizer used to extract IMT training load intensity. Adjusted analysis were provided.

Larson 1988

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): not reported/10
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 60 (6)

Control/sham

- N (randomized/analyzed): not reported/12
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 68 (3)

Overall

- N (randomized/analyzed): 45/22
- Loss to follow-up or excluded: 23
- Gender (M/F): 20/2
- COPD stage (GOLD): severe to very severe

Included criteria

- FEV₁ < 65%pred

Larson 1988 (Continued)

- Stable condition
- Patients with a recent history of exacerbation were allowed to enter the study only after they reported that their ability to function and day-to-day symptoms had returned to baseline
- All patients were enrolled > 2 months after recovery from an exacerbation except 1 who enrolled 1 month after recovery from an exacerbation because he insisted that his breathing would not improve any further
- Patients were advised against participation if they indicated that the 12MWD was too strenuous.

Excluded criteria

- Evidence of restrictive lung disease based on lung volumes
- A history of asthma, and if they had a history of cardiovascular disease or musculoskeletal conditions that could interfere with either the training or testing maneuvers
- Taking psychotropic drugs or abusing alcohol
- Participants were dropped from the study if they: experienced an exacerbation, required a change in their pharmacologic regimen, or reported < 80% compliance with the training protocol.

Interventions	<p>Intervention characteristics</p> <p>IMT: participants trained 7 d/week for 8 weeks using the Threshold IMT device at 30% of P_Imax. They initiated the training for 15 min/d during the first week and gradually increased the duration to 30 min/day for the remaining 7 weeks.</p> <p>Control/sham: this group received the same IMT protocol with the training load set at 15% of P_Imax</p>
Outcomes	<p>Respiratory muscle strength: P_Imax (RV)</p> <p>Functional exercise capacity: 12MWD</p> <p>Respiratory muscle endurance time: T_{lim}</p> <ul style="list-style-type: none"> • Note: measured through breathing against 66% of P_Imax
Identification	<p>Sponsorship source: supported in part by Grant No. HL-31558 from the National Institutes of Health and by a grant from Sigma Theta Tau, Psi Chapter</p> <p>Country: USA</p> <p>Author's name: Janet L. Larson</p> <p>Institution: College of Nursing, University of Illinois at Chicago</p> <p>Address: 845 S.Damen Avenue, Chicago,IL 60612</p>
Notes	

Larson 1999
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/14

Larson 1999 (Continued)

- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 68 (6)
- BMI, mean (SD), kg/m²: 27 (4)

PR

- N (randomized/analyzed): not reported/14
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 66 (6)
- BMI, mean (SD), kg/m²: 26 (4)

IMT

- N (randomized/analyzed): not reported/13
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 66 (5)
- BMI, mean (SD), kg/m²: 28 (4)

Control/sham

- N (randomized/analyzed): /12
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 62 (7)
- BMI, mean (SD), kg/m²: 26 (5)

Overall

- N (randomized/analyzed): 130/53
- Loss to follow-up or excluded: 83
- COPD stage (GOLD): moderate to severe

Included criteria

- Patients between 45 and 75 years of age;
- Moderate to severe airflow obstruction (FEV1 < 65%pred and FEV1/FVC < 70%)
- Complaints of dyspnea on exertion
- Clinically stable condition
- No participation in a pulmonary rehabilitation program in the last year

Excluded criteria

- A history of asthma
- Experienced a major exacerbation in the 2 months before enrollment
- Took > 10 mg of prednisone/d
- Required home oxygen therapy or experienced oxyhemoglobin desaturation < 85% with exercise, and/or had other health problems that would interfere with exercise

Interventions

Intervention characteristics

PR: participants performed cycle ergometer training at home on a calibrated stationary cycle ergometer (BodyGuard 990; BodyGuard, Sandnes, Norway), 20 min/d, 5 d/week for 2 months. An interval training protocol was used with participants performing 4 work sets, 5 min in duration, separated by rest intervals (2–4 min) of unloaded cycling. The training was initiated at 50% of the peak work rate, taken from the best baseline graded exercise test, and evaluated weekly with progressive increases as tolerated. Patients were instructed to pedal at a rate of 60 revolutions/minute (rpm), and they were encouraged to push themselves to the limits of their dyspnea, without exceeding a heart rate equal to 85% of the predicted maximal heart rate.

IMT: the participants trained with Threshold IMT (HealthScan, Cedar Grove, NJ), 30 min/d, 5 d/week for 2 months. the training was initiated at 30% of PImax with progressive increases up to 60% of PImax

Larson 1999 (Continued)

PR+IMT: participants received both PR and IMT as described above.

Control/sham: this group participated in a structured program of health education

Outcomes	<p>Dyspnea: Borg</p> <ul style="list-style-type: none"> • P_{thmax} • W_{max} • Submaximal exercise at 50% of W_{max} <p>HRQoL: CRQ</p> <ul style="list-style-type: none"> • Dyspnea • Fatigue <p>Respiratory muscle strength: P_{lmax} (RV)</p> <p>Respiratory muscle endurance: respiratory muscle endurance pressure (P_{thmax})</p> <p>Functional exercise capacity: W_{max}</p> <p>Laboratory exercise test: VO_{2peak} (L/min)</p>
Identification	<p>Sponsorship source: funded by a research grant from the National Institute of Nursing Research, National Institutes of Health, RO1-NR01428</p> <p>Country: USA</p> <p>Author's name: Janet L. Larson</p> <p>Institution: University of Illinois at Chicago</p> <p>Email: LLarson@uic.ed</p> <p>Address: 845 S. Damen, Chicago, IL60612</p>
Notes	

Leelarungrayub 2017

Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 12/10 • Loss to follow-up or excluded: 2 • Age, mean (SD) in years: 63.25 (1.49) • Gender (M/F): 4/6 • BMI, mean (SD), kg/m²: 20.6 (1.9) <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 12/10 • Loss to follow-up or excluded: 2

Leelarungrayub 2017 (Continued)

- Age, mean (SD) in years: 68.75 (2.39)
- Gender (M/F): 6/4
- BMI, mean (SD), kg/m²: 22.1 (2.4)

Overall

- N (randomized/analyzed): 24/20
- Loss to follow-up or excluded: 2
- Gender (M/F): 10/10
- COPD stage (GOLD): moderate

Included criteria

- All the participants were ex-smokers
- They had a stable clinical condition during the experiments

Excluded criteria

- Uncontrolled hypertension, unstable cardiac disease, recurrent symptoms of pneumothorax, thoracic or chest pain including neuromuscular disorders, liver diseases or endocrine abnormalities
- Participants taking supplements or any nutrients such as vitamins or N-acetylcysteine compounds during this study

Interventions	<p>Intervention characteristics</p> <p>IMT: participants trained with Portex (Smith Medical ASD), 20-30 min/session, 7 d/week for 6 weeks. They started breathing through a 6 mm hole once daily for the first 2 weeks, before changing to 4 mm and 2 mm holes in the 2nd and 4th week, respectively. 30 slowly repeated inspirations passed through the device, with a 3-min interval of rest in each of 4 training sessions.</p> <p>Control: this group did not receive any intervention</p>
Outcomes	<p>Functional exercise capacity: 6MWD</p> <p>HRQoL: CCQ</p> <ul style="list-style-type: none"> • Symptom score • Mental score • Function score • Total score <p>Respiratory muscle strength: P_{Imax} (RV)</p> <p>Respiratory function: FEV₁</p> <ul style="list-style-type: none"> • L • %pred
Identification	<p>Sponsorship source:</p> <p>Country: Thailand</p> <p>Setting: Sansai hospital, Sansai, Chiang Mai Province, Thailand</p> <p>Author's name: Jirakrit leelarungrayub</p> <p>Institution: Department of Physical Therapy, Faculty of associated Medical sciences, Chiang Mai University</p> <p>Email: donrawee.leela@cmu.ac.th</p> <p>Address: Intawaroraj road, Sripoom, Chiang Mai 50200, Thailand</p>

Leelarungrayub 2017 (Continued)

Notes	<p>We calculated the SD manually from the SE ($SD = SE \cdot \sqrt{N}$).</p> <p>This study also used a prototype that was excluded from our analysis because it hasn't been validated yet. We contacted the Portex manufacturer to ask for further information about the device.</p>
-------	---

Lisboa 1997
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 10/10 • Loss to follow-up or excluded: not reported • Age, mean (SD) in years: 61 (6.32) • Age (SD): 6.32 • Gender (M/F): 6/4 <p>Control/Sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 10/10 • Loss to follow-up or excluded: not reported • Age, mean (SD) in years: 64 (6.32) • Age (SD): 6.32 • Gender (M/F): 7/3 <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 20/20 • Loss to follow-up or excluded: 0 • Gender (M/F): 13/7 • COPD stage (GOLD): severe <p>Included criteria</p> <ul style="list-style-type: none"> • A stable period of their disease • Dyspnea during ADL • FEV1/FVC < 60% • The absence of cardiac or any other disease that could interfere with exercise performance <p>Excluded criteria: not reported</p>
Interventions	<p>Intervention characteristics</p> <p>IMT: participants trained at home using a Threshold IMT device (HealthScan Products Inc., NJ, USA), for 30 min/d, 6 d/week for 10 weeks at 30% of P_Imax</p> <p>Control/sham: this group received a similar IMT protocol and trained at 10% of P_Imax</p>
Outcomes	<p>Dyspnea: BDI-TDI (Total)</p> <p>Dyspnea: Borg</p> <ul style="list-style-type: none"> • Notes: It was evaluated under basal conditions

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

93

Lisboa 1997 (Continued)

Functional exercise capacity: 6MWD

Respiratory muscle strength: P_{Imax} (FRC)

Respiratory function: FEV₁ (L)

Functional exercise capacity: W_{max}

- Notes: we considered 1W = 6.11 kpm/min

Laboratory exercise test: VO_{2peak} (mL/min)

Identification

Country: Chile

Author's name: C. Lisboa

Institution: Dept of respiratory disease, Catholic University of Chile

Address: Santiago, Chile

Notes

The authors mentioned COPD in the discussion section.

Mador 2005
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 19/15
- Loss to follow-up or excluded: 4
- Age mean (SD) in years: 69.7 (7.7)
- BMI, mean (SD), kg/m²: 28.3 (8.9)

PR

- N (randomized/analyzed): 19/14
- Loss to follow-up or excluded: 5
- Age mean (SD) in years: 70.9 (7.4)
- BMI, mean (SD), kg/m²: 26.9 (3.3)

Overall

- N (randomized/analyzed): 38/29
- Loss to follow-up or excluded: 9
- COPD stage: moderate to very severe

Included criteria

- Confirmed diagnosis of COPD

Interventions

Intervention characteristics
PR+IMT

Mador 2005 (Continued)

- PR: participants trained on a cycle ergometer, initially at 50% of Wmax and keeping Borg score ≤ 5 during exercise. They also trained on a treadmill at a speed ranging from 1.1-2.0 miles per hour (1.7-3.2 km/h) at 0% elevation based on the participant's functional capacity (i.e. on 6MWD results). When the participants could exercise for 20 min without intolerable dyspnea or leg fatigue, the speed and/or elevation was increased.
- IMT: participants trained 15-20 min/d, 3 d/week for 8 weeks, and breathed from a rebreathing bag while obtaining fresh air through a side port. V \dot{V} E and PetCO $_2$ were continuously recorded using a metabolic cart (Medgraphics; St. Paul, MN), and oxygen saturation was measured by pulse oximetry. Rebreathing bags of 1.5-2.0 L, depending on the participant's vital capacity, were used, and the size of the bag was additionally adjusted with a clamp until stable normocapnia as estimated by the PetCO $_2$ was obtained during preliminary trials.

PR: this group received only the PR program described above

Outcomes	Functional exercise capacity: <ul style="list-style-type: none"> • 6MWD • Exercise time • Wmax HRQoL: CRQ Respiratory muscle strength: P I_{max} Respiratory muscle endurance time: T $_{lim}$
Identification	<p>Country: USA</p> <p>Author's name: M. Jeffery Mador</p> <p>Institution: Division of Pulmonary, Critical Care, and Sleep Medicine, Section 111S</p> <p>Email: mador@acsu.buffalo.edu</p> <p>Address: State University of New York at Buffalo, Veterans Administration Medical Center, 3495 Bailey Ave, Buffalo, NY 14215</p>
Notes	

Magadle 2007
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 16/14 • Loss to follow-up or excluded: 2 • Age (mean): 65.2 (13.6) <p>PR</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 15/13 • Loss to follow-up or excluded: 2

Magadle 2007 (Continued)

- Age (mean): 66.1 (12.39)

Overall

- N (randomized/analyzed): 31/29
- Loss to follow-up or excluded: 2
- COPD stage (GOLD): moderate to severe

Included criteria

- Patients with spirometric evidence of significant chronic airflow limitation (FEV1 < 50%pred, FEV1/FVC < 70%pred) and were diagnosed as having COPD according to ATC criteria

Excluded criteria

- Patients with cardiac disease, poor compliance, or requirement of supplemental oxygen

Interventions	Intervention characteristics PR+IMT <ul style="list-style-type: none"> • PR: this program included lower extremity endurance exercise (walking or cycling), upper extremity exercise and strength training with free weights. This phase included 36 sessions of 90 min duration (3 times/week for 12 weeks). In the 2nd phase, participants trained 1 h 3 times/week for 6 months. • IMT: started in the 2nd phase of the trial, and consisted of training 30 min/session, 3 times/week for 6 months. They used the Threshold IMT device at a training load ranged from 15%- 60% of PImax. PR (+sham IMT): this group received a similar training protocol and participants performed IMT at no load.
Outcomes	Functional exercise capacity: 6MWD HRQoL: SGRQ (Total) Respiratory muscle strength: PImax (RV) Respiratory function: FEV1 (%pred)
Identification	Country: Israel Setting: community-based rehabilitation center Author's name: Paltiel Weiner Institution: Department of Medicine A, Hillel Yaffe Medical Center Email: weiner@hillel-yaffe.health.gov.il Address: Hadera 38100, Israel
Notes	Participants were enrolled in a 1st phase that consisted of 36 sessions of PR for 3 months (90 min/session).

Majewska-Pulsakowska 2016
Study characteristics

Methods	Study design: RCT Study grouping: parallel-group
---------	---

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

96

Majewska-Pulsakowska 2016 (Continued)

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 13/13
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 61.5 (6.1)
- Gender (M/F): 10/3
- BMI, mean (SD), kg/m²: 28.8 (6.2)

PR

- N (randomized/analyzed): 9/9
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 62.3 (5.2)
- Gender (M/F): 6/3
- BMI, mean (SD), kg/m²: 28.2 (6.2)

IMT

- N (randomized/analyzed): 8/8
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 63.4 (9.8)
- Gender (M/F): 2/6
- BMI, mean (SD), kg/m²: 26.1 (5.9)

Control/sham

- N (randomized/analyzed): 13/13
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 65.5 (7.0)
- Gender (M/F): 7/6
- BMI, mean (SD), kg/m²: 27.8 (4.9)

Overall

- N (randomized/analyzed): 43/43
- Loss to follow-up or excluded: 0
- Gender (M/F): 25/18
- COPD stage (GOLD): moderate to severe

Included criteria

- COPD treated for at least 1 year, stage 2 and 3 according to GOLD (2014)
- 50–70 years old
- A stable clinical condition with no exacerbations over the period of 4 weeks before the study

Excluded criteria

- Participation in PR in the year preceding the study
- Diagnosed bronchial asthma
- Long-term home oxygen therapy
- Clinically significant diseases of the cardiovascular system
- Any uncontrolled chronic disease; muscle and nervous disorders reducing the patient's mobility
- Mental disorders preventing contact and co-operation with the patient

Interventions

Intervention characteristics

Majewska-Pulsakowska 2016 (Continued)

PR: consisted of training on a cycle ergometer, 3 times/week for 8 weeks in an ambulatory setting under the supervision of a cardiologist. Each training session began (warm-up) and finished (relaxation) with a pedalling load of 10 W for 3 min. The duration of a training session was initially 23 min, and then it was gradually increased up to 45 min.

IMT: consisted of home-based training using Threshold IMT device (Respironics; Philips Healthcare, DA Best, The Netherlands), twice a day (5-15 min), 5 d/week for 8 weeks. The training load ranged from 30%-60% of P_Imax

PR+IMT: this group received both interventions described above.

Control/sham: this group did not receive any intervention.

Outcomes	HRQoL: SGRQ Respiratory function: FEV1 <ul style="list-style-type: none"> • L • %pred
Identification	Country: Poland Author's name: K. Wytrychowski Institution: Department of Internal Diseases, Geriatrics and Allergology, Wrocław Medical University Email: Polande-mail:anhw@op.plk Address: 66 Skłodowskiej-Curie St., 50-369 Wrocław, Poland
Notes	

Masanga 2011

Study characteristics

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics PR+IMT <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/11 PR <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/9 Overall <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/21 Included criteria: stable patients with moderate to severe COPD Excluded criteria: not reported
Interventions	Intervention characteristics PR+IMT:

Masanga 2011 (Continued)

- PR: consisted of Occupational therapy, Education, Dietary instruction.
- IMT: at a training load that ranged from 10 cmH₂O to 40-90 cmH₂O.

PR:

Participants received only the PR program described above.

Outcomes	This abstract reported only adverse events: headache (6), jaw pain (6), neck pain (6), back pain (4), abdominal pain (2), cough (1), blood-streaked sputum (1), shoulder pain (1) and chest pain (1)
Identification	Country: Philippines Author's name: L. Masanga
Notes	

Nikoletou 2016
Study characteristics

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics IMT <ul style="list-style-type: none"> • N (randomized/analyzed): 34/21 • Loss to follow-up or excluded: 13 • Age, mean (SD) in years: 70.1 (8.4) • Gender (M/F): 14/9 • BMI, mean (SD), kg/m²: 24.9 (5.2) Control/sham <ul style="list-style-type: none"> • N (randomized/analyzed): 34/18 • Loss to follow-up or excluded: 16 • Age, mean (SD) in years: 71.1 (9.6) • Gender (M/F): 11/8 • BMI, mean (SD), kg/m²: 26.5 (8.1) Overall <ul style="list-style-type: none"> • N (randomized/analyzed): 68/39 • Loss to follow-up excluded: 29 • Gender (M/F): 25/17 • COPD stage (GOLD): moderate to very severe Included criteria <ul style="list-style-type: none"> • No exacerbation of COPD • Not changed their medication for at least 4 weeks prior to the initial assessment Excluded criteria <ul style="list-style-type: none"> • Patients with an a1-antitrypsin deficiency • Co-existing heart disease, hypertension, or long-term use of oral corticosteroids

Nikoleitou 2016 (Continued)

- Patients with significant thoracic musculoskeletal abnormalities, such as kyphosis or scoliosis
- Patients unsuitable for magnetic stimulation, for example, those with cardiac pacemakers

Interventions

Intervention characteristics

IMT: 2 sessions/d, 6 d/week for 7 weeks, home-based, using the Powerbreathe inspiratory muscle trainer (HaB International, Southam, Warwickshire, UK). Participants started training at 30% of their baseline P_{Imax} and increased the intensity once a week as tolerated. The average weekly increase in the intensity of training was 5% and the mean (SD) intensity at the end of the program was 62% (SD: 11.7) of the baseline P_{Imax}

Control/sham: participants received a similar IMT program and trained at 15% of P_{Imax}

Outcomes

Functional exercise capacity: incremental SWT

Dyspnea: Borg

- Notes: Borg post-incremental SWT

HRQoL: CRQ

- Dyspnea
- Fatigue
- Emotion
- Mastery

HRQoL: SF-36 Questionnaire

- Physical functioning
- Physical problems
- Emotional problems
- Social functioning
- Mental health
- Energy/ Vitality
- Pain
- General health perception
- Change in health

Respiratory muscle strength: P_{Imax} (FRC)

Respiratory muscle endurance time: T_{lim}

Identification

Country: UK

Setting: respiratory outpatient clinics at King's College Hospital, GP practices and British Lung Foundation Breathe Easy groups

Author's name: Dimitra Nikoleitou

Institution: School of Rehabilitation Sciences, Faculty of Health, Social care and Education, Kingston University and St George's University of London

Email: D.Nikoleitou@sgul.kingston.ac.uk

Address: Grosvenor Wing, Cranmer Terrace, London SW17 0RE, UK

Notes

Paneroni 2018
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 15/12 • Loss to follow-up or excluded: 3 • Age, mean (SD) in years: 67.8 (9.8) • Gender (M/F): 10/2 • BMI, mean (SD), kg/m²: 26.9 (5.3) <p>PR</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 11/10 • Loss to follow-up or excluded: 1 • Age, mean (SD) in years: 65.8 (25.0) • Gender (M/F): 8/2 • BMI, mean (SD), kg/m²: 25.0 (5.1) <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 26/26 • Loss to follow-up or excluded: 4 • Gender (M/F): 18/4 • COPD stage (GOLD): mild to very severe <p>Included criteria</p> <ul style="list-style-type: none"> • MVV of < 90% of predicted • The ability to perform the 6MWD • The absence of chronic respiratory failure (arterial oxygen (PaO₂) and carbon dioxide (PaCO₂) tensions at rest > 60 and < 45 mmHg, respectively) <p>Excluded criteria</p> <ul style="list-style-type: none"> • Refusal to participate • Cognitive impairment as assessed by a Mini-Mental State Examination score of < 25 points • Hemodynamic instability, and severe clinical events (e.g. COPD exacerbations or heart failure treated with infusions of vasopressors and/or inotropes) in the month before enrollment
Interventions	<p>Intervention characteristics</p> <p>PR+IMT:</p> <ul style="list-style-type: none"> • PR: incremental exercise training according to Maltais et al (Maltais 1997) until performing 30 min of continuous cycling at 50%–70% of the maximal load calculated based on the initial 6MWD plus 30-min sessions of abdominal, upper, and lower limb muscle activities, lifting progressively increasing weights (300–500 g), shoulder and full arm circling, and other exercises according to Clark et al (Clark 1996) • IMT: participants received twice a day, 5 d/week for 2 weeks, 20 15-min sessions of normocapnic hyperpnea training using SpiroTiger device (Idiag; Fehraltorf, Switzerland). The initial target of minute ventilation (VE) was 66% of participants' MVV. Once the participant was able to perform 15-min training sessions without interruptions, the target VE was increased to 75% of MVV, by increasing only the respiratory rate. Thereafter, training involved further 10% increments of MVV, when the participant could perform without interruption three 15-min sessions at the set level

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

101

Paneroni 2018 (Continued)

PR (+sham IMT): participants in this group received a similar training PR program, and they performed IMT using Threshold IMT device at no resistance

Outcomes	Functional exercise capacity: 6MWD Respiratory muscle strength: P _{lmax} Respiratory muscle endurance time: T _{lim} (normocapnic hyperpnea) <ul style="list-style-type: none"> • Notes: it is the sustained VE at 66% of their MVV. If there was no exhaustion in 15 min, the test was repeated the following day at 75% of MVV Respiratory muscle endurance time: T _{lim} (Threshold device) <ul style="list-style-type: none"> • Notes: the pressure was set at 30% of the patients' baseline P_{lmax} Respiratory muscle endurance: MVV
----------	---

Identification	<p>Country: Italy</p> <p>Setting: Respiratory Rehabilitation Divisions of the Lumezzane and Pavia Institutes of the Istituti Clinici Scientifici Maugeri, Italy</p> <p>Author's name: Mara Paneroni</p> <p>Institution: Istituti Clinici Scientifici Maugeri, IRCCS</p> <p>Address: Salvatore Maugeri 2, 27100 Pavia, Italy</p> <p>Clinical trial register: NCT01556139</p>
----------------	---

Notes

Petrovic 2012
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 10/10 • Loss to follow-up or excluded: 0 • Age, mean (SD) in years: 58.7 (5.2) • Gender (M/F): 6/4 • BMI, mean (SD), kg/m²: 24.3 (4.0) <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 10/10 • Loss to follow-up or excluded: 0 • Age, mean (SD) in years: 60.3 (5.3) • Gender (M/F): 5/5 • BMI, mean (SD), kg/m²: 24.2 (3.4)

Petrovic 2012 (Continued)

Overall

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0
- Gender (M/F): 11/9
- COPD stage (GOLD): moderate to severe

Included criteria

- Required participants to be < 70 years old
- Moderate to severe non-reversible airflow obstruction

Excluded criteria

- Exacerbation of COPD within the previous 6 weeks
- Neoplastic disease or the presence of a disease that could contribute to dyspnea, or exercise limitation (cardiovascular, neuromuscular, or other respiratory diseases)
- No patients had received treatment with systemic cortisone over the last 6 months

Interventions

Intervention characteristics

IMT: performed once/d, 7 d/week, for 8 weeks using Respifit Sdevice (Mauerbach, Australia). The training consisted of strength training at 80% of P_{Imax} (≈ 12 min each session) and endurance training at 60% of P_{Imax} (≈ 13.5 min each session)

Control/sham: this group did not receive any intervention

Outcomes

Dyspnea: Borg

- Constant cycle ergometer test
- Incremental cycle ergometer test
- Notes: the sensation of dyspnea was assessed every 2 min during exercise

Respiratory muscle strength: P_{Imax}

Respiratory muscle endurance time: T_{lim}

- Notes: measured through breathing against 60% P_{Imax}

Laboratory exercise test: VO_{2peak} (mL/min)

- Constant cycle ergometer test
- Incremental cycle ergometer test

Identification

Country: Austria

Setting: outpatient clinic

Author's name: Milos Petrovic

Institution: Pulmonary Department and Karl Landsteiner Institute for Clinical and Experimental Pulmology

Email: milos.petrovic@wienkav.at

Address: Hietzing Hospital, Wolkensbergenstrasse 1, 1130 Vienna, Austria

Clinical trial register: NCT00469313

Notes

Preusser 1994

Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): /12 • Loss to follow-up or excluded: 1 • Age, mean (SD) in years: 68/10 • Gender (M/F): 6/6 <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): /8 • Loss to follow-up or excluded: 1 • Age, mean (SD) in years: 62/17 • Gender (M/F): 1/7 <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 22/20 • Loss to follow-up or excluded: 2 • Gender (M/F): 7/13 • COPD stage (GOLD): severe to very severe <p>Included criteria</p> <ul style="list-style-type: none"> • The presence of COPD as defined by the ATS • Willingness to participate in a 3-month inspiratory muscle training protocol <p>Excluded criteria</p> <ul style="list-style-type: none"> • Uncontrolled cardiac disease, diabetes, hypertension, musculoskeletal, and neuromuscular diseases
Interventions	<p>Intervention characteristics</p> <p>IMT: consisted of supervised sessions, 3 days/week for 12 weeks, using Threshold IMT device at 52% of P_Imax. The training duration ranged from 5 min in week 1 to 18 min in week 12</p> <p>Control/sham: this group received a similar IMT protocol and trained at 22% of P_Imax</p>
Outcomes	<p>Functional exercise capacity: 12MWD</p> <p>Respiratory muscle strength: P_Imax (FRC)</p> <p>Respiratory muscle endurance: Respiratory muscle endurance pressure (P_{th}max)</p> <ul style="list-style-type: none"> • Notes: "After quietly breathing through an unloaded system for 2 min, the threshold load began at -4 cm H₂O pressure and increased by -2 cm after every fifth breath (every 15 s) until the subject either signalled to stop the test or was unable to match respiratory rate, duty cycle and flow rate during three of the five breaths for that pressure load. The maximal threshold load successfully completed during the ramp test was recorded"
Identification	<p>Sponsorship source: supported by grant F31 NR06378, National Center for Nursing Research; a grant from the American Nurses Foundation; and the Epsilon Chapter of Sigma Theta Tau</p>

Preusser 1994 (Continued)

Country: USA

Author's name: Barbara A. Preusser

Institution: College of Nursing, The Ohio State University, and the Department of Pulmonary Medicine, The Ohio State University Hospitals, Columbus

Address: University of Utah College of Nursing, 25 South Medical Drive, Salt Lake City 84112

Notes

Ramirez Sarmiento 2002

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): not reported/7
- Loss to follow-up or excluded: not reported
- Age, mean (SD) in years: 65
- BMI, mean (SD), kg/m²: 29

Control/sham

- N (randomized/analyzed): not reported/7
- Loss to follow-up or excluded: 2
- Age, mean (SD) in years: 66 (6)
- BMI, mean (SD), kg/m²: 26 (4)

Overall

- N (randomized/analyzed): 16/14
- Loss to follow-up or excluded: 2
- BMI, mean (SD), kg/m²:
- COPD stage (GOLD): severe to very severe

Included criteria

- Patients with COPD

Excluded criteria

- Hypoxemia (PaO₂ < 60 mmHg breathing room air)
- Asthma; coronary disease; undernourishment (BMI < 20 kgm²)
- Chronic metabolic disease
- Orthopedic disease
- Previous abdominal or thoracic surgery
- Treatment with steroids, hormones or cancer chemotherapy

Interventions

Intervention characteristics

Ramirez Sarmiento 2002 (Continued)

IMT: participants trained 30 min/day, 5 d/week for 5 weeks. The sessions were supervised and using Threshold IMT device set at 50% of P_Imax

Control/sham: participants received a similar IMT protocol and trained at no load

Outcomes	Functional exercise capacity: <ul style="list-style-type: none"> • 6MWD • Wmax Respiratory muscle strength: P _I max Respiratory muscle endurance: Respiratory muscle endurance pressure (P _{th} max) <ul style="list-style-type: none"> • Notes: breathing against incremental loads (8 cmH₂O) every 2 min until maximal sustainable threshold pressure was reached Respiratory muscle endurance time: T _{lim} <ul style="list-style-type: none"> • Notes: breathing against a constant a submaximal constant load (equivalent to 80% of maximal threshold pressure) Laboratory exercise test: VO ₂ peak (mL/kg/min) <ul style="list-style-type: none"> • Notes: measured during incremental cycle test Respiratory function: FEV ₁ (%pred) Respiratory function: RV (%pred)
Identification	<p>Sponsorship source: supported, in part, by grants BIOMED (BMH-4-CT98-3406), FIS, SEPAR, and SIBEL</p> <p>Country: Spain</p> <p>Author's name: Mauricio Orozco-Levi</p> <p>Institution: Servei Grup de Recerca de Pneumologica, Hospital del MarIMIM</p> <p>Email: morozco@imim.es</p> <p>Address: Passeig Maritim 25, E-080003, Barcelona (Catalonia), Spain</p>
Notes	

Saher 2021
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 23/17 • Loss to follow-up or excluded: 6 • Age, mean (SD) in years: 62.8 (6.72) • Gender (M/F): 9/5

Saher 2021 (Continued)

- BMI, mean (SD), kg/m² : 22.7 (4.25)

Control

- N (randomized/analyzed): 23/17
- Loss to follow-up excluded: 6
- Age, mean (SD) in years: 61.8 (11.93)
- Gender (M/F): 12/8
- BMI, mean (SD), kg/m² : 23.7 (5.51)

Overall

- N (randomized/analyzed): 46/34
- Loss to follow-up or excluded: 12
- Gender (M/F): 21/13
- COPD stage (GOLD): moderate to very severe

Included criteria

- Hypercapnic moderate to severe COPD patients
- Patients were required to have hypercapnic respiratory failure (PaCO₂ > 46 mmHg), already receiving non-invasive ventilation for at least 8 h/d, and with a decreased inspiratory muscle strength (PImax < 60 cm H₂O)

Excluded criteria

- Patients with any other severe respiratory disease apart from COPD, or with any other severe non-pulmonary disease limiting prognosis (e.g. metastatic cancer, active tuberculosis, congestive heart failure, liver cirrhosis)

Interventions

Intervention characteristics

IMT: participants performed IMT 15 min/session, 2 session/day, for 10 days. They used an IMT threshold device (Powerbreath, Gaia Ltd., Southam, UK). The training load started at 30% of PImax, and it was increased by 5%-10% daily until 60%

Control: this group did not receive any training

Note: both groups received non-invasive ventilation (NIV) as part of COPD management.

Outcomes

Functional exercise capacity: 6MWD

Inspiratory muscle strength (PImax)

Identification

Country: India

Setting: Metro Center for Respiratory Disease, Metro Hospital

Author's name: T. Saher

Institution: Center for Physiotherapy and Rehabilitation Sciences

Email: jmoiz@jmi.ac.in

Address: Jamia Millia Islamia, New Delhi, 110025, India

Notes

Saka 2021
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 23/20 • Loss to follow-up or excluded: 3 • Age, mean (SD) in years: 62.30 (7.43) • Gender (M/F): 19/1 • BMI, mean (SD), kg/m² : 26.04 (4.41) <p>Sham IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 22/20 • Loss to follow-up excluded: 2 • Age, mean (SD) in years: 62.10 (7.76) • Gender (M/F): 18/2 • BMI, mean (SD), kg/m² : 27.11 (4.88) <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 45/40 • Loss to follow-up or excluded: 5 • Gender (M/F): 37/3 • COPD stage (GOLD): moderate to severe <p>Included criteria</p> <ul style="list-style-type: none"> • FEV1:FVC ratio < 70% in PFT • Age: ≥ 18 years old • Being able to read written and understand spoken language <p>Excluded criteria</p> <ul style="list-style-type: none"> • A history of COPD exacerbation in the last 6 weeks • The presence of comorbidities affecting ambulation/activity (e.g. severe cardiac or neurological disorders, cancer, musculoskeletal problems) and cognitive disorders (Mini-Mental State Examination < 24)
Interventions	<p>Intervention characteristics</p> <p>IMT: the training was performed 15 min twice a day, 5 d/week, for 8 weeks. Home-based training with 1 session supervised by a physiotherapist. Participants used Threshold IMT device (Threshold IMT® Philips Respironics, UK) at a load of 30% of PImax.</p> <p>Sham IMT: participants received the same protocol with a training load set at 15% of PImax.</p>
Outcomes	<p>Dyspnea: mMRC</p> <p>Functional exercise capacity: 6MWD</p> <p>HRQoL: SGRQ</p> <ul style="list-style-type: none"> • Symptoms • Activity

Saka 2021 (Continued)

- Impacts
 - Total
- HRQoL: CAT
- Inspiratory muscle strength (PImax)
- Respiratory function: FEV1
- L
 - %pred

Identification

Sponsorship: Scientific Research Projects Unit of Bezmialem Vakif University, project number 9.2017/31

Country: Turkey

Setting: Bezmialem Vakif University Division of Physiotherapy and Rehabilitation, Cardiopulmonary Physiotherapy and Rehabilitation Department.

Author's name: Seda Saka

Institution: Cardiopulmonary Physiotherapy Rehabilitation Department, Institute of Health Sciences, Bezmialem Vakif University, Istanbul, Turkey

Email: fztседasaka@gmail.com

Address: Division of Physiotherapy and Rehabilitation, Faculty of Health Sciences, Bezmialem Vakif University, Silahataraga St. No:189, Alibeykoy, 34060, Istanbul, Turkey

Clinical trial register: NCT03517839

Notes

Sanchez Riera 2001

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): 10/10
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 67 (4)
- Gender (M/F): 9/1

Control/Sham

- N (randomized/analyzed): 10/10
- Loss to follow-up/excluded: 0
- Age, mean (SD) in years: 67.6 (5)
- Gender (M/F): 9/1

Overall

Sanchez Riera 2001 (Continued)

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0
- Gender (M/F): 18/2
- COPD stage (GOLD): moderate to severe

Included criteria

- The presence of COPD as defined by the ATS
- Patients were in stable condition

Excluded criteria

- Clinical evidence of cardiovascular, musculoskeletal, or neuromuscular disease or of any other disease that might interfere with exercise

Interventions	<p>Intervention characteristics</p> <p>IMT: participants trained at home 30 min/d, 6 d/week for 6 months using incentive flowmeter device (INSPIRx; Intertech Resources Inc; Ft. Myers, FL) set at 30% of P_Imax. The breathing pattern was controlled during the exercise.</p> <p>Control/sham: participants received a similar IMT protocol and trained at no load.</p>
Outcomes	<p>Dyspnea: Borg (Constant cycle ergometer test)</p> <p>Functional exercise capacity: SWT</p> <ul style="list-style-type: none"> • Notes: "Patients were required to walk 10 m back and forth. The walking speed was paced by an audio signal from a cassette that emitted beeps at regular intervals. The speed was increased each minute by 0.17 m/s until the next level was attained. The end of the test was determined by patients when they were too breathless to maintain the required speed, or by operators, if patients failed to complete a shuttle in the time allowed" <p>Functional exercise capacity: W_{max}</p> <ul style="list-style-type: none"> • Notes: "performed on a cycle ergometer. After 1 min of unloaded pedalling, the work rate was increased 10 W/min at a time. The test was stopped when patients were unable to continue because of dyspnea or leg fatigue" <p>Respiratory muscle strength: P_Imax (FRC)</p> <p>Laboratory exercise test: VO₂peak (L/min)</p> <p>Respiratory muscle endurance: Respiratory muscle endurance pressure (P_{thmax})</p>
Identification	<p>Sponsorship source: Supported by "Junta de Andalucia" grant No. 94//535-119</p> <p>Country: Spain</p> <p>Setting: home-based training</p> <p>Author's name: Hildegard Sanchez Riera</p> <p>Institution: Pneumology Service, Virgen Del Rocio University Hospital</p> <p>Email: ablucil@mx2.redestb.es</p> <p>Address: "La Motilla," C/Rayo 4, 41700 Dos Hermanas, Sevilla, Spain</p>
Notes	<p>CRQ was not included because study did not fully report control group data</p>

Scherer 2000
Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/15 • Loss to follow-up or excluded: not reported • Age, mean (SD) in years: 66.9 (9.2) • Gender (M/F): 9/6 • BMI, mean (SD), kg/m²: 23.8 (3) <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 15 • Loss to follow-up or excluded: not reported • Age, mean (SD): 71.0 (4.6) • Gender (M/F): 10/5 • BMI, mean (SD), kg/m²: 25.9 (3.48) <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 34/30 • Loss to follow-up or excluded: 4 • Range age (min, max): 46,80 • Gender (M/F): 19/11 • COPD stage (GOLD): moderate to severe <p>Included criteria</p> <ul style="list-style-type: none"> • Chronic airflow obstruction (FEV1 < 70%pred, FEV1/FVC < 70%pred, < 15% improvement in FEV1 after bronchodilatation with 200 mg of albuterol inhaled from a pressurized metered-dose inhaler with a spacer) • Aged 20-80 years • A stable clinical condition for at least 1 month • The patients' physical activity had to be limited by pulmonary dyspnea only <p>Excluded criteria</p> <ul style="list-style-type: none"> • Patients with dyspnea at rest • Cardiac disease, poor compliance, drug or alcohol abuse, pregnancy or lactation, a requirement for supplemental oxygen, CO2 retention, or use of any mechanical ventilatory support
Interventions	<p>Intervention characteristics</p> <p>IMT: respiratory muscle endurance training was conducted twice daily, 5 d/week for 8 weeks. The device was developed and consisted of tubing (I.D. 5 19 mm) connecting a rebreathing bag with a mouth-piece at a 90° angle. The breathing frequency was 60% of MVV.</p> <p>Control/sham: participants trained following the same pattern of the description above, at no load using incentive spirometer (COACH 2 Volumetric Incentive Spirometer; DHD Healthcare, Canastota, NY). The respiratory rate was 6-8 breaths/min, and the target inspiration was set at 70% of each participant's ventilatory capacity.</p>

Scherer 2000 (Continued)

Outcomes	<p>Dyspnea: BDI-TDI (Total)</p> <p>Functional exercise capacity:</p> <ul style="list-style-type: none"> • 6MWD • Exercise time (treadmill) <p>Respiratory muscle strength: P_Imax (RV)</p> <p>Laboratory exercise test: VO₂peak (mL/kg/min)</p> <p>Respiratory muscle endurance time: T_{lim}</p> <ul style="list-style-type: none"> • Notes: respiratory muscle endurance was measured as sustained ventilation at 66% of each participant's highest MVV. The time during which participants were able to sustain this target ventilation was recorded. If a participant surpassed 15 min of breathing at this level, the test was repeated on the following day at 75% of MVV. <p>Respiratory function: FEV₁ (%pred)</p>
----------	---

Identification	<p>Sponsorship source: supported by grants from Astra Pharmaceutica, Dietikon, and Merck Sharpe and Dohme-Chibret, and Rhône-Poulenc Rorer</p> <p>Country: Switzerland</p> <p>Setting: outpatient clinic of the Pulmonary Division of the Triemli Hospital</p> <p>Author's name: Thomas A. Scherer</p> <p>Institution: LungenZentrum Hirslanden</p> <p>Email: thsche@swissonline.ch</p> <p>Address: Witellikerstrasse 36, 8008 Zurich, Switzerland</p>
----------------	---

Notes

Schultz 2018

Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
---------	--

Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 307/300 • Loss to follow-up or excluded: 26 • Age, mean (SD) in years: 57.7 (8.2) • Gender (M/F): 188/112 • BMI, mean (SD), kg/m²: 26.6 (6.4) <p>PR</p> <ul style="list-style-type: none"> • N (randomized): 304/302 • Loss to follow-up or excluded: 24 • Age, mean (SD) in years: 57.9 (6.6)
--------------	--

Schultz 2018 (Continued)

- Gender (M/F): 201/101
- BMI, mean (SD), kg/m²: 26.9 (6.6)

Overall

- N (randomized): 611/602
- Loss to follow-up or excluded: 50
- Gender (M/F): 389/213
- COPD stage (GOLD): moderate to severe

Included criteria

- Medical history of COPD
- FEV1/FVC < 70% and FEV1 %pred < 80% post-bronchodilation

Excluded criteria

- Lack of language or cognitive abilities to fill out questionnaires
- Hypercapnic respiratory failure (arterial carbon dioxide tension > 50 mmHg at rest)
- Indication for intermittent noninvasive ventilation
- Contraindications for inspiratory muscle training (e.g. a history of recent lung surgery, recent pulmonary embolism; a history of recurrent spontaneous pneumothorax)
- Severe comorbidities that confer significantly greater morbidity than COPD (e.g. active cancer without successfully completed curative tumor therapy)

Interventions	Intervention characteristics PR+IMT: <ul style="list-style-type: none"> • PR: consists of 2 components <ul style="list-style-type: none"> ◦ Obligatory components: (mostly 30- to 60-min sessions) including physical training (endurance training: 4 or 5 sessions per week; strength training: three sessions/week; whole-body vibration muscle training: 7 sessions/week), patient education (≥ 7 sessions) and respiratory physiotherapy in groups (2-4 sessions/week). ◦ Optional components: smoking cessation (8 sessions), mucolytic physiotherapy, saline inhalation, psychological interventions, social counselling, nutritional counselling and occupational therapy • IMT: conducted 21 min/d, 7 d/week for 3 weeks, using Threshold IMT device (POWERbreathe Medic; POWERbreathe International, Southam, UK). The initial training load was at least 30% of P_Imax and was progressively increased to at least 60%, and about half of the sessions were supervised <p>PR (+ sham IMT): this group received a similar training protocol as described above and the IMT was conducted at no load.</p>
Outcomes	Dyspnea: BDI-TDI (Total) Functional exercise capacity: 6MWD HRQoL <ul style="list-style-type: none"> • SGRQ (Total) • CAT • CCQ: (Total score) Respiratory muscle strength: P _I max Respiratory function: FEV1 (L)
Identification	<p>Sponsorship source: this study was supported by Deutsche Rentenversicherung Bayern Süd. Funding information for this article has been deposited with the Crossref Funder Registry</p> <p>Country: Germany</p>

Schultz 2018 (Continued)

Setting: Bad Reichenhall Clinic

Author's name: Konrad Schultz

Institution: Center for Rehabilitation, Pulmonology and Orthopedics, Klinik Bad Reichenhall

Email: konrad.schultz@klinik-bad-reichenhall.de

Address: Salzburger Strasse 8–11, 83435 Bad Reichenhall, Germany

Clinical trial register: DRKS00004609

Notes

Adjusted mean differences with 95% CI were reported

Sykes 2005

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0

PR

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0

Overall

- N (randomized/analyzed): 40/40
- Loss to follow-up or excluded: 0
- Range age (min, max): 60,84
- Gender (M/F): 34/6

Included criteria: not reported

Excluded criteria: not reported

Interventions

Intervention characteristics

PR+IMT: participants received exercise training and IMT with Threshold IMT at a load ranged from 30%-60% of P_{lmax}

PR: this group received only exercise training

Outcomes

Identification

Country: China

Setting: Tai Po Hospital

Author's name: Sykes

Sykes 2005 (Continued)

Notes

 The systematic review ([Gosselink 2011](#)) reported data from this trial.

Tounsi 2021
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 17/16
- Loss to follow-up or excluded: 1
- Age, mean (SD) in years: 62 (5)
- Gender (M/F): 17/0
- BMI, mean (SD), kg/m²: 23.13 (4.37)

PR

- N (randomized): 18/16
- Loss to follow-up or excluded: 2
- Age, mean (SD) in years: 63 (4)
- Gender (M/F): 18/0
- BMI, mean (SD), kg/m²: 23.41 (5.19)

Overall

- N (randomized): 35/32
- Loss to follow-up or excluded: 3
- Gender (M/F): 35/0
- COPD stage (GOLD): moderate to very severe

Included criteria

- patients with forced expiratory volume in 1 s (FEV1) <80% predicted and FEV1 /forced vital capacity (FVC) <70%
- Age between 45–75 years
- Clinically stable

Exclusion criteria

- Cardiovascular problem
- Psychiatric or cognitive disorders
- Progressive neuromuscular diseases
- Severe orthopedic problems with a significant impact on daily activities
- Prior inclusion in a rehabilitation program (<1 year)

Interventions

Intervention characteristics
PR+IMT:

- PR: an 8-week program with three sessions per week. the training program consists of aerobic supervised exercise training. The training consisted of 30 min of supervised treadmill exercise per session.

Tounsi 2021 (Continued)

Each session ended with upper and lower limb stretching. Each participant received an individualized program based on 60% to 80% of the average speed achieved during the six-minute walk test.

- **IMT:** The training was performed once a day, 7 days per week for 8 weeks, using a handle device (PowerBreathe1 Medic, IMT Technologies Ltd, Birmingham, UK). The training consists in making two sets of 30 breaths (4–5 min/set) with 5–10 min of rest between each set. The Respiratory resistive load was set at 50% of the initial P_Imax and then increased by 10% of the initial P_Imax every two weeks of training. Part of the IMT was performed and well instructed in the pulmonary rehabilitation center (3 days/ week) for 8 weeks; the other part was home-based training.

PR: this group received a similar training protocol as described above without IMT.

Outcomes	Functional exercise capacity: 6MWD Respiratory muscle strength: P _I max
Identification	<p>Sponsorship source: the authors received no funding for this work</p> <p>Country: Tunisia</p> <p>Setting: Farhat Hached Hospital of Sousse</p> <p>Author's name: Bilel Tounsi</p> <p>Institution: 1/ Laboratory of Exercise Physiology and Rehabilitation (APERE, UR-EA 3300), Sport Sciences Department, Picardie Jules Verne University, Amiens, France, 2/ Research Laboratory of Exercise Physiology and Pathophysiology: From Integral to Molecular Biology, Medicine and Health (LR19ES09), Faculty of Medicine of Sousse, University of Sousse, Sousse, Tunisia</p> <p>Email: bilel.tounsi@u-picardie.fr</p> <p>Address: Department of sports sciences, Picardie Jules Verne University, Amiens, France</p> <p>Clinical trial register: NCT04084405</p>
Notes	

Tout 2013

Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 10/10 • Loss to follow-up or excluded: 0 • Age, mean (SD) in years: 61 (9.32) <p>PR</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 10/10 • Loss to follow-up or excluded: 0 • Age, mean (SD) in years: 58.1 (8.72) <p>Overall</p>

Tout 2013 (Continued)

- N (randomized/analyzed): 20/20
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 60.38 (8.02)
- Gender (M/F): 19/21
- BMI, mean (SD), kg/m²: 28.3 (3.11)
- COPD stage (GOLD): mild to moderate

Included criteria

- Co-operating COPD patients
- Diagnosed clinically and through spirometric measurement (grades 1 and 2 in the Gold classification); presenting with 50% < FEV1 < 80% of predicted or theoretical value in the spirometric test
- Presenting with an improvement < 15% of the FEV1 following use of bronchodilators
- From 45-75 years of age
- Of either sex

Excluded criteria

- Heart failure or associated cardiac pathology
- Previous pulmonary or cardiac surgery
- Patient depending on oxygen therapy or undergoing cortisone treatment
- Associated neuromuscular pathologies

Interventions

Intervention characteristics

PR+IMT:

- PR: 16 rehabilitation sessions that consisted of bronchial decluttering meant to clear the airways; diaphragmatic rehabilitation (solicitation of the physiological diaphragmatic contraction) aimed at improved stamina; reinforcing of the lower limb muscles so as to limit functional deconditioning; psychological support and therapeutic education
- IMT: 20-30 min/session, twice/week for 8 weeks, using Threshold IMT at a training load ranged from 30%-60% of P_Imax

PR: this group received only the PR protocol described above.

Outcomes

Functional exercise capacity: 6MWD

HRQoL: SGRQ

- Symptoms
- Activity
- Impact
- Total

Respiratory function: FEV1 (L)

Identification

Country: Lebanon

Author's name: Rola Tout

Institution: Institut de Physiothérapie, université Saint-Joseph

Email: rolatout@yahoo.com

Address: URAF, rue de Damas, Beyrouth, Lebanon

Notes

Wang 2017

Study characteristics

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p> <p>Subgroup analysis: P_{Imax} less or more than 60 cmH₂O</p>
Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 28/28 • Loss to follow-up or excluded: 0 • Age, mean (SD) in years: 70.8 (4.5) • BMI, mean (SD), kg/m²: 21.31 (2.76) <p>PR</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 27/27 • Loss to follow-up/ excluded: 0 • Age, mean (SD) in years: 70.6 (6.3) • BMI, mean (SD), kg/m²: 22.40 (2.85) <p>Overall</p> <ul style="list-style-type: none"> • N (randomized/analyzed): 55/55 • Loss to follow-up or excluded: 0 • COPD stage (GOLD): moderate to severe <p>Included criteria</p> <ul style="list-style-type: none"> • Age > 40 years • Diagnosis of stable COPD based on the GOLD guidelines; • No participation in any PR program in the previous 2 months • The participants had to be able to understand the investigator's instructions and complete the tests required in the study <p>Excluded criteria</p> <ul style="list-style-type: none"> • Acute or chronic airway diseases other than COPD, cardiovascular disorders (such as acute coronary syndrome), metabolic conditions (such as diabetes or hyperthyroidism), or other health problems that would interfere with exercise performance or the testing procedures
Interventions	<p>Intervention characteristics</p> <p>PR+IMT:</p> <ul style="list-style-type: none"> • PR: consisted of 30 min of cycle ergometer training 3 times/week for 8 weeks. The exercise intensity threshold in lower limbs was calculated as 70% VO₂max during cardiopulmonary exercise training • IMT: conducted 3 times/week for 8 weeks using a Threshold IMT device (model HS730, Philips, Amsterdam, the Netherlands), around 14 min/session at 30% of P_{Imax} <p>PR: this group received only the PR protocol described above</p>
Outcomes	<p>Dyspnea: Borg</p> <ul style="list-style-type: none"> • 6MWD

Wang 2017 (Continued)

- Incremental cycle ergometer test
 - Notes: Borg was measured during peak cycle endurance test
- Dyspnea: mMRC
- Functional exercise capacity:
- 6MWD
 - Wmax: measured through an incremental load of 5 w/min or 10 w/min
- HRQoL: CAT
- HRQoL: SGRQ (Total)
- Respiratory muscle strength: PImax
- Respiratory muscle endurance: MVV
- Laboratory exercise test: VO₂peak
- mL/min
 - mL/kg/min
- Respiratory function: FEV1
- %pred
 - L

Identification

Sponsorship source: this work was supported by Guangzhou Municipal Science and Technology Project (201507020033), Medical Scientific Research Foundation of Guangdong Province (A2016399), Open Project of State Key Laboratory of Respiratory Disease (SKLRD2016OP019), and Clinical Research training program of Southern Medical University (LC2016PY032).

Country: China

Setting: Zhujiang Hospital affiliated to Southern Medical University

Author's name: Xin Chen

Institution: Department of Respiratory Medicine, Zhujiang Hospital, Southern Medical University

Email: chen_xin1020@163.com

Address: 253 Gongye Road, Guangzhou 510282, China

Clinical trial register: NCT02285400

Notes

Adjusted P values were reported.

Authors reported a subgroup analysis for intervention group participants with or without respiratory muscle weakness.

Weiner 1992

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

Weiner 1992 (Continued)

PR+IMT

- N (randomized/analyzed): 12/12
- Loss to follow-up or excluded: 0
- Age, mean (SD in years): 67.2 (9)

PR

- N (randomized/analyzed): 12/12
- Loss to follow-up or excluded: 0
- Age, mean (SD in years): 64.4 (10.3)

Overall

- N (randomized/analyzed): 24/24
- Loss to follow-up or excluded: 0
- COPD stage (GOLD): severe to very severe

Included criteria

- Patients with spirometric evidence of chronic airflow limitation that was not corrected by bronchodilator therapy

Excluded criteria: not reported

Interventions

Intervention characteristics

PR+IMT:

- PR: consisted of
 - 20 min of cycling on cycle ergometer. Participants started cycling with low load that was then gradually increased, about 5% each session, to reach 50% of the initial Wmax.
 - 10 min of rowing
 - 15 min of muscle exercises to strengthen upper and lower extremities and abdominal muscles
- IMT: 15 min/session, 3 times/week for 6 months using the Threshold IMT device. Participants breathed at 15% of their P_Imax for 1 week. The resistance was then increased by 5% to reach 60% of P_Imax at the end of the first month and 80% at the end of the second month.

PR+ (sham IMT): this group received the same training protocol as described above, and IMT was conducted at no load.

Outcomes

Functional exercise capacity:

- 12MWD
- Exercise time (constant cycle ergometer test)
- Notes: endurance work time at 2/3 of Wmax

Respiratory muscle strength: P_Imax (RV)

Respiratory muscle endurance: respiratory muscle endurance pressure (P_{thmax})

- Notes: participantss inspired through a 2-way HansRudolph valve whose inspiratory port was connected to a chamber and plunger to which weights could be added externally. Inspiratory elastic work was then increased by the progressive addition of 25-100 g weights at 2-min intervals. The pressure achieved with the heaviest load (tolerated for at least 60 s) was defined as the peak pressure (P_{thmax})

Respiratory function: FEV1 (%pred)

Identification

Country: Israel

Author's name: Paltiel Weiner

Weiner 1992 (Continued)

Institution: Department of Medicine, Hillel-Yaffe Medical Center, and the Institute for Respiratory Disease

Address: Hadera, Israel

Notes

Weiner 2000
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

PR+IMT

- N (randomized/analyzed): 12/11
- Loss to follow-up or excluded: 1
- Age, mean (SD) in years: 62.5 (8.31)

PR

- N (randomized/analyzed): 5/4
- Loss to follow-up or excluded: 1
- Age, mean (SD) in years: 61.0 (5.4)

Overall

- N (randomized/analyzed): 17/15
- Loss to follow-up or excluded: 2
- COPD stage (GOLD): severe

Included criteria

- Spirometric evidence of chronic airflow limitation
- Diagnosis of moderate-to-severe COPD according to the criteria of the ATS

Excluded criteria: not reported

Interventions

Intervention characteristics

PR+IMT: the training program is the same as described in (Weiner 1992). Participants trained 3 times/week and each session consisted of 1 h of supervised training. When only exercise training was performed, participants trained the whole hour, and when IMT was added, exercise training was cut to 30 min.

PR+ SHAM: participants in this group received the same rehabilitation protocol and IMT was conducted at no load

Outcomes

Respiratory muscle strength: P_{lmax} (RV)

Identification

Country: Israel

Author's name: Paltiel Weiner

Institution: Department of Medicine A, Hillel-Yaffe Medical Center

Weiner 2000 (Continued)

Address: Hadera, Israel

Notes

Weiner 2003
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): 8/8
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 63.1 (8.7)
- Gender (M/F): 6/2

Control/sham

- N (randomized/analyzed): 8/8
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 61.8 (9)
- Gender (M/F): 7/1

Overall

- N (randomized/analyzed): 16/16
- Loss to follow-up or excluded: 0
- Gender (M/F): 13/3
- COPD stage (GOLD): severe

Included criteria

- Spirometric evidence of significant chronic airflow limitation (i.e. FEV1 of < 50% of predicted and FEV1/FVC ratio of 70% of predicted) in whom COPD had been diagnosed, according to the criteria of the ATS

Excluded criteria

- Patients with cardiac disease
- Poor compliance
- A requirement for supplemental oxygen therapy
- CO2 retention

Interventions

Intervention characteristics

IMT: participants trained daily, 1 h/d, 6 times/week for 3 months using Threshold IMT device. They started breathing at a resistance equal to 15% of their P_Imax or P_Emax for 1 week. The resistance then was increased incrementally 5% to 10% each session, to reach 60% of their P_Imax or P_Emax at the end of the first month of training.

Control/sham: participants trained at 7 cmH₂O throughout the trial

Outcomes

Dyspnea: BDI-TDI:

Weiner 2003 (Continued)

- Functional impairment
- Magnitude of task
- Magnitude of effort
- Focal score

Functional exercise capacity: 6MWD

Respiratory muscle strength: P_Imax (RV)

Respiratory muscle endurance: respiratory muscle endurance pressure (P_{thmax})

Identification

Country: Israel

Author's name: Paltiel Weiner

Institution: Department of Medicine A, Hillel Yaffe Medical Center

Email: weiner@hillel-yaffe.health.gov.il

Address: Hadera, Israel 38100

Notes

Because only the IMT group data were reported in the results, and there were differences between the graphs and what was reported numerically, we included both groups' data from the graphs to guarantee consistency.

Weiner 2006
Study characteristics
Methods

Study design: RCT

Study grouping: parallel-group

Participants
Baseline characteristics

IMT

- N (randomized/analyzed): 14/14
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 63.0 (7.66)
- Gender (M/F): 8/6
- BMI, mean (SD), kg/m²: 28.0 (5.72)

Control/sham

- N (randomized/analyzed): 14/14
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 62.7 (7.58)
- Gender (M/F): 8/6
- BMI, mean (SD), kg/m²: 28.0 (5.68)

Overall

- N (randomized/analyzed): 28/28
- Loss to follow-up or excluded: 0
- Gender (M/F): 16/12
- COPD stage (GOLD): severe

Weiner 2006 (Continued)

Included criteria

- Spirometric evidence of severe chronic air flow limitation (FEV 1 < 50% of predicted) and FEV 1/FVC < 70% of predicted

Excluded criteria: not reported

Interventions

Intervention characteristics

IMT: participants trained daily, 1 h/d, 6 d/week for 8 weeks using Powerbreathe (Southam, UK). They started breathing at a resistance equal to 15% of their P_{lmax} for 1 week. The resistance was then increased incrementally (5%–10% each session), to reach 60% of their P_{lmax} at the end of the 1st month. IMT was then continued at 60% of their P_{lmax} adjusted weekly to the new P_{lmax} achieved

Control/sham: participants trained with a resistance of 7 cmH₂O following the same protocol described above

Outcomes

Respiratory muscle strength: P_{lmax} (RV)

Identification

Country: Israel

Author's name: Paltiel Weiner

Institution: Department of Medicine A, Hillel Yaffe Medical Center

Email: weiner@hillel-yaffe.health.gov.il

Address: Hadera 38100 Israel

Notes

Wu 2017
Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Baseline characteristics

IMT (Threshold device)

- N (randomized/analyzed): 19/19
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 59.74 (6.14)
- BMI, mean (SD), kg/m²: 18.40 (2.19)

IMT (resistive device)

- N (randomized/analyzed): 21/21
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 62.24 (7.36)
- BMI, mean (SD), kg/m²: 19.25 (2.17)

Control/sham

- N (randomized/analyzed): 20/29
- Loss to follow-up or excluded: 0

Wu 2017 (Continued)

- Age, mean (SD) in years: 60.30 (6.55)
- BMI, mean (SD), kg/m²: 18.54 (2.58)

Overall

- N (randomized/analyzed): 60/60
- Loss to follow-up or excluded: 0
- COPD stage (GOLD): moderate to severe

Included criteria

- Moderate, severe and very severe COPD (post-bronchodilator FEV₁/FVC < 70% and FEV₁ < 50% of predicted (GOLD B, C and D, respectively)
- Inspiratory muscle weakness (P_Imax < 60 cm H₂O)
- Bronchial dilation test (BDT) negative
- No history of PR

Excluded criteria

- Time from most recent exacerbation > 2 months
- With no medication changes in 1 month prior to enrollment
- Obesity (BMI > 30 m²/kg)
- Severe orthopedic problems having a major impact on ADL
- Previous inclusion in a rehabilitation program (< 1 year)
- Concomitant heart failure and pulmonary vascular diseases
- Diagnosed psychiatric or cognitive disorder
- Progressive neurological or neuromuscular disorder

Interventions

Intervention characteristics

IMT: participants trained twice a day, 15 min/session for 8 weeks using either Threshold IMT (Respironics Inc; Pittsburgh, PA, USA) or Pflex (Respironics Inc, Pittsburgh, PA, USA) devices set at 60% of P_Imax

Control/sham: no intervention received by this group

Outcomes

Dyspnea: BDI-TDI

- Functional impairment
- Magnitude of task
- Magnitude of effort
- Focal score

Functional exercise capacity:

- W_{max} (Incremental cycle ergometer test: measured by increasing the work rate by 10 w/min after one minute of unloaded pedalling)
- Exercise time (Incremental cycle ergometer test)

HRQoL: CRQ

- Dyspnea
- Fatigue
- Emotion
- Mastery
- Total

Respiratory muscle strength: P_Imax (RV)

Laboratory exercise test: VO₂peak (L/min) (Incremental cycle ergometer test)

Wu 2017 (Continued)

Respiratory function: FEV1

- L
- %pred

Identification

Sponsorship source: the study was supported by the Science and Technology Project of Guangdong Province (2017A020211018) and the Guangzhou Healthcare collaborative innovation major project (201604040012) and State's Key Project of Research and Development Plan(2017YFSF11078).

Country: China

Setting: Guangzhou Institute of Respiratory Disease

Author's name: Rongchang Chen

Institution: Guangzhou Institute of Respiratory Disease, State Key Laboratory of Respiratory Disease

Email: chenrc_vip@163.com

Address: Guangzhou, China

Clinical trial register: NCT03101774

Notes

Xu 2018

Study characteristics

Methods

Study design: RCT

Study grouping: parallel-group

Subgroup analysis: PImax: < or > 60 cmH2O

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): 23/23
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 67.49 (6.17)
- BMI, mean (SD), kg/m²: 22.09 (3.37)

Control/sham

- N (randomized/analyzed): 23/23
- Loss to follow-up or excluded: 0
- Age, mean (SD) in years: 69.43 (6.44)
- BMI, mean (SD), kg/m²: 20.86 (4.41)

Overall

- N (randomized/analyzed): 46/46
- Loss to follow-up or excluded: 0
- COPD stage (GOLD): moderate to severe

Included criteria

- Patients with clinically stable COPD

Xu 2018 (Continued)

- Naive to PR and willing to participate

Excluded criteria

- Cognitive disorders
- Organ failure
- Malignant tumors
- Metabolic diseases

Interventions

Intervention characteristics

IMT: participants trained daily, at home, 48 min/d for 8 weeks using Threshold IMT (Respironics, USA). The training load ranged from 30%-45% of P_{lmax}

Control/sham: participants received the same protocol and trained at no load.

Outcomes

Dyspnea: mMRC

Functional exercise capacity: 6MWD

HRQoL:

- SGRQ (Total)
- CAT

Respiratory muscle strength: P_{lmax}

Respiratory function: FEV₁

- L
- %pred

Identification

Sponsorship source: this work was funded unconditionally by Clinical Research training program of Southern Medical University (LC2016PY032), National Key R&D Program of China (2017YFC1310601), The Guangzhou Healthcare Collaborative Innovation Major Project (201604020012), Guangzhou Innovation and Entrepreneurship Education Project of Universities (201709T26), Special Funds for the Cultivation of Guangdong College Students' Scientific and Technological Innovation (PDJHB0101). The sponsors have no any role in design, conduct, data interpretation of the study, and preparation, review or approval of this manuscript

Country: China

Setting: Zhujiang Hospital of Southern Medical University

Author's name: Xin Chen

Institution: Department of Respiratory Medicine, Zhujiang Hospital, Southern Medical University

Email: chen_xin1020@163.com

Address: 253 Gongye Road, Guangzhou 510282, China

Clinical trial register: NCT02326181

Notes

ZhouL 2016

Study characteristics

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

ZhouL 2016 (Continued)

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics IMT <ul style="list-style-type: none"> • N (randomized/analyzed): 22/22 • Loss to follow-up or excluded: 1 • Age, mean (SD) in years: 66.4 (5.5) • Gender (M/F): 17/5 • BMI, mean (SD), kg/m²: 22.5 (2.1) Control/sham <ul style="list-style-type: none"> • N (randomized/analyzed): 22/22 • Loss to follow-up or excluded: 1 • Age, mean (SD) in years: 66.8 (6.2) • Gender (M/F): 16/6 • BMI, mean (SD), kg/m²: 21.8 (2.2) Overall <ul style="list-style-type: none"> • N (randomized/analyzed): 44/44 • Loss to follow-up or excluded: 2 • Gender (M/F): 33/11 • BMI, mean (SD), kg/m² • COPD stage (GOLD): severe to very severe Included criteria <ul style="list-style-type: none"> • Age: 40-80 years • Severe or extremely severe COPD in lung functions • Combined with chronic respiratory failure (under normal inhalation, blood gas analysis PCO₂ ≥ 50 mmHg after 1 h of rest) • No acute occurrence of exacerbating episode in the past 4 weeks Excluded criteria <ul style="list-style-type: none"> • People who smoke > 10 cigarettes/d • Patients with unstable cardiac hemodynamics, such as acute left heart failure, unstable angina • Combined with other respiratory diseases, such as typical bronchiectasis, typical pulmonary fibrosis, sleep apnea, lung tumors, sequelae of tuberculosis (damage to the lung) • Those suffering from neuromuscular diseases or sequelae of severe cerebrovascular accidents
Interventions	Intervention characteristics IMT: participants trained twice a day, without supervision, 30 min/d, 6 d/week for 8 weeks using Threshold IMT device set at 60% of P _I max. This group also received non-invasive positive pressure ventilation Control: this group received long-term oxygen therapy
Outcomes	Dyspnea: mMRC Functional exercise capacity: 6MWD Respiratory muscle strength: P _I max

ZhouL 2016 (Continued)

Identification

Sponsorship source: National Natural Science Foundation of China (81361128004); Public welfare research and Capacity building special fund project (2014A020215033); Guangzhou Medical University Scientific Research Fund (2014c22)

Country: China

Setting: Guangzhou Institute of Respiratory diseases

Comments:

Author's name: Chen Rongchang

Institution: Frist Affiliated Hospital of Guangzhou Medical University

Email: member@wc.rf.org

Address: Guangzhou 510120, China

Clinical trial register: 0192675

Notes

6MWD: six-minute walk distance; **12MWD:** 12-minute walk distance; **ADL:** activities of daily living; **ATS:** American Thoracic Society; **BDI:** Baseline Dyspnea Index; **BMI:** body mass index; **CAT:** Chronic Obstructive Pulmonary Disease Assessment Test; **CCQ:** Clinical COPD Questionnaire; **CI:** confidence interval; **COPD:** chronic obstructive pulmonary disease; **CPET:** cardiopulmonary exercise testing; **CRQ:** Chronic Respiratory Disease Questionnaire; **ERS:** European Respiratory Society; **FEV1:** forced expiratory volume at 1 second; **FRC:** functional residual capacity; **FVC:** forced vital capacity; **GOLD:** Global Initiative for Chronic Obstructive Lung Disease; **HRQoL:** health-related quality of life; **IMT:** inspiratory muscle training; **ISWT:** incremental shuttle walk test; **IVC:** inspiratory vital capacity; **MDP:** Multidimensional Dyspnea Profile; **MIP:** maximal inspiratory pressure; **mMRC:** Modified Medical Research Council; **MVV:** maximal voluntary ventilation; **PEmax:** maximal expiratory pressure; **PFT:** pulmonary function test; **PImax:** maximal inspiratory pressure; **PR:** pulmonary rehabilitation; **P_{thmax}:** respiratory muscle endurance pressure; **RCT:** randomized controlled trial; **RV:** residual volume; **SD:** standard deviation; **SE:** standard error; **SEM:** standard error of the mean; **SGRQ:** St George's Respiratory Questionnaire; **SpO₂:** Peripheral oxygen saturation; **SWT:** shuttle walk test; **TDI:** Transition Dyspnea Index; **VO_{2max}:** maximal oxygen consumption; **Wmax:** maximum exercise workload

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Ahmad 2013	Ineligible intervention
Aldrich 1985	Ineligible study design
Anand 2013	Ineligible intervention
Baines 2005	Ineligible comparator
Basso Vanelli 2016	Ineligible comparator
Battaglia 2009	Ineligible intervention
Belman 1994	Ineligible comparator
Bgin 1991	Ineligible study design
Bissett 2016	Ineligible patient population

Study	Reason for exclusion
Bjerre Jepsen 1981	Ineligible intervention
Cader 2010	Ineligible patient population
Chen 1985	Ineligible intervention
Daynes 2018	Ineligible study design
de Andrade 2005	Ineligible study design
de Lucas Ramos 1998	Ineligible study design
Di Mambro 2007	Ineligible study design
DRKS00005637	Ineligible comparator
DRKS00006021	Cancelled Clinical trial
Elbouhy 2014	Ineligible patient population
Elmorsi 2016	Ineligible study design
Enright 2005	Ineligible comparator
Garcia 2008	Ineligible study design
Goldstein 1989	Ineligible intervention
Gregg 1989	Ineligible study design
Guyatt 1992	Ineligible intervention
Hart 2000	Ineligible patient population
Heydari 2015	Ineligible comparator
Hopp 1996	Ineligible study design
Ibakordor 2013	Ineligible comparator
Ionescu 2005	Non-RCT
Izumizaki 2008	Ineligible study design
Johnson 1996	Ineligible study design
Kivastik 2015	Ineligible study design
Koch 2020	Ineligible study design
Kolesnikova 2016	Ineligible intervention
Levine 1986	Ineligible comparator
Liao 2015	Ineligible intervention

Study	Reason for exclusion
Lin 2012	Ineligible intervention
Lisboa 1994	Ineligible patient population
Lisboa 1995a	Ineligible study design
Lisboa 1995b	Ineligible patient population
Lisboa 1998	Ineligible study design
Madariaga 2007	Ineligible comparison
Madsen 1985	Ineligible comparator
Martin 2006	Ineligible intervention
McKeon 1986	Ineligible patient population
Meshcheriakova 2006	Ineligible intervention
Minoguchi 2002	Ineligible study design
NCT01218295	Ineligible study design
NCT01556139	Ineligible patient population
NCT01747694	Ineligible intervention
NCT01945398	Ineligible study design
NCT01956565	Ineligible study design
NCT02186340	Ineligible study design
NCT02278523	Ineligible comparator
NCT02579200	Ineligible patient population
NCT02914093	Ineligible comparator
NCT02935166	Ineligible study design
NCT03186092	Ineligible patient population
NCT03500042	Ineligible outcomes
NCT03739879	Ineligible study design
NCT03844711	Ineligible comparator
NCT03880630	Ineligible study design
NCT04084405	Ineligible comparator
NCT04117399	Ineligible outcomes

Study	Reason for exclusion
NCT04460261	Ineligible study design
Neves 2014a	Ineligible intervention
Neves 2014b	Ineligible study design
Nield 2007	Ineligible intervention
Nosedá 1987	Ineligible comparator
O'Connor 2019	Ineligible study design
Okura 2019	Ineligible study design
Okura 2020	Ineligible study design
PACTR201703002095224	Ineligible comparator
Padula 2001	Ineligible patient population
Perez 2010	Ineligible intervention
Pescaru 2016	Ineligible study design
Quintero 1999	Ineligible study design
Richardson 1989	Ineligible study design
Rocha 2015	Ineligible intervention
Sassoon 1992	Ineligible intervention
Serón 2005	Ineligible patient population
Shahin 2008	Ineligible study design
Shioya 2007	Ineligible intervention
Similowski 1994	Ineligible study design
Sivashanmugam 2019	Ineligible comparator
Soicher 1998	Ineligible study design
Sonne 1982	Ineligible study design
Sudo 1997	Ineligible intervention
Sugiyama 2010	Ineligible study design
Sun 2003	Ineligible intervention
TCTR20191009004	Ineligible intervention
UMIN000030937	Ineligible intervention

Study	Reason for exclusion
Van't Hul 2006	Ineligible intervention
Villafranca 1998	Ineligible study design
Wada 2016	Ineligible intervention
Wu 2006	Ineligible intervention
Xi 2015	Ineligible intervention
Yamaguti 2012	Ineligible intervention
Yan 1996	Ineligible intervention
Yang 2005	Ineligible intervention
Zhang 2008	Ineligible study design

Characteristics of studies awaiting classification *[ordered by study ID]*

Barter 1987

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Bustamante 1997

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Cassidy 2009

Methods	Study design: RCT
	Study grouping: parallel-group

Cassidy 2009 (Continued)

Participants

Baseline characteristics

IMT

- N (randomized/analyzed): 14/14
- Loss to follow-up or excluded: 0
- Gender (M/F): 7/7

Control/sham

- N (randomized/analyzed): 14/14
- Loss to follow-up or excluded: 0
- Gender (M/F): 5/9

Overall

- N (randomized/analyzed): 28/28
- Loss to follow-up or excluded: 0
- Gender (M/F): 12/16
- COPD stage: moderate to severe

Included criteria: patients with COPD, following an acute exacerbation

Excluded criteria: not reported

Interventions

Intervention characteristics
IMT: participants underwent 8 weeks of unsupervised IMT using a Threshold IMT device

Control/sham: participants received sham IMT

Outcomes

 Respiratory muscle strength: P_{lmax}

Functional exercise capacity: 6MWD

HRQoL: CRQ

Notes

Country: Ireland

Setting: home-based training

Author's name: C.Cassidy

Institution: Respiratory Assessment Unit, CREST Directorate

Address: St. James's Hospital, Dublin 8, Ireland

Cejudo 1998

Methods

Unfound

Participants

Unfound

Interventions

Unfound

Outcomes

Unfound

Notes

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)
134

Chen 2017

Methods	Study design: RCT Study grouping: parallel-group Clinical trial register: NCT02200549
Participants	Baseline characteristics PR+IMT <ul style="list-style-type: none"> • N (randomized/analyzed): 24/not reported PR <ul style="list-style-type: none"> • N (randomized/analyzed): 25/not reported Overall <ul style="list-style-type: none"> • N (randomized/analyzed): 49 not reported Included criteria: not reported Excluded criteria: not reported
Interventions	Intervention characteristics PR+IMT: <ul style="list-style-type: none"> • PR: 8 weeks of cycle ergometer training. • IMT: 8 weeks of IMT PR: training with a cycle ergometer for 8 weeks
Outcomes	Dyspnea Functional exercise capacity HRQoL: CRQ Respiratory muscle strength
Notes	Country: China Author's name: X.Chen Institution: Zhujiang Hospital- Southern Medical University, Department of Respiratory Medicine Address: Guangzhou, China

Croitoru 2013

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics PR+IMT

Croitoru 2013 (Continued)

- N (randomized/analyzed): 14/14
- Loss to follow-up or excluded: 0
- Age mean (SD) in years: 63.4 (8)

PR

- N (randomized/analyzed): 14/14
- Loss to follow-up or excluded: 0
- Age mean (SD) in years: 60.3 (11)

Overall

- N (randomized/analyzed): 28/28
- Loss to follow-up or excluded: 0
- COPD stage: moderate to severe

Included criteria: not reported

Excluded criteria: not reported

Interventions	<p>Intervention characteristics</p> <p>PR+IMT:</p> <ul style="list-style-type: none"> • PR: outpatient, 8 weeks, lower and upper limbs training, education, and psychological support • IMT: daily, at home, with Threshold device, 30 min/d <p>PR: participants received the same PR protocol described above.</p>
Outcomes	<p>Functional exercise capacity: 6MWD</p> <p>HRQoL: SGRQ</p> <p>Respiratory muscle strength: P_{Imax}</p>
Notes	

Del Castillo Otero 1998

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Di Marzo 2000

Methods	Unfound
Participants	Unfound

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)
136

Di Marzo 2000 *(Continued)*

Interventions	Unfound
Outcomes	Unfound
Notes	

Di Marzo 2002

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Downes Vogel 2002

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Eastwood 2005

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Gething 2001

Methods	Unfound
Participants	Unfound

Gething 2001 (Continued)

Interventions	Unfound
Outcomes	Unfound
Notes	

Göhl 2006

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

IRCT201104266299N1

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics Overall <ul style="list-style-type: none"> N (randomized/analyzed): 30 Included criteria <ul style="list-style-type: none"> Patients with mild to very severe COPD Having an established treatment plan BMI < 35 Absence of other diseases such as neurological disorders, musculoskeletal system, peripheral vascular disease, and independence to the long-term oxygen therapy Exclusion criteria <ul style="list-style-type: none"> Changes in the treatment plan during the study.
Interventions	Intervention characteristics IMT: participants trained at home daily for 21 min, 6 d/week for 8 weeks Control/sham: no intervention
Outcomes	Dyspnea HRQoL: SGRQ
Notes	

IRCT20180205038633N1

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>PR+IMT</p> <ul style="list-style-type: none"> N (randomized/analyzed): 15/unknown <p>PR</p> <ul style="list-style-type: none"> N (randomized/analyzed): 15/unknown <p>IMT</p> <ul style="list-style-type: none"> N (randomized/analyzed): 16/unknown <p>Control/sham:</p> <ul style="list-style-type: none"> N (randomized/analyzed): 15/unknown <p>Overall</p> <ul style="list-style-type: none"> N (randomized/analyzed): 61/unknown COPD stage: moderate to severe <p>Inclusion/exclusion criteria</p> <ul style="list-style-type: none"> Willing to participate in the study Grade 2 or 3 COPD based on GOLD criteria Age 40-70 years; Lack of other pulmonary diseases Lack of severe limb limbs Lack of pulmonary surgery in the last 12 months Recent fracture failure at the level of ribs (6 months) No history of psychotropic disease and related drugs and alcohol and psychotropic substances
Interventions	<p>Intervention characteristics</p> <p>IMT: the intensity of training was 40%-60% (S-Index 40%-60%), 2 d/week, and 5 sessions of dermal muscle training with repeat 50 (5 Repeat 10) for about 15 min/session.</p> <p>PR: aerobic exercise method: with a treadmill and foot pedometer, 2 times a week, with 40%-60% heart rate reserve for 40 min/session</p> <p>PR+IMT: in each session, breathing exercises were performed first and then aerobic exercises were performed on the lower extremities</p> <p>Control: this group did not take any special intervention other than the usual treatments (control group)</p>
Outcomes	<p>Dyspnea: mMRC</p> <p>Respiratory muscle strength: P_{lmax}</p> <p>Functional exercise capacity: 6MWD</p> <p>Respiratory function: FEV₁</p>
Notes	

ISRCTN19258620

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> 80 non-hypercapnic patients with moderate (FEV < 40%) recruited from consultant hospital and community chest clinics All patients will be receiving optimum medical management and will have been stable for at least 4 weeks prior to their initial assessment <p>Exclusion criteria</p> <ul style="list-style-type: none"> Hypercapnia (PaCO₂ > 45 mmHg) Any patient who is unsuitable for magnetic stimulation (pacemakers, artificial heart valves, metal prosthesis)
Interventions	<p>IMT: using Powerbreathe device</p>
Outcomes	<p>Dyspnea: Borg</p> <p>Functional exercise capacity: SWT</p> <p>Respiratory muscle strength: P_{lmax}</p> <p>Respiratory muscle endurance pressure: P_{thmax}</p> <p>Respiratory muscle endurance time: T_{lim}</p>
Notes	

Jones 1985

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>Overall</p> <ul style="list-style-type: none"> N (randomized/analyzed): 21 <p>Inclusion criteria: FEV1 < 1.2 L</p>
Interventions	<p>Intervention characteristics</p> <p>IMT: training with resistive device for 10 weeks</p> <p>Control/sham: placebo training group</p>
Outcomes	<p>Functional exercise capacity: W_{max}</p> <p>Functional exercise capacity: 12MWD</p>

Jones 1985 (Continued)

Notes

Koppers 2004

Methods	Unfound
---------	---------

Participants	Unfound
--------------	---------

Interventions	Unfound
---------------	---------

Outcomes	Unfound
----------	---------

Notes

Liu 1989

Methods	Unfound
---------	---------

Participants	Unfound
--------------	---------

Interventions	Unfound
---------------	---------

Outcomes	Unfound
----------	---------

Notes

Manuel Vargas 1995

Methods	Study design: RCT Study grouping: parallel-group
---------	---

Participants	Baseline characteristics Overall <ul style="list-style-type: none"> N (randomized/analyzed): 23
--------------	--

Interventions	Intervention characteristics IMT: participants trained with a Threshold device at 30% of P _{Imax} Control/sham: no training was provided
---------------	--

Outcomes	Functional exercise capacity
----------	------------------------------

Notes

Mendoza 2007

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	Unfound

Meshcherykova 2018

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics IMT <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/17 • Age mean (SD) in years: 62.5 (7.5) Control/sham <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/11 • Age mean (SD) in years: 63.5 (9.8) Overall <ul style="list-style-type: none"> • N (randomized/analyzed): not reported/ 28 • COPD stage: severe Included criteria: patients with severe COPD
Interventions	Intervention characteristics IMT: participants received IMT for 3 months Control/sham: training with no load
Outcomes	Dyspnea: BDI-TDI Functional exercise capacity: 6MWD Respiratory function: FEV1 Respiratory function: RV
Notes	

NCT01056081

Methods	Study design: RCT Study grouping: parallel-group
---------	---

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

142

NCT01056081 (Continued)

Participants

Baseline characteristics

Overall:

- N (randomized/analyzed): 19/unknown

Inclusion criteria

- Patients with a clinical and spirometric diagnosis of moderate to severe COPD according to GOLD
- In a stable condition (without exacerbations or infections for at least a month)
- Had to be former smokers (> 6 months without smoking)
- Patients referred by a physician to the pulmonary rehabilitation program

Exclusion criteria

- Patients with a known history of asthma, or severe and/or unstable heart disease or any other pathological condition that could impair their physical activities

Interventions

Intervention characteristics

PR+IMT

- PR: no information
- IMT: participants trained with a Threshold inspiratory muscle trainer (Respironics HealthScan, Inc, Cedar Grove, New York, USA). The participants performed the IMT training in a seated position, with the upper limbs supported. The total duration of the respiratory training was 30 min, with sequences of 3 min of training followed by pauses of 2 min. The initial load was equivalent to 30% of the individual's MIP. This load was progressively increased over the first 4 weeks, according to the participant's tolerance, to reach 60% of the MIP. This level was then maintained until the end of the training.

PR: this group received only PR

Outcomes

Dyspnea

Functional exercise capacity: exercise time

Respiratory muscle strength: P_{lmax}

Notes

NCT01903772

Methods

Study design: RCT

Study grouping: parallel-group

Participants

Inclusion Criteria:

- Patients with COPD
- Eligible to participate in an inpatient pulmonary rehabilitation program of 3 weeks
- P_{lmax} <60 cmH₂O or <50% of the predicted normal value

Exclusion Criteria:

- Major comorbidities preventing successful participation in an 8-week exercise training intervention
- use of non-invasive ventilation

NCT01903772 (Continued)

Interventions	<p>Intervention characteristics</p> <p>IMT: Inspiratory Muscle Training Three times daily inspiratory muscle training (2x30 breaths) at an intensity of >50% P_{Imax}</p> <p>Sham IMT: Twice daily inspiratory muscle training (3x30 breaths) at an intensity of 5 centimeters of water (H₂O)</p>
Outcomes	<p>Dyspnea</p> <p>Functional exercise capacity</p> <p>Respiratory muscle strength: P_{Imax}</p> <p>Inspiratory muscle endurance capacity</p>
Notes	<p>It is unclear whether the study was completed, and no contact details were found.</p>

NCT02392715

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> • COPD demonstrated by spirometry using GOLD criteria • Patient referred by a pneumologist to the ambulatory PR program at the Riviera-Chablais Hospital, Monthey • Patient with maximal inspiratory pressure < 60 cmH₂O • Patient > 40 years <p>Exclusion criteria</p> <ul style="list-style-type: none"> • Orthopaedic or neurological troubles that could slant the 6MWD
Interventions	<p>Intervention characteristics</p> <p>PR+IMT:</p> <ul style="list-style-type: none"> • PR: general exercise training • IMT: participants trained with a Threshold device 3 times/week with a total of 36 sessions. The training load was increased from 15%-60% of P_{Imax}. <p>PR: (+sham IMT): this group received general exercise training with a sham IMT set at 5 cmH₂O.</p>
Outcomes	<p>Functional exercise capacity: 6MWD</p> <p>HRQoL: SGRQ</p> <p>Inspiratory muscle strength (P_{Imax})</p>
Notes	

NCT02673242

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> • COPD grade 3-4 • Inhabitant of Hedmark (Løten, Våler, Åsnes, Hamar, Elverum) <p>Exclusion criteria</p> <ul style="list-style-type: none"> • Undergoing exercise-based physical therapy treatment • Not able to do IMT physically or mentally
Interventions	<p>Intervention characteristics</p> <p>IMT: using a Threshold device for 6 weeks</p> <p>Control/sham: participants received sham training or another intervention</p>
Outcomes	<p>Dyspnea: mMRC</p> <p>Functional exercise capacity: 6MWD</p> <p>HRQoL: CAT</p>
Notes	

NCT03080662

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> • To be eligible for the PR program participants must have stable COPD (at least 4 weeks), inspiratory muscle weakness (P_Imax < 70%) and pulmonary hyperinflation (TLC > 120%). Patients that have signed informed consent <p>Exclusion criteria</p> <ul style="list-style-type: none"> • Hospitalization within the previous 14 days • Current participation in a rehabilitation program • Locomotor or neurological condition or disability limiting the ability to perform exercise • Lung transplantation or lung volume reduction surgery foreseen within 1 month after discharge
Interventions	<p>Intervention characteristics</p> <p>IMT: training with ORYGEN DUAL Sham Valve for 5 weeks</p> <p>Control/sham: training with the same protocol with no load</p>
Outcomes	

NCT03080662 (Continued)

Notes

NCT03438019

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Baseline characteristics</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> • Male or female aged > 40 years with a confirmed diagnosis of COPD by a pulmonologist (presence of risk factors and airflow obstruction) • Evidence of inspiratory muscle weakness as defined by the latest ATS/ERS statement on respiratory muscle training <p>Exclusion criteria</p> <ul style="list-style-type: none"> • Refusal to participate in the study • Patients actively undergoing PR • Inability to perform the required manoeuvres (i.e. patients with a cerebrovascular accident or tracheostomy) • Patients not in their stable state (i.e. having an acute exacerbation or within 4 weeks of having one) or the presence of important comorbidities that may confound the interpretation of TIRE (test of incremental respiratory endurance) measures (i.e. decompensated heart failure, diaphragmatic paralysis, prior lung surgery, active cancer treatment, etc.).
Interventions	<p>Intervention characteristics</p> <p>The TIRE IMT group: will receive a tablet with the TIRE software installed and a PrO2 device through which they will train. Training consists of 6 levels (A-F) with 6 inspirations at each level for a total of 36 breaths. Recovery times between breaths range from 40-5 seconds as the participant advances each level. TIRE data will be stored in the tablet for subsequent interrogation and data retrieval.</p> <p>The standard IMT group: will receive a Threshold Inspiratory Muscle Trainer. This device incorporates a flow-independent one-way valve to ensure consistent resistance and features an adjustable specific pressure setting to be set based on MIP values of each participant. Participants will be instructed to perform up to 36 breaths daily. To compare with TIRE training, we will ask participants to perform this within a 30-min session.</p> <p>The sham IMT group: will also receive a Threshold device and undergo the exact protocol of group 2 but with minimal resistance applied (7 cm H2O, the lowest in the device).</p>
Outcomes	<p>Dyspnea: mMRC</p> <p>Functional exercise capacity: 6MWD</p> <p>HRQoL:</p> <ul style="list-style-type: none"> • SGRQ • CAT <p>Respiratory muscle strength: P_{lmax}</p>
Notes	

NCT03790410

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics Inclusion criteria <ul style="list-style-type: none"> • Stable COPD • Do not take part in other treatment • Be able to learn the usage of inspiratory muscle trainer Exclusion criteria <ul style="list-style-type: none"> • Acute exacerbation • Balance problems with a neurological cause • The patient unable to cooperate • Participation in pulmonary rehabilitation 3 months before the study • IMT use in the 3 months before the study
Interventions	Intervention characteristics PR+IMT: PR consists of breathing exercises, bicycle ergometer conditionings, relaxations, and strength and endurance training. IMT consists of strengthening exercises on the diaphragm muscle. PR: This group undertook only PR.
Outcomes	Respiratory muscle strength: P _{lmax}
Notes	

Newall 1998

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Newall 2000

Methods	Unfound
Participants	Unfound
Interventions	Unfound

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

Newall 2000 *(Continued)*

Outcomes	Unfound
Notes	Unfound

NTR2990

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Pertuze 1994

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Ramirez Sarmiento 2000

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Reidi 2005

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics Overall

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

148

Reidi 2005 (Continued)

- N (randomized/analyzed): 18/unknown
- Age mean (SD) in years: 65.1 (6.6)

Inclusion criteria: FEV1 < 60%

Interventions	<p>Intervention characteristics</p> <p>IMT: the training was performed through the Threshold set at 30 of P_{Imax}. The program consisted of 4 weeks, with 3 weekly sessions where both groups were reassessed weekly</p> <p>Control/sham: the training consisted of performing the threshold with or without the natural resistance of the equipment < 7 cmH₂O</p>
Outcomes	Respiratory muscle strength: P _{Imax}
Notes	

Valderramas 2009

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Vargas 1995

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	
Interventions	<p>Intervention characteristics</p> <p>IMT: participants used an inexpensive pressure threshold load valve constructed according to the Appropriate Technology principles of the WHO, adjusted at 30% of MIP for 3 months</p> <p>Control/sham: this group did not receive any intervention.</p>
Outcomes	Respiratory muscle strength: P _{Imax}
Notes	

Vargas 1998

Methods	Unfound
---------	---------

Vargas 1998 (Continued)

Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Wang 2004

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics Overall <ul style="list-style-type: none"> N (randomized/analyzed): 64
Interventions	Intervention characteristics IMT: twice a day for 6 months Control/sham: no intervention was received
Outcomes	Respiratory function: FEV1 (%pred)
Notes	

Wanke 1994

Methods	Study design: RCT Study grouping: parallel-group
Participants	Baseline characteristics PR+IMT <ul style="list-style-type: none"> N (randomized/analyzed): 21/unknown PR <ul style="list-style-type: none"> N (randomized/analyzed): 21/unknown
Interventions	Intervention characteristics PR+IMT: <ul style="list-style-type: none"> PR: cycle ergometer training for 8 weeks IMT: for 8 weeks PR: cycle ergometer training alone
Outcomes	Functional exercise capacity

Wanke 1994 *(Continued)*

Notes

Weiner 2006a

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

Wolstenholme 1998

Methods	Unfound
Participants	Unfound
Interventions	Unfound
Outcomes	Unfound
Notes	

6MWD: six-minute walk distance; **12MWD:** 12-minute walk distance; **ATS:** American Thoracic Society; **BDI:** Baseline Dyspnea Index; **BMI:** body mass index; **CAT:** COPD Assessment Test ; **COPD:** chronic obstructive pulmonary disease; **CRQ:** Chronic Respiratory Disease Questionnaire; **ERS:** European Respiratory Society; **FEV1:** forced expiratory volume at 1 second; **GOLD:** Global Initiative for Chronic Obstructive Lung Disease; **HRQoL:** health-related quality of life; **IMT:** inspiratory muscle training; **MIP:** maximal inspiratory pressure; **mMRC:** Modified Medical Research Council; **PImax:** maximal inspiratory pressure; PR: pulmonary rehabilitation; **RCT:** randomized controlled trial; **RV:** residual volume; **SGRQ:** St George's Respiratory Questionnaire; **SWT:** shuttle walk test; **TDI:** Transition Dyspnea Index; **TIRE:** test of incremental respiratory endurance; **WHO:** World Health Organization; **Wmax:** maximum exercise workload

Characteristics of ongoing studies *[ordered by study ID]*
CTRI/2020/11/029226

Study name	Effects of inspiratory muscle training vs autogenic drainage in hospitalised COPD patients
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> • Diagnosed case of COPD from 4-10 years • Patients of both genders aged between 50-70 years • Smokers and non-smokers • Clinically stable patients Exclusion criteria

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) 151
 (Review)

CTRI/2020/11/029226 (Continued)

- Resting SBP > 200 mm Hg or DBP > 110 mm Hg
- Pre-hypertension and hypotension
- Diagnosed cases of orthopnea
- Diagnosed cases of chronic kidney disease
- Injured musculoskeletal or fractured since 3 months.

Interventions	IMT: conducted for 30 min with a total of 10 sessions Control: this group did not receive any intervention
Outcomes	HRQoL: SGRQ Pulmonary function tests
Starting date	20 November 2020
Contact information	Sponsorship source: KLE Academy of Higher Education and Research Country: India Author's name: Varun Naik Institution: Physiotherapy Department Nehru Nagar Belgaum Belgaum Email: drvarunnaik@gmail.com Address: KARNATAKA 590010 Clinical trial register: CTRI/2020/11/029226
Notes	

CTRI/2021/05/033469

Study name	To test the efficacy of inspiratory muscle training device Airofit in reducing breathlessness in COPD patients
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> • All patients with spirometry-proven, stable COPD (GOLD stage ≥ 2) screened for inclusion • Patients with MIP < 100% of predicted • Patients aged 35-65 years Exclusion criteria <ul style="list-style-type: none"> • Patients without access to a suitable smartphone or tablet for the duration of intervention • Patients with an inability to read and understand written and verbal instructions in English • Patients with a history of hospitalization during the previous 4 weeks • Patients with severe orthopedic problems during the previous 4 weeks • Patients with diagnosed psychiatric or cognitive disorders • Patients with a progressive neurological or neuromuscular disorder • Patients on the waiting list for lung transplantation • Patients with previous inclusion in a rehabilitation program < 1 year • Patients with previous experience with IMT

CTRI/2021/05/033469 (Continued)

- Patients with underlying bronchiectasis, innate lymphoid cell, post-COVID and post-tuberculosis sequelae
- All COPD patients with active COVID status or within 15 days of post-COVID recovery

Interventions	<p>Intervention characteristics</p> <p>IMT (Airofit device group): the Airofit is a new IMT device, which combines a calibrated inspiratory flow resistance with a pressure transducer and mobile device app. As the user inhales through the device, a pressure load is created; the magnitude of the load is proportional to the rate of air flow, as well as to the flow resistance properties of the load setting. The former has been identified as a limitation of flow-resistive IMT. To overcome the flow-dependency of the training load, the Airofit uses a pressure transducer to communicate with bespoke software, in real time. The measurement of pressure provides the user with instantaneous visual feedback of the pressure created by their inspiratory muscles, i.e. their training effort. The software also provides the user with a personalized visual training intensity target (50% of MIP). A total of 19 patients with a spirometry-proven stable COPD will be enrolled in the experimental arm.</p> <p>IMT (Powerbreathe device group): 19 participants will train at a load of 50% of PImax.</p> <p>Sham IMT: the participants will use control device 'Breathing pacer'</p>
Outcomes	<p>Inspiratory muscle strength (PImax)</p> <p>Adverse events</p>
Starting date	1 June 2021
Contact information	<p>Sponsorship source: Airofit AS Copenhagen, Denmark</p> <p>Country: India</p> <p>Author's name: Atul Deshmukh</p> <p>Institution: Padmashree Dr. D Y Patil University Navi Mumbai</p> <p>Email: atul.deshmukh@dypatil.edu</p> <p>Address: Centre for Interdisciplinary Research, Ground floor, Central Research Facility, Near Simulation Laboratory Padmashree Dr. D Y Patil University Campus Sector 7 Nerul Navi Mumbai, MAHARASHTRA, 400706, India</p>
Notes	

CTRI201712010952

Study name	Effect of breathing exercises to improve the strength of respiratory muscles in COPD population
Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Inclusion criteria: diagnosed COPD patients</p> <p>Exclusion criteria: women with COPD, those unable to comprehend</p>
Interventions	<p>Intervention : IMT+PR</p> <p>Control: PR</p>

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

153

CTRI201712010952 (Continued)

Outcomes	6MWD Respiratory muscle strength (P _I max) CRQ
Starting date	10 April 2017
Contact information	veenakiran_nambiar@yahoo.co.in
Notes	

De Souza 2019

Study name	Does inspiratory muscle training (IMT) reduce depression in patients with COPD?
Methods	Study design: RCT Study grouping: parallel-group
Participants	Patients with COPD presenting with inspiratory muscle weakness
Interventions	Intervention characteristics IMT: participants trained at 50% of P _I max, 2 sessions/d of 30 breaths for 8 weeks, with a weekly face-to-face session Control/sham: participants trained at 10% of P _I max with a similar protocol
Outcomes	Dyspnea: mMRC Functional exercise capacity: 6MWD Respiratory muscle strength: P _I max
Starting date	
Contact information	souzayr@gmail.com
Notes	

Formiga 2020

Study name	Novel versus traditional inspiratory muscle training regimens as home-based, stand-alone therapies in COPD: protocol for a randomized controlled trial
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> • Patients > 40 years with a clinical and functional diagnosis of COPD according to GOLD guidelines – stages 1-4

Formiga 2020 (Continued)

- Evidence of inspiratory muscle weakness, defined as a MIP \leq 80 cmH₂O and a SMIP \leq 427 pressure-time units
- The ability to operate a computer, tablet or smartphone independently and follow the training instructions
- Clinical stability with no history of infections or exacerbation of respiratory symptoms for at least 2 months prior to study enrollment
- Non-participation in exercise programs in the past 12 months

Exclusion criteria

- History of lung surgery, lung cancer, as well as individuals with any diagnosed cognitive (i.e. Mini Mental State Examination score $<$ 24), orthopedic, neurological or neuromuscular disorders that might prevent them from appropriately performing the required physical tests and/or completing the study questionnaires
- Patients will not be excluded based upon their current bronchodilator regimen. If they experience acute exacerbations or respiratory infections during the training period, they will be examined by a pneumologist who will decide whether the participant should continue with the training or not.

Interventions	<p>Intervention characteristics</p> <p>All groups will train once a day for 8 weeks</p> <p>TIRE: participants will use an on-screen training template set at 50% of their MIP and SMIP. The training will consist of 6 levels (A-F) with 6 inspirations per level for up to 36 efforts per session. Pre-set recovery times between breaths: 60 seconds at level A to 50, 40, 30, 20 and 10 seconds at levels B to F, respectively</p> <p>Threshold IMT: participants will train using a one-way spring-loaded valve set at 50% of their MIP. The training will consist of 36 inspirations performed using the device within a 30-min period.</p> <p>Sham-IMT: participants will use a one-way spring-loaded valve set to its minimal resistance (-9 cmH₂O). The training will consist of 36 inspirations performed using the device within a 30-min period.</p>
Outcomes	<p>Dyspnea: mMRC</p> <p>Respiratory muscle strength: P_Imax</p> <p>HRQoL: CAT</p> <p>Functional exercise capacity: 6MWD</p> <p>Respiratory muscle strength (P_Imax)</p> <p>Pulmonary function test</p>
Starting date	1 May 2021
Contact information	dosbaba.filip@fnbrno.cz
Notes	Clinical trial register: NCT04415788

JPRN-UMIN00039893

Study name	The effectiveness of inspiratory muscle training in patients with chronic obstructive pulmonary disease
Methods	Study design: RCT

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

155

JPRN-UMIN000039893 (Continued)

Study grouping: parallel-group

Participants	<p>Inclusion criteria</p> <ul style="list-style-type: none"> • Age minimum: 20 years • Age maximum: not applicable • Gender: male <p>Exclusion criteria</p> <ul style="list-style-type: none"> • Participants have no severe and/or unstable cardiac disease, orthopedic disease or mental disorder
Interventions	<p>Intervention characteristics</p> <p>PR+IMT:</p> <ul style="list-style-type: none"> • PR: • IMT: 30%-50% of P_{lmax} intensity, 30 breaths/session, 2 sessions/d, every day for 3 months <p>PR +sham IMT: this group will undergo the same rehabilitation program with an IMT load set at 10% of P_{lmax}</p>
Outcomes	<p>Functional exercise capacity: 6MWD</p> <p>HRQoL: CAT</p> <p>Inspiratory muscle strength (P_{lmax})</p>
Starting date	01/11/2016
Contact information	<p>Sponsorship source: Akita University Graduate School of Health Sciences</p> <p>Country: Japan</p> <p>Author's name: Takanobu Shioya</p> <p>Institution: Akita University Graduate School of Health Sciences Department of Physical Therapy</p> <p>Email: shioya@hos.akita-u.ac.jp</p> <p>Address: 1-1-1, Hondo, Akita Japan</p> <p>Clinical trial register: JPRN-UMIN000039893</p>
Notes	

JPRN-UMIN000043099

Study name	Effect of inspiratory muscle training on diaphragm and exercise tolerance in patients with COPD
Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>Inclusion criteria</p> <ul style="list-style-type: none"> • Age minimum: 65 years • Age maximum: 85 years • Gender: male and female

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

156

JPRN-UMIN000043099 (Continued)

Exclusion criteria

- The exclusion criteria include unstable medical conditions that cause or contribute to breathlessness (i.e. metabolic, cardiovascular, or other respiratory diseases) or any other disorders that interfere with exercise testing, such as neuromuscular diseases or musculoskeletal problems

Interventions	Intervention characteristics IMT: IMT will be performed for 12 weeks Control: this group will not receive any intervention
Outcomes	Exercise capacity
Starting date	22 January 2021
Contact information	Sponsorship source: Kindai University Hospital Country: Japan Author's name: Masashi Shiraishi Institution: Kindai University Hospital Department of Rehabilitation Email: masashi-shiraishi@med.kindai.ac.jp Address: 377-2 Onohigashi, Osakasayama-city 589-8511 Japan Clinical trial register: JPRN-UMIN000043099
Notes	

NCT04120142

Study name	Effect of inspiratory muscle training during PR on dyspnoea and exercise tolerance in COPD patients
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> • COPD diagnosed by pulmonary function testing • Clinically stable • Absence of other obstructive diseases • Signed written consent Exclusion criteria <ul style="list-style-type: none"> • Were previous pneumonectomy or lobectomy in the past 6 months • Spontaneous risk of pneumothorax or rib fracture • Incapacity to follow a standard rehabilitation programme (locomotor deficits, acute cardiac failure and acute exacerbation of COPD at the beginning of the program) • The absence of written informed consent
Interventions	Intervention characteristics PR+IMT: IMT + aerobic exercise

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

157

NCT04120142 (Continued)

PR: aerobic exercise alone

Outcomes	Dyspnea Respiratory muscle strength (P _I max)
Starting date	1 February 2019
Contact information	
Notes	

NCT04201522

Study name	The effect of respiratory training on exercise tolerance in COPD (ERTET)
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> Age ≥ 40 years Chronic airflow obstruction: FEV₁/FVC < 0.7, FEV₁ of 30%-80%pred, after bronchodilation Exclusion criteria <ul style="list-style-type: none"> Inability to perform a cycling exercise Diagnosed of ≥ 1 comorbidities that may limit exercise tolerance: cardiovascular, metabolic, endocrine, gastrointestinal, renal, neurological or rheumatological disease Recent COPD exacerbation (< 3 months) Recent cancer A daily dose of Prednisone > 10 mg Hypoxemia at rest or during exercise: PaO₂ < 60 mmHg or SpO₂ ≤ 88% BMI > 30 kg/m² Pregnancy Skinfold at intercostal or vastus lateralis muscle > 1.5 cm
Interventions	Intervention characteristics IMT: participants will train for 6 weeks, 15 min twice daily, 5 d/week at 60% of the peak of minute ventilation, at home by means of a respiratory device (SpiroTiger, Idiag, Fehraltorf, CH) Control/sham: participants will undergo the training protocol at rest's minute ventilation
Outcomes	Respiratory muscle strength (P _I max)
Starting date	14 March 2017
Contact information	ferid.oueslati@criucpq.ulaval.ca
Notes	

NCT04387318

Study name	Inspiratory muscle training and neuromuscular electrical stimulation in chronic obstructive pulmonary disease
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> • Clinical diagnosis of COPD, in stages 2, 3 and 4 according to GOLD criteria • Clinically stable, i.e. absence of infections or exacerbations in the last 3 months • The medical team allows the patient to exercise • Availability of attending the rehabilitation program Exclusion criteria <ul style="list-style-type: none"> • Unstable primary pathologies (cardiovascular, renal, metabolic, psychiatric) • Hemodynamic instability • Nutritional supplementation in the 4 weeks preceding the study • Severe hearing or a visual impairment recorded on patient chart or self-referred • Obesity (BMI > 30 kg/m²) • A neurological or musculoskeletal condition that severely limits mobility and postural control, thus making it impossible to carry out the assessments • Electronic devices, such as heart pacemakers and implantable cardioverter-defibrillator • Skin injuries and infection where electrodes would be placed • Prior participation in pulmonary rehabilitation programs 3 months previous to the study • Vertigo
Interventions	PR+IMT: <ul style="list-style-type: none"> • PR: aerobic and resistance exercise for 8 weeks • IMT: participants will train with POWERbreathe Medic Plus (POWERbreathe International Ltd., England, UK) inspiratory training device for 5 sets of 10 repetitions each, with a 1-minute interval between each set. The initial load set will be 30% of P_Imax, during the first 2 weeks to allow for an adjustment period. After that, load increases occurred as follows: 35% of P_Imax in week 3, 40% of P_Imax in week 4, 45% of P_Imax in week 5, 50% of P_Imax at week 6, 55% of P_Imax in week 7, and 60% of P_Imax in week 8 PR: this group will undergo only the PR protocol described above
Outcomes	HRQoL: SGRQ Respiratory muscle strength (P _I max) Respiratory muscle endurance
Starting date	1 October 2019
Contact information	albuisa@gmail.com
Notes	

NCT04802096

Study name	Effects of inspiratory muscle training in addition to pulmonary rehabilitation in patients with moderate to severe COPD exacerbation
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> • Age > 20 years old • Diagnosed as moderate exacerbation of COPD • MIP < 80 cmH2O Exclusion criteria <ul style="list-style-type: none"> • Any clinical diagnosis that will influence the measurement, including any history of neuromyopathy • Angina, acute myocardial infarction in the previous month • Pregnancy • Participated in IMT program in the previous 3 months • Any psychiatric or cognitive disorders, e.g. Mini-Mental State Examination < 24, that will disturb the communication and co-operation of the study
Interventions	PR+IMT: <ul style="list-style-type: none"> • PR: aerobic exercise training, strength training, and education related to airway clearance and drug utilization. All the participants will receive 8-week PR • IMT: the intensity of IMT will be set at 30% of MIP. Participants in this group will perform 15 breaths/set, 6 sets/d on a daily basis. The intervention will last for 8 weeks PR: this group will undergo the same PR program as the intervention group
Outcomes	Functional exercise capacity: 6MWD HRQoL: SGRQ HRQoL: CAT
Starting date	15 March 2021
Contact information	Sponsorship source: National Taiwan University Hospital Country: Taiwan Author's name: Wei-Yu Huang Institution: School and Graduate Institute of Physical Therapy of National Taiwan University Email: r08428013@ntu.edu.tw Address: Taipei, Zhongzheng Dist, Taiwan, 100 Clinical trial register: NCT04802096
Notes	

RBR-10nyzqc

Study name	Effects of Inspiratory Muscle Training on Breathlessness, Exercise Capacity and Postural Control in Patients with COPD
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> • Age > 40 years • Diagnosis of COPD according to GOLD (2016) • Absence of exacerbations in the last 30 days • Presence of inspiratory muscle weakness (P_Imax < 70 cmH₂O or < 70% of predicted) Exclusion criteria <ul style="list-style-type: none"> • Diagnosis of psychiatric or cognitive diseases that compromise the understanding of the study's guidelines • Progressive neurological disease • Neuromuscular disease; Vestibular disorders • Orthopedic changes that compromise the results of field and range tests Presence of other co-morbidities that, at the researcher's discretion, may interfere with the results of the study
Interventions	IMT: 22 patients will train at around 50% of their P _I max. The training consists of 30 breaths in a row, with deep and strong inspiration (lasting 4-5 minutes), twice a day, every day for 8 weeks. Each week, the patient will return to our center to perform a new manovacuometry and update the load value, maintaining the ~ 50% of the P _I max value, and after this update, one of the sessions of the day will be held in person. Sham IMT: 16 patients will train at around 10% of their P _I max. The training consists of 30 breaths in a row, with deep and strong inspiration (lasting 4-5 minutes), twice a day, every day for 8 weeks. Every week, the patient will return to our center to perform a new manovacuometry, the data will be recorded, however the load will be kept at around 10% of the initial P _I max value, after this registration, one of the sessions of the day will be held in person. This value will not make any changes during the study period. After the eight-week protocol, patients will return to perform the initial evaluations and tests again, in the same order they did the first time, and will receive a new medical order for serum vitamin D and calcium measurement for comparison with the first. The CG participants after the study period will undergo the same protocol applied to the IG as treatment.
Outcomes	Dyspnea: mMRC Functional exercise capacity: 6MWD
Starting date	
Contact information	yves.souza@uva.br
Notes	

RBR 42rmqy

Study name	Inspiratory muscle training in chronic obstructive pulmonary disease oxygen-dependent patients: a randomized controlled trial
------------	---

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

161

RBR 42rmqy (Continued)

Methods	<p>Study design: RCT</p> <p>Study grouping: parallel-group</p>
Participants	<p>IMT</p> <ul style="list-style-type: none"> • N (randomized): 30 <p>Control/sham</p> <ul style="list-style-type: none"> • N (randomized): 30 <p>Inclusion criteria</p> <ul style="list-style-type: none"> • Individuals with COPD • Both genders • Stable clinical condition • Without cardiac disease (heart failure, angina pectoris) • No history of respiratory infection at least 30 days before inclusion of study <p>Exclusion criteria</p> <ul style="list-style-type: none"> • Inability to walk • Inability to do 6MWD • Neuromuscular disease • Non-controlled comorbidities such as arterial hypertension and diabetes mellitus
Interventions	<p>Intervention characteristics</p> <p>IMT: the experimental group will have 30 individuals that will receive home-based IMT, 7 d/week, over 4 weeks. The daily training will be split into 2 daily sessions (i.e. morning and afternoon). Each daily session will be comprised of four 4-min sets of respiratory training with 1-min rest intervals between each set. The resistance will be provided by the Power Breathe Classic, which allows individuals to exercise the inspiratory muscles during the training session. Training will be individually tailored for each participant. The initial training load for each participant will be set at 50% of his/her P_Imax. The participants will be trained and instructed to do the exercise program on their own with no supervision. Once a week, during the physical therapist's home visit, the clinician will determine the new values for maximal inspiratory strength, and will adjust the load to 50% of the new values. The device will be covered with opaque material, so that participants will be blinded to the training load.</p> <p>Control/sham: the control group with 30 individuals will receive a sham intervention. Sham respiratory muscle training will be delivered using a Power Breathe with no resistance (0 cmH₂O) or progression, over the 4-week period, 40 min/d, 7 d/week</p>
Outcomes	<p>HRQoL: ADL</p> <p>Respiratory muscle strength (P_Imax)</p> <p>Laboratory exercise test: VO₂peak</p>
Starting date	30 October 2019
Contact information	viniciusmaldaner@gmail.com
Notes	

TCTR20210604001

Study name	Is single bout of inspiratory muscle training alter blood pressure and cardio autonomies modulation in COPD patients? : a pilot study
Methods	Study design: RCT Study grouping: parallel-group
Participants	Inclusion criteria <ul style="list-style-type: none"> • GOLD stage 2-3 COPD patients, which was defined by FEV1 30%-80%pred • Free of acute exacerbation of COPD > 8 weeks before the study • BMI 18.5 kg/m²-29.9 kg/m² Exclusion criteria <ul style="list-style-type: none"> • History of recent lung surgery, spontaneous pneumothorax or rib fracture < 12 months prior to the study • History of stroke, diabetes, cardiovascular or neuromuscular disorder • Resting blood pressure > 180/110 mmHg or < 90/60 mmHg • Fracture at upper, lower extremity or spine < 6 months prior to the study • Lower extremity and spine pain which was assessed by visual analog scale > 3 • Regular exercise which was defined as having exercise or perform physical activity • Unable to communicate and follow instructions • Resting supplemental oxygen • Age minimum: 40 years • Age maximum: 70 years • Gender: male
Interventions	Intervention characteristics IMT: participants will perform single session of IMT exercise with 60% of MIP. They will complete 6 breaths/set for 5 sets with 1-min rest between each set Sham IMT: participants will undergo the same program described above at no load
Outcomes	
Starting date	11 January 2021
Contact information	Sponsorship source: Khon Kaen University Khon Kaen, Thailand Country: Thailand Author's name: EAKARACH WONGSAYA Institution: School of Physical Therapy, Faculty of Associated Medical Sciences, Khon Kaen University Email: eakarach.wo@up.ac.th Address: 123 School of Physical Therapy, Faculty of Associated Medical Sciences, Khon Kaen University University 40002 Khon Kaen Thailand Clinical trial register: TCTR20210604001
Notes	

Tctr 2022

Study name	Effect of inspiratory muscle training on cardiovascular autonomic functions in chronic obstructive pulmonary disease patients
Methods	Study design: RCT Study grouping: parallel-group
Participants	Patients with COPD
Interventions	IMT Home-based IMT will be performed daily using PowerBreathe device. Total eight week of IMT will be provided. The IMT group will start training at 40% of their initial P _{Imax} . The new P _{Imax} value will be measured every week by the researcher. The training load in each week will be increased continuously over time by adjusting to at least 50% of P _{Imax} or gradually increase to the highest tolerable intensity during each of the supervised sessions. The highest tolerable intensity means that the rates of perceived inspiratory effort using a modified Borg dyspnea scale will be the range 4 to 6 of 10 (moderate to very severe). Daily training will consist of 2 sessions of 30 breaths.,The participants will receive only the standard care treatment from the physicians such as pharmacological treatment. They will meet the physician at week 1 and week 8. Control This group will not receive any intervention
Outcomes	Dyspnea: <ul style="list-style-type: none">• mMRC• BDI-TDI Functional exercise capacity: 6MWD Respiratory muscle strength: P _{Imax} Spirometry measurements
Starting date	2022
Contact information	
Notes	

6MWD: six-minute walk distance; **ADL:** activities of daily living; **BMI:** body mass index; **CAT:** Chronic Obstructive Pulmonary Disease Assessment Test; **COPD:** chronic obstructive pulmonary disease; **CRQ:** Chronic respiratory disease questionnaire; **DBP:** diastolic blood pressure; **FEV1:** forced expiratory volume at 1 second; **FVC:** forced vital capacity; **GOLD:** Global Initiative of Chronic Obstructive Lung Disease; **HRQoL:** health-related quality of life; **IMT:** inspiratory muscle training; **MIP:** maximal inspiratory pressure; **mMRC:** Modified Medical Research Council; **P_{Imax}:** maximal inspiratory pressure; **PR:** pulmonary rehabilitation; **RCT:** randomized controlled trial; **SBP:** systolic blood pressure; **SGRQ:** St George's Respiratory Questionnaire; **SMIP:** sustained maximal inspiratory pressure; **TIRE:** test of incremental respiratory endurance

RISK OF BIAS

Legend:  Low risk of bias  High risk of bias  Some concerns

Risk of bias for analysis 1.1 Dyspnea: Borg (at submaximal exercise: 50% to 80% of Wmax)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Charususin 2018	✓	✓	✓	✓	✓	✓
Larson 1999	⚠	✗	✗	✗	⚠	✗

Risk of bias for analysis 1.2 Dyspnea: Modified Medical Research Council (mMRC)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Beaumont 2018	✓	✓	✓	✗	✓	✗
Wang 2017	✓	✓	✓	✗	⚠	✗

Risk of bias for analysis 1.3 Functional exercise capacity: 6-minute walk distance (6MWD) (meters)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Beaumont 2015	⚠	✓	✓	✓	✓	⚠
Beaumont 2018	✓	✓	✓	✓	✓	✓
Charususin 2018	✓	✓	✓	✓	✓	✓
De Farias 2019	⚠	⚠	✓	✓	⚠	⚠
De Farias 2019	⚠	⚠	✓	✓	⚠	⚠
Dellweg 2017	✓	✓	✓	✓	✓	✓
Mador 2005	⚠	⚠	⚠	✓	⚠	⚠

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Magadle 2007	~	✓	✓	✓	~	~
Paneroni 2018	✓	✓	✗	✓	✓	✗
Schultz 2018	~	✓	✓	✓	✓	~
Tounsi 2021	✓	✓	✓	✓	✓	✓
Tout 2013	✗	✓	✓	✓	✓	✗
Wang 2017	✓	✓	✓	✓	~	~

Risk of bias for analysis 1.4 Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: duration of the intervention)

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Subgroup 1.4.1 Short-term (<4 weeks)						
Beaumont 2015	~	✓	✓	✓	✓	~
Dellweg 2017	✓	✓	✓	✓	✓	✓
Paneroni 2018	✓	✓	✗	✓	✓	✗
Schultz 2018	~	✓	✓	✓	✓	~
Subgroup 1.4.2 Medium-term (≥4 weeks and <8 weeks)						
Beaumont 2018	✓	✓	✓	✓	✓	✓
Dellweg 2017	✓	✓	✓	✓	✓	✓
Subgroup 1.4.3 Long-term (≥8 weeks)						

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Charususin 2018	✓	✓	✓	✓	✓	✓
De Farias 2019	~	~	✓	✓	~	~
De Farias 2019	~	~	✓	✓	~	~
Mador 2005	~	~	~	✓	~	~
Magadle 2007	~	✓	✓	✓	~	~
Tounsi 2021	✓	✓	✓	✓	✓	✓
Tout 2013	✗	✓	✓	✓	✓	✗
Wang 2017	✓	✓	✓	✓	~	~

Risk of bias for analysis 1.5 Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: with or without respiratory muscle weakness)

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Subgroup 1.5.1 With respiratory muscle weakness						
Charususin 2018	✓	✓	✓	✓	✓	✓
Dellweg 2017	✓	✓	✓	✓	✓	✓
Tout 2013	✗	✓	✓	✓	✓	✗
Subgroup 1.5.2 Without respiratory muscle weakness						
Beaumont 2015	~	✓	✓	✓	✓	~
Beaumont 2018	✓	✓	✓	✓	✓	✓

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
De Farias 2019	~	~	✓	✓	~	~
De Farias 2019	~	~	✓	✓	~	~
Mador 2005	~	~	~	✓	~	~
Magadle 2007	~	✓	✓	✓	~	~
Paneroni 2018	✓	✓	✗	✓	✓	✗
Schultz 2018	~	✓	✓	✓	✓	~
Tounsi 2021	✓	✓	✓	✓	✓	✓
Wang 2017	✓	✓	✓	✓	~	~

Risk of bias for analysis 1.7 Functional exercise capacity: Wmax (watt)

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Charusisin 2018	✓	✓	✓	✓	✓	✓
Dekhuijzen 1991	~	✓	✓	✓	✗	✗
Larson 1999	~	✗	✗	✓	~	✗
Mador 2005	~	~	~	✓	~	~
Wang 2017	✓	✓	✓	✓	~	~

Risk of bias for analysis 1.9 Health-related quality of life (HRQoL): St George's Respiratory Questionnaire (SGRQ)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 1.9.1 Symptoms						
Beaumont 2018	✓	✓	✓	✗	✓	✗
Tout 2013	✗	✓	✓	✗	~	✗
Subgroup 1.9.2 Activity						
Beaumont 2018	✓	✓	✓	✗	✓	✗
Tout 2013	✗	✓	✓	✗	~	✗
Subgroup 1.9.3 Impact						
Beaumont 2018	✓	✓	✓	✗	✓	✗
Tout 2013	✗	✓	✓	✗	~	✗
Subgroup 1.9.4 Total						
Abedi Yekta 2019	~	✓	✓	✗	✓	✗
Beaumont 2018	✓	✓	✓	✗	✓	✗
Magadle 2007	~	✓	✓	✓	~	~
Majewska-Pul-sakowska 2016	~	✓	✓	✗	~	✗
Schultz 2018	~	✓	✓	✓	✓	~
Tout 2013	✗	✓	✓	✗	~	✗
Wang 2017	✓	✓	✓	✗	~	✗

Risk of bias for analysis 1.10 Health-related quality of life (HRQoL): Chronic Respiratory Disease Questionnaire (CRQ)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 1.10.1 Dyspnea						
Charususin 2018	✓	✓	✓	✓	✓	✓
Larson 1999	~	✗	✗	~	~	✗
Mador 2005	~	✓	~	✗	~	✗
Subgroup 1.10.2 Fatigue						
Charususin 2018	✓	✓	✓	✓	✓	✓
Larson 1999	~	✗	✗	~	~	✗
Mador 2005	~	✓	~	✗	~	✗
Subgroup 1.10.3 Emotion						
Charususin 2018	✓	✓	✓	✓	✓	✓
Mador 2005	~	✓	~	✗	~	✗
Subgroup 1.10.4 Mastery						
Charususin 2018	✓	✓	✓	✓	✓	✓
Mador 2005	~	✓	~	✗	~	✗

Risk of bias for analysis 1.11 Health-related quality of life (HRQoL): COPD Assessment Test (CAT)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Schultz 2018	~	✓	✓	✓	✓	~

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Wang 2017	✓	✓	✓	✗	~	✗

Risk of bias for analysis 1.12 Inspiratory muscle strength: PImax (cmH20)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Beaumont 2015	~	✓	✓	✓	✓	~
Beaumont 2018	✓	✓	✓	✓	✓	✓
Berry 1996	~	✓	✓	✓	~	~
Charususin 2018	✓	✓	✓	✓	✓	✓
De Farias 2019	~	~	✓	✓	~	~
De Farias 2019	~	~	✓	✓	~	~
Dekhuijzen 1991	~	✓	✓	✓	~	~
Dellweg 2017	✓	✓	✓	✓	✓	✓
Fanfa Bordin 2020	✓	✓	✓	✓	✓	✓
Larson 1999	~	~	✗	✓	~	✗
Mador 2005	~	~	~	✓	~	~
Magadle 2007	~	✓	✓	✓	~	~
Paneroni 2018	✓	✓	✗	✓	✓	✗
Schultz 2018	~	✓	✓	✓	✓	✓

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Tounsi 2021	✓	✓	✓	✓	✓	✓
Wang 2017	✓	✓	✓	✓	~	~
Weiner 1992	~	✓	✓	✓	~	~
Weiner 2000	✗	✓	✓	✓	~	✗

Risk of bias for analysis 1.13 Inspiratory muscle strength: P_{Imax} (cmH₂O) (subgroup analysis: duration of the intervention)

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Subgroup 1.13.1 Short-term (<4 weeks)						
Beaumont 2015	~	✓	✓	✓	✓	~
Dellweg 2017	✓	✓	✓	✓	✓	✓
Paneroni 2018	✓	✓	✗	✓	✓	✗
Schultz 2018	~	✓	✓	✓	✓	✓
Subgroup 1.13.2 Medium-term (≥ 4 weeks and <8 weeks)						
Beaumont 2018	✓	✓	✓	✓	✓	✓
Dekhuijzen 1991	~	✓	✓	✓	~	~
Dellweg 2017	✓	✓	✓	✓	✓	✓
Weiner 2000	✗	✓	✓	✓	~	✗
Subgroup 1.13.3 Long-term (≥ 8 weeks)						
Berry 1996	~	✓	✓	✓	~	~

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Charususin 2018	✓	✓	✓	✓	✓	✓
De Farias 2019	~	~	✓	✓	~	~
De Farias 2019	~	~	✓	✓	~	~
Dekhuijzen 1991	~	✓	✓	✓	~	~
Fanfa Bordin 2020	✓	✓	✓	✓	✓	✓
Larson 1999	~	~	✗	✓	~	✗
Mador 2005	~	~	~	✓	~	~
Magadle 2007	~	✓	✓	✓	~	~
Tounsi 2021	✓	✓	✓	✓	✓	✓
Wang 2017	✓	✓	✓	✓	~	~
Weiner 1992	~	✓	✓	✓	~	~

Risk of bias for analysis 1.14 Inspiratory muscle strength: P_{Imax} (cmH₂O) (subgroup analysis: with or without respiratory muscle weakness)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 1.14.1 With respiratory muscle weakness						
Charususin 2018	✓	✓	✓	✓	✓	✓
Dekhuijzen 1991	~	✓	✓	✓	~	~
Dellweg 2017	✓	✓	✓	✓	✓	✓

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Weiner 1992	~	✓	✓	✓	~	~
Weiner 2000	✗	✓	✓	✓	~	✗
Subgroup 1.14.2 Without respiratory muscle weakness						
Beaumont 2015	~	✓	✓	✓	✓	~
Beaumont 2018	✓	✓	✓	✓	✓	✓
De Farias 2019	~	~	~	✓	~	~
De Farias 2019	~	~	~	✓	~	~
Fanfa Bordin 2020	✓	✓	✓	✓	✓	✓
Larson 1999	~	~	✗	✓	~	✗
Mador 2005	~	~	~	✓	~	~
Magadle 2007	~	✓	✓	✓	~	~
Paneroni 2018	✓	✓	✗	✓	✓	✗
Schultz 2018	~	✓	✓	✓	✓	✓
Tounsi 2021	✓	✓	✓	✓	✓	✓
Wang 2017	✓	✓	✓	✓	~	~

Risk of bias for analysis 2.1 Dyspnea: Borg (at submaximal exercise capacity)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Dacha 2019						
Hill 2006						
Koppers 2006						
Langer 2018						
Larson 1999						
Petrovic 2012						

Risk of bias for analysis 2.2 Dyspnea: Baseline and Transition Dyspnea Indexes (BDI-TDI)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.2.1 Functional impairment						
Harver 1989						
Weiner 2003						
Wu 2017						
Wu 2017						
Subgroup 2.2.2 Magnitude of task						
Harver 1989						
Weiner 2003						
Wu 2017						

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Wu 2017	~	✓	✓	✗	~	✗
Subgroup 2.2.3 Magnitude of effort						
Harver 1989	~	✓	✗	✓	~	✗
Weiner 2003	~	✓	✓	✓	✗	✗
Wu 2017	~	✓	✓	✗	~	✗
Wu 2017	~	✓	✓	✗	~	✗
Subgroup 2.2.4 Focal score						
Chuang 2017	~	✓	✗	✗	~	✗
Harver 1989	~	✓	✗	✓	~	✗
Langer 2018	✓	✓	✓	✓	✓	✓
Lisboa 1997	~	✓	✓	✓	~	~
Sanchez Riera 2001	~	✓	✓	✓	✗	✗
Scherer 2000	~	✓	✓	✓	~	~
Weiner 2003	~	✓	✓	✓	✗	✗
Wu 2017	~	✓	✓	✗	~	✗
Wu 2017	~	✓	✓	✗	~	✗

Risk of bias for analysis 2.3 Dyspnea: Transition Dyspnea Index (TDI): Focal score (subgroup analysis: with or without respiratory muscle weakness)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.3.1 With respiratory muscle weakness						
Chuang 2017	~	✓	✗	✗	~	✗
Harver 1989	~	✓	✗	✓	~	✗
Sanchez Riera 2001	~	✓	✓	✓	✗	~
Wu 2017	~	✓	✓	✗	~	✗
Wu 2017	~	✓	✓	✗	~	✗
Subgroup 2.3.2 Without respiratory weakness						
Langer 2018	✓	✓	✓	✓	✓	✓
Lisboa 1997	~	✓	✓	✓	~	~
Scherer 2000	~	✓	✓	✓	~	~

Risk of bias for analysis 2.4 Dyspnea: Modified Medical Research Council (mMRC)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Langer 2018	✓	✓	✓	✓	✓	✓
Saka 2021	~	✓	✓	✓	~	~
Xu 2018	✓	✓	✓	✓	✓	✓
ZhouL 2016	✓	~	~	✗	~	✗

Risk of bias for analysis 2.5 Functional exercise capacity: 6-minute walk distance (6MWD) (meters)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Bavarsad 2015	~	~	✓	✓	~	~
Beckerman 2005	~	✗	✗	✓	~	✗
Chuang 2017	~	✓	✓	✓	~	~
Cutrim 2019	✓	✓	✓	✓	✓	✓
Hill 2006	~	✓	✓	✓	~	~
Hsiao 2003	~	✗	~	✓	~	✗
Hsiao 2003	~	✗	~	✓	~	✗
Koppers 2006	~	✓	✓	✓	~	~
Leelarungrayub 2017	~	✓	✗	✓	~	✗
Lisboa 1997	~	✓	✓	✓	~	~
Ramirez Sarmiento 2002	~	✗	✓	✓	~	✗
Saher 2021	~	✗	~	✓	~	✗
Saka 2021	~	✓	✓	✓	~	~
Scherer 2000	~	✓	~	✓	~	~
Weiner 2003	~	✓	✓	✓	✗	✗
Xu 2018	✓	✓	✓	✓	✓	✓
ZhouL 2016	✓	~	~	✓	~	~

Risk of bias for analysis 2.6 Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: duration of the intervention)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.6.1 Short-term (<4 weeks)						
Saher 2021	~	✗	~	✓	~	✗
Subgroup 2.6.2 Medium-term (≥4 weeks and <8 weeks)						
Chuang 2017	~	✓	✓	✓	~	~
Koppers 2006	~	✓	✓	✓	~	~
Leelarungrayub 2017	~	✓	✗	✓	~	✗
Lisboa 1997	~	✓	✓	✓	~	~
Subgroup 2.6.3 Long-term (≥8 weeks)						
Bavarsad 2015	~	~	✓	✓	~	~
Beckerman 2005	~	✗	✗	✓	~	~
Cutrim 2019	✓	✓	✓	✓	✓	✓
Hill 2006	~	✓	✓	✓	~	~
Hsiao 2003	~	✗	~	✓	~	✗
Hsiao 2003	~	✗	~	✓	~	✗
Ramirez Sarmiento 2002	~	✗	✓	✓	~	✗
Saka 2021	~	✓	✓	✓	~	~
Scherer 2000	~	✓	~	✓	~	~
Weiner 2003	~	✓	✓	✓	✗	✗
Xu 2018	✓	✓	✓	✓	✓	✓

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
ZhouL 2016						

Risk of bias for analysis 2.7 Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: with or without respiratory muscle weakness)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.7.1 With respiratory muscle weakness						
Chuang 2017						
Leelarungrayub 2017						
Saher 2021						
Xu 2018						
ZhouL 2016						
Subgroup 2.7.2 Without respiratory muscle weakness						
Bavarsad 2015						
Beckerman 2005						
Hill 2006						
Hsiao 2003						
Koppers 2006						
Lisboa 1997						
Ramirez Sarmiento 2002						

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Saka 2021	~	✓	✓	✓	~	~
Scherer 2000	~	✓	~	✓	~	~
Weiner 2003	~	✓	✓	✓	✗	✗
Xu 2018	✓	✓	✓	✓	✓	✓

Risk of bias for analysis 2.9 Functional exercise capacity: Wmax (watt)

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Hill 2006	~	✓	✓	✓	✓	~
Koppers 2006	~	✓	✓	✓	~	~
Larson 1999	~	✗	✗	✓	~	✗
Lisboa 1997	~	✓	✓	✓	~	~
Ramirez Sarmiento 2002	~	✗	✓	✓	~	✗
Sanchez Riera 2001	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~

Risk of bias for analysis 2.12 Health-related quality of life (HRQoL): St George Respiratory Questionnaire (SGRQ)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.12.1 Symptoms						
Berton 2015	~	~	~	✓	~	~
Saka 2021	~	✓	✓	✓	~	~
Subgroup 2.12.2 Activity						
Berton 2015	~	~	~	✓	~	~
Saka 2021	~	✓	✓	✓	~	~
Subgroup 2.12.3 Impact						
Berton 2015	~	~	~	✓	~	~
Saka 2021	~	✓	✓	✓	~	~
Subgroup 2.12.4 Total						
Abedi Yekta 2019	~	✓	~	✗	✓	✗
Beckerman 2005	~	✗	✗	✓	~	✗
Berton 2015	~	~	~	✓	~	~
Majewska-Pul-sakowska 2016	~	✓	✓	✗	~	✗
Saka 2021	~	✓	✓	✓	~	~
Xu 2018	✓	✓	✓	✓	✓	✓

Risk of bias for analysis 2.13 Health-related quality of life (HRQoL): Chronic Respiratory Disease Questionnaire (CRQ)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.13.1 Dyspnea						
Bustamante 2007	~	✓	✓	✓	~	~
Hill 2006	~	✓	✓	✓	~	~
Larson 1999	~	✗	✗	~	~	✗
Nikoleitou 2016	~	~	~	✓	~	~
Wu 2017	~	✓	✓	✗	~	✗
Wu 2017	~	✓	✓	✗	~	✗
Subgroup 2.13.2 Fatigue						
Bustamante 2007	~	✓	✓	✓	~	~
Hill 2006	~	✓	✓	✓	~	~
Larson 1999	~	✗	✗	~	~	✗
Nikoleitou 2016	~	~	~	✓	~	~
Wu 2017	~	✓	✓	✗	~	✗
Wu 2017	~	✓	✓	✗	~	✗
Subgroup 2.13.3 Emotion						
Bustamante 2007	~	✓	✓	✓	~	~
Hill 2006	~	✓	✓	✓	~	~
Nikoleitou 2016	~	~	~	✓	~	~
Wu 2017	~	✓	✓	✗	~	✗

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Wu 2017	⚠	✓	✓	✗	⚠	✗
Subgroup 2.13.4 Mastery						
Bustamante 2007	⚠	✓	✓	✓	⚠	⚠
Hill 2006	⚠	✓	✓	✓	⚠	⚠
Nikoleitou 2016	⚠	⚠	⚠	✓	⚠	⚠
Wu 2017	⚠	✓	✓	✗	⚠	✗
Wu 2017	⚠	✓	✓	✗	⚠	✗

Risk of bias for analysis 2.14 Health-related quality of life (HRQoL): COPD Assessment Test (CAT)

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Saka 2021	⚠	✓	✓	✓	⚠	⚠
Xu 2018	✓	✓	✓	✓	✓	✓

Risk of bias for analysis 2.15 Inspiratory muscle strength: PImax (cmH₂O)

Bias						
Study	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	Overall
Beckerman 2005	⚠	✗	✗	✓	⚠	✗
Belman 1988	⚠	✓	✓	✓	⚠	⚠

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Berton 2015	~	✓	✓	✓	~	~
Bustamante 2007	~	✓	✓	✓	~	~
Chuang 2017	~	✓	✓	✓	~	~
Covey 2001	~	~	~	✓	~	~
Cutrim 2019	✓	✓	✓	✓	✓	✓
Dacha 2019	✓	✓	✓	✓	~	~
Harver 1989	~	✓	✗	✓	~	✗
Heijdra 1996	~	✓	✓	✓	~	~
Hill 2006	~	✓	✓	✓	~	~
Hsiao 2003	~	✗	~	✓	~	✗
Hsiao 2003	~	✗	~	✓	~	✗
Kim 1993	~	✓	~	✓	~	~
Koppers 2006	~	✓	✓	✓	~	~
Langer 2018	✓	✓	✓	✓	✓	✓
Larson 1988	✓	✗	~	✓	~	✗
Larson 1999	~	~	✗	✓	~	✗
Leelarungrayub 2017	~	✓	✗	✓	~	✗
Lisboa 1997	~	✓	✓	✓	~	~
Nikoletou 2016	~	~	~	✓	~	~

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Petrovic 2012	~	✓	✓	✓	~	~
Preusser 1994	~	✓	✓	✓	~	~
Ramirez Sarmiento 2002	~	✗	✓	✓	~	✗
Saher 2021	~	✗	~	✓	~	✗
Saka 2021	~	✓	✓	✓	~	~
Sanchez Riera 2001	~	✓	✓	✓	~	~
Scherer 2000	~	✓	~	✓	~	~
Weiner 2003	~	✓	✓	✓	✗	✗
Weiner 2006	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Xu 2018	✓	✓	✓	✓	✓	✓
ZhouL 2016	✓	~	~	✓	~	~

Risk of bias for analysis 2.16 Inspiratory muscle strength: P_{lmax} (cmH₂O) (subgroup analysis: duration of the intervention)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.16.1 Short-term (<4 weeks)						
Saher 2021	~	✗	~	✓	~	✗

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.16.2 Medium-term (≥4 weeks and <8 weeks)						
Belman 1988	~	✓	✓	✓	~	~
Bustamante 2007	~	✓	✓	✓	~	~
Chuang 2017	~	✓	✓	✓	~	~
Koppers 2006	~	✓	✓	✓	~	~
Larson 1988	✓	✗	~	✓	~	✗
Leelarungrayub 2017	~	✓	✗	✓	~	✗
Nikoletou 2016	~	~	~	✓	~	~
Ramirez Sarmiento 2002	~	✗	✓	✓	~	✗
Subgroup 2.16.3 Long-term (≥8 weeks)						
Beckerman 2005	~	✗	✗	✓	~	✗
Berton 2015	~	✓	✓	✓	~	~
Chuang 2017	~	✓	✓	✓	~	~
Covey 2001	~	~	~	✓	~	~
Cutrim 2019	✓	✓	✓	✓	✓	✓
Dacha 2019	✓	✓	✓	✓	~	~
Harver 1989	~	✓	✗	✓	~	✗
Heijdra 1996	~	✓	✓	✓	~	~
Hill 2006	~	✓	✓	✓	~	~

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Hsiao 2003	~	✗	~	✓	~	✗
Hsiao 2003	~	✗	~	✓	~	✗
Kim 1993	~	✓	~	✓	~	~
Langer 2018	✓	✓	✓	✓	✓	✓
Larson 1988	✓	✗	~	✓	~	✗
Larson 1999	~	~	✗	✓	~	✗
Lisboa 1997	~	✓	✓	✓	~	~
Petrovic 2012	~	✓	✓	✓	~	~
Preusser 1994	~	✓	✓	✓	~	~
Ramirez Sarmiento 2002	~	✗	✓	✓	~	✗
Saka 2021	~	✓	✓	✓	~	~
Sanchez Riera 2001	~	✓	✓	✓	~	~
Scherer 2000	~	✓	~	✓	~	~
Weiner 2003	~	✓	✓	✓	✗	✗
Weiner 2006	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Xu 2018	✓	✓	✓	✓	✓	✓
ZhouL 2016	✓	~	~	✓	~	~

Risk of bias for analysis 2.17 Inspiratory muscle strength: P_{lmax} (cmH₂O) (subgroup analysis: with or without respiratory muscle weakness)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.17.1 With respiratory muscle weakness						
Chuang 2017	~	✓	✓	✓	~	~
Harver 1989	~	✓	✗	✓	~	✗
Leelarungrayub 2017	~	✓	✗	✓	~	✗
Preusser 1994	~	✓	✓	✓	~	~
Saher 2021	~	✗	~	✓	~	✗
Sanchez Riera 2001	~	✓	✓	✓	~	~
Weiner 2006	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Xu 2018	✓	✓	✓	✓	✓	✓
ZhouL 2016	✓	~	~	✓	~	~
Subgroup 2.17.2 Without respiratory muscle weakness						
Beckerman 2005	~	✗	✗	✓	~	✗
Belman 1988	~	✓	✓	✓	~	~
Berton 2015	~	✓	✓	✓	~	~
Bustamante 2007	~	✓	✓	✓	~	~
Covey 2001	~	~	~	✓	~	~

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Dacha 2019	✓	✓	✓	✓	⚠	⚠
Hill 2006	⚠	✓	✓	✓	⚠	⚠
Hsiao 2003	⚠	✗	⚠	✓	⚠	✗
Kim 1993	⚠	✓	⚠	✓	⚠	⚠
Koppers 2006	⚠	✓	✓	✓	⚠	⚠
Langer 2018	✓	✓	✓	✓	✓	✓
Larson 1999	⚠	⚠	✗	✓	⚠	✗
Lisboa 1997	⚠	✓	✓	✓	⚠	⚠
Nikoletou 2016	⚠	⚠	⚠	✓	⚠	⚠
Petrovic 2012	⚠	✓	✓	✓	⚠	⚠
Ramirez Sarmiento 2002	⚠	✗	✓	✓	⚠	✗
Saka 2021	⚠	✓	✓	✓	⚠	⚠
Scherer 2000	⚠	✓	⚠	✓	⚠	⚠
Xu 2018	✓	✓	✓	✓	✓	✓

Risk of bias for analysis 2.18 Inspiratory muscle strength: P_Imax (cmH₂O) (subgroup analysis: method of measurement)

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Subgroup 2.18.1 Residual Volume (RV)						

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Beckerman 2005	~	✗	✗	✓	~	✗
Covey 2001	~	~	~	✓	~	~
Cutrim 2019	✓	✓	✓	✓	✓	✓
Heijdra 1996	~	✓	✓	✓	~	~
Hsiao 2003	~	✗	~	✓	~	✗
Hsiao 2003	~	✗	~	✓	~	✗
Kim 1993	~	✓	~	✓	~	~
Koppers 2006	~	✓	✓	✓	~	~
Larson 1988	✓	✗	~	✓	~	✗
Larson 1999	~	~	✗	✓	~	✗
Leelarungrayub 2017	~	✓	✗	✓	~	✗
Saher 2021	~	✗	~	✓	~	✗
Scherer 2000	~	✓	~	✓	~	~
Weiner 2003	~	✓	✓	✓	✗	✗
Weiner 2006	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Wu 2017	~	✓	✓	✓	~	~
Subgroup 2.18.2 Functional Residual Capacity (FRC)						
Belman 1988	~	✓	✓	✓	~	~

Study	Bias					Overall
	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported results	
Harver 1989	⊘	✓	✗	✓	⊘	✗
Hill 2006	⊘	✓	✓	✓	⊘	⊘
Langer 2018	✓	✓	✓	✓	✓	✓
Lisboa 1997	⊘	✓	✓	✓	⊘	⊘
Nikoleitou 2016	⊘	⊘	⊘	✓	⊘	⊘
Petrovic 2012	⊘	✓	✓	✓	⊘	⊘
Preusser 1994	⊘	✓	✓	✓	⊘	⊘
Sanchez Riera 2001	⊘	✓	✓	✓	⊘	⊘
Subgroup 2.18.3 Not reported						
Berton 2015	⊘	✓	✓	✓	⊘	⊘
Bustamante 2007	⊘	✓	✓	✓	⊘	⊘
Chuang 2017	⊘	✓	✓	✓	⊘	⊘
Dacha 2019	✓	✓	✓	✓	⊘	⊘
Ramirez Sarmiento 2002	⊘	✗	✓	✓	⊘	✗
Saka 2021	⊘	✓	✓	✓	⊘	⊘
Xu 2018	✓	✓	✓	✓	✓	✓
ZhouL 2016	✓	⊘	⊘	✓	⊘	⊘

DATA AND ANALYSES


Comparison 1. PR+IMT vs PR

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 Dyspnea: Borg (at submaximal exercise: 50% to 80% of Wmax)	2	202	Mean Difference (IV, Random, 95% CI)	0.19 [-0.42, 0.79]
1.2 Dyspnea: Modified Medical Research Council (mMRC)	2	204	Mean Difference (IV, Random, 95% CI)	-0.12 [-0.39, 0.14]
1.3 Functional exercise capacity: 6-minute walk distance (6MWD) (meters)	12	1199	Mean Difference (IV, Random, 95% CI)	5.95 [-5.73, 17.63]
1.4 Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: duration of the intervention)	12		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.4.1 Short-term (<4 weeks)	4	687	Mean Difference (IV, Random, 95% CI)	10.25 [-24.98, 45.49]
1.4.2 Medium-term (≥4 weeks and <8 weeks)	2	178	Mean Difference (IV, Random, 95% CI)	36.72 [-67.67, 141.11]
1.4.3 Long-term (≥8 weeks)	7	363	Mean Difference (IV, Random, 95% CI)	7.82 [-6.90, 22.54]
1.5 Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: with or without respiratory muscle weakness)	12		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.5.1 With respiratory muscle weakness	3	219	Mean Difference (IV, Random, 95% CI)	31.17 [-10.50, 72.84]
1.5.2 Without respiratory muscle weakness	9	981	Mean Difference (IV, Random, 95% CI)	0.38 [-11.65, 12.42]
1.6 Functional exercise capacity: 12-minute walk distance (12MWD) (meters)	3	80	Mean Difference (IV, Random, 95% CI)	155.77 [-84.53, 396.08]
1.7 Functional exercise capacity: Wmax (watt)	5	326	Mean Difference (IV, Random, 95% CI)	-1.01 [-6.96, 4.94]
1.8 Functional exercise capacity: exercise time (seconds)	3	192	Mean Difference (IV, Random, 95% CI)	58.62 [-25.09, 142.32]
1.9 Health-related quality of life (HRQoL): St George's Respiratory Questionnaire (SGRQ)	7		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.9.1 Symptoms	2	169	Mean Difference (IV, Random, 95% CI)	-2.33 [-6.28, 1.62]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.9.2 Activity	2	169	Mean Difference (IV, Random, 95% CI)	0.28 [-1.65, 2.20]
1.9.3 Impact	2	169	Mean Difference (IV, Random, 95% CI)	-1.63 [-5.38, 2.11]
1.9.4 Total	7	908	Mean Difference (IV, Random, 95% CI)	0.13 [-0.93, 1.20]
1.10 Health-related quality of life (HRQoL): Chronic Respiratory Disease Questionnaire (CRQ)	3		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.10.1 Dyspnea	3		Mean Difference (IV, Random, 95% CI)	-0.30 [-1.90, 1.29]
1.10.2 Fatigue	3		Mean Difference (IV, Random, 95% CI)	0.28 [-0.76, 1.31]
1.10.3 Emotion	2		Mean Difference (IV, Random, 95% CI)	-0.63 [-2.53, 1.26]
1.10.4 Mastery	2		Mean Difference (IV, Random, 95% CI)	-0.05 [-1.18, 1.08]
1.11 Health-related quality of life (HRQoL): COPD Assessment Test (CAT)	2	657	Mean Difference (IV, Random, 95% CI)	0.13 [-0.80, 1.06]
1.12 Inspiratory muscle strength: P_{lmax} (cmH₂O)	17	1329	Mean Difference (IV, Random, 95% CI)	11.46 [7.42, 15.50]
1.13 Inspiratory muscle strength: P_{lmax} (cmH₂O) (subgroup analysis: duration of the intervention)	17		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.13.1 Short-term (<4 weeks)	4	687	Mean Difference (IV, Random, 95% CI)	12.63 [4.14, 21.11]
1.13.2 Medium-term (≥ 4 weeks and <8 weeks)	4	233	Mean Difference (IV, Random, 95% CI)	12.27 [3.75, 20.79]
1.13.3 Long-term (≥ 8 weeks)	11	478	Mean Difference (IV, Random, 95% CI)	11.52 [5.50, 17.53]
1.14 Inspiratory muscle strength: P_{lmax} (cmH₂O) (subgroup analysis: with or without respiratory muscle weakness)	16		Mean Difference (IV, Random, 95% CI)	Subtotals only
1.14.1 With respiratory muscle weakness	5	282	Mean Difference (IV, Random, 95% CI)	14.84 [11.35, 18.34]
1.14.2 Without respiratory muscle weakness	11	1031	Mean Difference (IV, Random, 95% CI)	10.57 [5.23, 15.91]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.15 Laboratory exercise test: VO ₂ peak (L/min)	5	313	Mean Difference (IV, Random, 95% CI)	-0.01 [-0.05, 0.03]
1.16 Respiratory muscle endurance: respiratory muscle endurance pressure (P _{thmax}) (cmH ₂ O)	2	52	Std. Mean Difference (IV, Random, 95% CI)	1.22 [-0.18, 2.62]
1.17 Respiratory muscle endurance time: Tlim (seconds) (sustained ventilation according to PImax)	3	236	Mean Difference (IV, Random, 95% CI)	84.63 [-50.77, 220.02]
1.18 Respiratory muscle endurance time: Tlim (seconds) (sustained ventilation according to MVV)	2	51	Mean Difference (IV, Random, 95% CI)	477.69 [215.43, 739.94]
1.19 Maximal voluntary ventilation (MVV)	3	93	Std. Mean Difference (IV, Random, 95% CI)	0.40 [-0.02, 0.83]
1.20 Respiratory function: forced expiratory volume at 1 second (FEV1) (%Pred)	6	173	Mean Difference (IV, Random, 95% CI)	0.77 [-1.72, 3.26]
1.21 Respiratory function: forced expiratory volume at 1 second (FEV1) (Liters)	6	889	Mean Difference (IV, Random, 95% CI)	0.04 [-0.04, 0.13]


Analysis 1.1. Comparison 1: PR+IMT vs PR, Outcome 1: Dyspnea: Borg (at submaximal exercise: 50% to 80% of Wmax)



Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 1.2. Comparison 1: PR+IMT vs PR, Outcome 2: Dyspnea: Modified Medical Research Council (mMRC)



Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 1.3. Comparison 1: PR+IMT vs PR, Outcome 3: Functional exercise capacity: 6-minute walk distance (6MWD) (meters)




Footnotes

- (1) Data from adjusted analysis
- (2) Threshold device
- (3) Normocapnic hyperpnea device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 1.7. Comparison 1: PR+IMT vs PR, Outcome 7: Functional exercise capacity: Wmax (watt)



Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 1.8. Comparison 1: PR+IMT vs PR, Outcome 8: Functional exercise capacity: exercise time (seconds)



Footnotes

- (1) Cycling at 80% of Wmax (data from adjusted analysis)
- (2) Cycling at 60 to 70% of Wmax
- (3) Cycling at 40% of Wmax


Analysis 1.9. Comparison 1: PR+IMT vs PR, Outcome 9: Health-related quality of life (HRQoL): St George's Respiratory Questionnaire (SGRQ)



Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 1.10. Comparison 1: PR+IMT vs PR, Outcome 10: Health-related quality of life (HRQoL): Chronic Respiratory Disease Questionnaire (CRQ)



Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 1.11. Comparison 1: PR+IMT vs PR, Outcome 11: Health-related quality of life (HRQoL): COPD Assessment Test (CAT)



Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 1.12. Comparison 1: PR+IMT vs PR, Outcome 12: Inspiratory muscle strength: PImax (cmH2O)




Footnotes

- (1) Threshold device
- (2) Normocapnic hyperpnea

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 1.13. Comparison 1: PR+IMT vs PR, Outcome 13: Inspiratory muscle strength: P1max (cmH2O) (subgroup analysis: duration of the intervention)




Footnotes

- (1) Normocapnic hyperpnea
- (2) Threshold device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 1.14. Comparison 1: PR+IMT vs PR, Outcome 14: Inspiratory muscle strength: PImax (cmH2O) (subgroup analysis: with or without respiratory muscle weakness)




Footnotes

- (1) Threshold device
- (2) Normocapnic hyperpnea

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 1.15. Comparison 1: PR+IMT vs PR, Outcome 15: Laboratory exercise test: VO2peak (L/min)




Footnotes

- (1) We used the mean weight of each group to convert from ml/kg/min to L/min

Analysis 1.16. Comparison 1: PR+IMT vs PR, Outcome 16: Respiratory muscle endurance: respiratory muscle endurance pressure (P_{thmax}) (cmH₂O)




Analysis 1.17. Comparison 1: PR+IMT vs PR, Outcome 17: Respiratory muscle endurance time: T_{lim} (seconds) (sustained ventilation according to P_{lmax})



Footnotes

- (1) Breathing against 50-60% of P_{lmax}
- (2) Breathing against 70% of P_{lmax}
- (3) Breathing against 30% of P_{lmax}

Analysis 1.18. Comparison 1: PR+IMT vs PR, Outcome 18: Respiratory muscle endurance time: T_{lim} (seconds) (sustained ventilation according to MVV)



Footnotes

- (1) Sustained ventilation at 70% of MVV
- (2) Sustained ventilation at 60-75% of MVV

Analysis 1.19. Comparison 1: PR+IMT vs PR, Outcome 19: Maximal voluntary ventilation (MVV)

Study or Subgroup	PR			PR+IMT			Weight	Std. Mean Difference IV, Random, 95% CI	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total			
Berry 1996	57.9	14.81	7	61.9	15.6	9	18.0%	-0.25 [-1.24, 0.74]	
Paneroni 2018	61.4	26.1	12	47.5	14	10	23.7%	0.62 [-0.24, 1.48]	
Wang 2017	4.14	6.19	28	0.9	6.13	27	58.3%	0.52 [-0.02, 1.06]	
Total (95% CI)			47			46	100.0%	0.40 [-0.02, 0.83]	

Heterogeneity: Tau² = 0.01; Chi² = 2.07, df = 2 (P = 0.35); I² = 4%
 Test for overall effect: Z = 1.86 (P = 0.06)
 Test for subgroup differences: Not applicable

Analysis 1.20. Comparison 1: PR+IMT vs PR, Outcome 20: Respiratory function: forced expiratory volume at 1 second (FEV1) (%Pred)

Study or Subgroup	PR+IMT			PR			Weight	Mean Difference IV, Random, 95% CI	Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total			
Berry 1996	47	6.6	7	48.1	6.9	9	14.0%	-1.10 [-7.75, 5.55]	
Dellweg 2017	37.8	13.6	15	33.3	12.7	14	6.8%	4.50 [-5.07, 14.07]	
Magadle 2007	46.5	5.98	14	47.2	5.76	13	31.6%	-0.70 [-5.13, 3.73]	
Majewska-Pulsakowska 2016	61.9	21	13	51.1	14.6	9	2.8%	10.80 [-4.08, 25.68]	
Wang 2017	1.23	8.1	28	0.16	8.1	27	33.8%	1.07 [-3.21, 5.35]	
Weiner 1992	35.2	9	12	33.6	9.7	12	11.1%	1.60 [-5.89, 9.09]	
Total (95% CI)			89			84	100.0%	0.77 [-1.72, 3.26]	

Heterogeneity: Tau² = 0.00; Chi² = 3.12, df = 5 (P = 0.68); I² = 0%
 Test for overall effect: Z = 0.61 (P = 0.54)
 Test for subgroup differences: Not applicable

Analysis 1.21. Comparison 1: PR+IMT vs PR, Outcome 21: Respiratory function: forced expiratory volume at 1 second (FEV1) (Liters)

Study or Subgroup	PR+IMT			PR			Weight	Mean Difference IV, Random, 95% CI	Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total			
Berry 1996	1.39	0.21	7	1.49	0.21	9	11.6%	-0.10 [-0.31, 0.11]	
Charusisin 2018	1.1	0.4	89	1.1	0.5	85	19.4%	0.00 [-0.13, 0.13]	
Majewska-Pulsakowska 2016	1.8	0.7	13	1.3	0.3	9	3.5%	0.50 [0.07, 0.93]	
Schultz 2018	0.21	0.33	300	0.19	0.31	302	33.7%	0.02 [-0.03, 0.07]	
Tout 2013	1.44	0.57	10	0.9	0.47	10	3.1%	0.54 [0.08, 1.00]	
Wang 2017	0.06	0.15	28	0.01	0.15	27	28.7%	0.05 [-0.03, 0.13]	
Total (95% CI)			447			442	100.0%	0.04 [-0.04, 0.13]	

Heterogeneity: Tau² = 0.00; Chi² = 11.49, df = 5 (P = 0.04); I² = 56%
 Test for overall effect: Z = 1.02 (P = 0.31)
 Test for subgroup differences: Not applicable

Comparison 2. IMT vs control/sham


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.1 Dyspnea: Borg (at submaximal exercise capacity)	6	144	Mean Difference (IV, Random, 95% CI)	-0.94 [-1.36, -0.51]
2.2 Dyspnea: Baseline and Transition Dyspnea Indexes (BDI-TDI)	8		Mean Difference (IV, Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.2.1 Functional impairment	3	95	Mean Difference (IV, Random, 95% CI)	0.88 [0.51, 1.25]
2.2.2 Magnitude of task	3	95	Mean Difference (IV, Random, 95% CI)	0.73 [0.35, 1.12]
2.2.3 Magnitude of effort	3	95	Mean Difference (IV, Random, 95% CI)	0.86 [0.42, 1.30]
2.2.4 Focal score	8	238	Mean Difference (IV, Random, 95% CI)	2.98 [2.07, 3.89]
2.3 Dyspnea: Transition Dyspnea Index (TDI): Focal score (subgroup analysis: with or without respiratory muscle weakness)	7		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.3.1 With respiratory muscle weakness	4	152	Mean Difference (IV, Random, 95% CI)	3.52 [2.55, 4.49]
2.3.2 Without respiratory weakness	3	70	Mean Difference (IV, Random, 95% CI)	2.28 [1.10, 3.46]
2.4 Dyspnea: Modified Medical Research Council (mMRC)	4	150	Mean Difference (IV, Fixed, 95% CI)	-0.59 [-0.76, -0.43]
2.5 Functional exercise capacity: 6-minute walk distance (6MWD) (meters)	16	501	Mean Difference (IV, Random, 95% CI)	35.71 [25.68, 45.74]
2.6 Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: duration of the intervention)	16		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.6.1 Short-term (<4 weeks)	1	34	Mean Difference (IV, Random, 95% CI)	33.06 [23.05, 43.07]
2.6.2 Medium-term (≥4 weeks and <8 weeks)	4	131	Mean Difference (IV, Random, 95% CI)	31.15 [1.50, 60.81]
2.6.3 Long-term (≥8 weeks)	11	336	Mean Difference (IV, Random, 95% CI)	38.47 [22.75, 54.20]
2.7 Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: with or without respiratory muscle weakness)	15		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.7.1 With respiratory muscle weakness	5	178	Mean Difference (IV, Random, 95% CI)	33.74 [25.08, 42.40]
2.7.2 Without respiratory muscle weakness	11	291	Mean Difference (IV, Random, 95% CI)	29.80 [12.86, 46.73]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.8 Functional exercise capacity: 12-minute walk distance (12MWD) (meters)	3	101	Mean Difference (IV, Random, 95% CI)	-33.31 [-158.10, 91.48]
2.9 Functional exercise capacity: Wmax (watt)	7	206	Mean Difference (IV, Random, 95% CI)	0.66 [-6.44, 7.76]
2.10 Functional exercise capacity: exercise time (seconds)	5		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.11 Functional exercise capacity: shuttle walk test (SWT) (meters)	2	57	Mean Difference (IV, Random, 95% CI)	-7.45 [-92.74, 77.83]
2.12 Health-related quality of life (HRQoL): St George Respiratory Questionnaire (SGRQ)	6		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.12.1 Symptoms	2	53	Mean Difference (IV, Random, 95% CI)	-2.10 [-3.50, -0.71]
2.12.2 Activity	2	53	Mean Difference (IV, Random, 95% CI)	-9.86 [-15.08, -4.63]
2.12.3 Impact	2	53	Mean Difference (IV, Random, 95% CI)	-6.06 [-13.76, 1.65]
2.12.4 Total	6	182	Mean Difference (IV, Random, 95% CI)	-3.85 [-8.18, 0.48]
2.13 Health-related quality of life (HRQoL): Chronic Respiratory Disease Questionnaire (CRQ)	5		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.13.1 Dyspnea	5	178	Mean Difference (IV, Random, 95% CI)	1.63 [0.23, 3.03]
2.13.2 Fatigue	5	178	Mean Difference (IV, Random, 95% CI)	1.32 [0.08, 2.55]
2.13.3 Emotion	4	163	Mean Difference (IV, Random, 95% CI)	2.64 [0.82, 4.46]
2.13.4 Mastery	4	154	Mean Difference (IV, Random, 95% CI)	1.57 [0.07, 3.06]
2.14 Health-related quality of life (HRQoL): COPD Assessment Test (CAT)	2	86	Mean Difference (IV, Random, 95% CI)	-2.97 [-3.85, -2.10]
2.15 Inspiratory muscle strength: P _{Imax} (cmH ₂ O)	32	916	Mean Difference (IV, Random, 95% CI)	14.57 [9.85, 19.29]
2.16 Inspiratory muscle strength: P _{Imax} (cmH ₂ O) (subgroup analysis: duration of the intervention)	32		Mean Difference (IV, Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.16.1 Short-term (<4 weeks)	1	34	Mean Difference (IV, Random, 95% CI)	12.60 [6.94, 18.26]
2.16.2 Medium-term (≥ 4 weeks and <8 weeks)	8	223	Mean Difference (IV, Random, 95% CI)	11.89 [6.76, 17.02]
2.16.3 Long-term (≥ 8 weeks)	26	748	Mean Difference (IV, Random, 95% CI)	15.02 [9.26, 20.78]
2.17 Inspiratory muscle strength: P_{Imax} (cmH₂O) (subgroup analysis: with or without respiratory muscle weakness)	28		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.17.1 With respiratory muscle weakness	10	323	Mean Difference (IV, Random, 95% CI)	11.08 [7.51, 14.64]
2.17.2 Without respiratory muscle weakness	19	499	Mean Difference (IV, Random, 95% CI)	13.82 [5.36, 22.29]
2.18 Inspiratory muscle strength: P_{Imax} (cmH₂O) (subgroup analysis: method of measurement)	32		Mean Difference (IV, Random, 95% CI)	Subtotals only
2.18.1 Residual Volume (RV)	15	466	Mean Difference (IV, Random, 95% CI)	12.63 [8.46, 16.81]
2.18.2 Functional Residual Capacity (FRC)	9	206	Mean Difference (IV, Random, 95% CI)	11.87 [7.33, 16.41]
2.18.3 Not reported	8	244	Mean Difference (IV, Random, 95% CI)	19.52 [4.63, 34.41]
2.19 Laboratory exercise test: VO_{2peak}	11	286	Std. Mean Difference (IV, Random, 95% CI)	0.31 [0.05, 0.57]
2.20 Respiratory muscle endurance: respiratory muscle endurance pressure (P_{thmax}) (cmH₂O)	8	179	Mean Difference (IV, Random, 95% CI)	9.71 [4.93, 14.50]
2.21 Respiratory muscle endurance time: T_{lim} (seconds)	10	260	Mean Difference (IV, Random, 95% CI)	270.57 [182.44, 358.71]
2.22 Maximal voluntary ventilation (MVV)	2	36	Std. Mean Difference (IV, Random, 95% CI)	0.99 [0.28, 1.69]
2.23 Respiratory function: forced expiratory volume at 1 second (FEV1) (%pred)	10	314	Mean Difference (IV, Random, 95% CI)	2.62 [0.20, 5.04]
2.24 Respiratory function: forced expiratory volume at 1 second (FEV1) (Liters)	12	362	Mean Difference (IV, Random, 95% CI)	0.04 [-0.06, 0.14]

Analysis 2.1. Comparison 2: IMT vs control/sham, Outcome 1: Dyspnea: Borg (at submaximal exercise capacity)




Footnotes

- (1) Isotime at 70%-80% of Wmax
- (2) 50% of Wmax
- (3) Isotime at 75% of Wmax
- (4) at 25%-50% of Wmax
- (5) 75% of Wmax

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.2. Comparison 2: IMT vs control/sham, Outcome 2: Dyspnea: Baseline and Transition Dyspnea Indexes (BDI-TDI)




Footnotes

- (1) Threshold device
- (2) Pflax device
- (3) This study did not report the BDI score.

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

**Analysis 2.3. Comparison 2: IMT vs control/sham, Outcome 3: Dyspnea: Transition
Dyspnea Index (TDI): Focal score (subgroup analysis: with or without respiratory muscle weakness)**




Footnotes

- (1) Threshold device
- (2) Pflex device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.4. Comparison 2: IMT vs control/sham, Outcome 4: Dyspnea: Modified Medical Research Council (mMRC)




Footnotes

- (1) The scale is from 1 to 5

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

**Analysis 2.5. Comparison 2: IMT vs control/sham, Outcome 5:
Functional exercise capacity: 6-minute walk distance (6MWD) (meters)**




Footnotes

- (1) Incentive spirometer
- (2) Threshold device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.6. Comparison 2: IMT vs control/sham, Outcome 6: Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: duration of the intervention)




Footnotes

- (1) Incentive spirometer
- (2) Threshold device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.7. Comparison 2: IMT vs control/sham, Outcome 7: Functional exercise capacity: 6-minute walk distance (6MWD) (meters) (subgroup analysis: with or without respiratory muscle weakness)




Footnotes

(1) Incentive spirometer


Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.8. Comparison 2: IMT vs control/sham, Outcome 8: Functional exercise capacity: 12-minute walk distance (12MWD) (meters)



Analysis 2.9. Comparison 2: IMT vs control/sham, Outcome 9: Functional exercise capacity: Wmax (watt)




Footnotes

- (1) Threshold device
- (2) Pfllex device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 2.10. Comparison 2: IMT vs control/sham, Outcome 10: Functional exercise capacity: exercise time (seconds)




Footnotes

- (1) Cycling at 75% of Wmax
- (2) Cycling at 50% of Wmax
- (3) Exercise time on treadmill which was set to 80% of the inclination and to 100% of the speed reached at VO₂peak
- (4) Incremental cycle test (Threshold device)
- (5) Incremental cycle test (Pflflex device)

Analysis 2.11. Comparison 2: IMT vs control/sham, Outcome 11: Functional exercise capacity: shuttle walk test (SWT) (meters)




Analysis 2.12. Comparison 2: IMT vs control/sham, Outcome 12: Health-related quality of life (HRQoL): St George Respiratory Questionnaire (SGRQ)



Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.13. Comparison 2: IMT vs control/sham, Outcome 13: Health-related quality of life (HRQoL): Chronic Respiratory Disease Questionnaire (CRQ)




Footnotes

- (1) Pflx device
- (2) Threshold device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 2.14. Comparison 2: IMT vs control/sham, Outcome 14: Health-related quality of life (HRQoL): COPD Assessment Test (CAT)



Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.15. Comparison 2: IMT vs control/sham, Outcome 15: Inspiratory muscle strength: PImax (cmH₂O)




Footnotes

- (1) Incentive spirometer
- (2) Threshold device
- (3) Pflflex device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.16. Comparison 2: IMT vs control/sham, Outcome 16: Inspiratory muscle strength: P1max (cmH₂O) (subgroup analysis: duration of the intervention)




Footnotes

- (1) Threshold device
- (2) Incentive spirometer
- (3) Pflflex device

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias

Analysis 2.17. Comparison 2: IMT vs control/sham, Outcome 17: Inspiratory muscle strength: P_{lmax} (cmH₂O) (subgroup analysis: with or without respiratory muscle weakness)




Footnotes

- (1) Pfler device
- (2) Threshold device
- (3) Incentive spirometer

Risk of bias legend

- (A) Bias arising from the randomization process
- (B) Bias due to deviations from intended interventions
- (C) Bias due to missing outcome data
- (D) Bias in measurement of the outcome
- (E) Bias in selection of the reported result
- (F) Overall bias


Analysis 2.18. Comparison 2: IMT vs control/sham, Outcome 18: Inspiratory muscle strength: P_{lmax} (cmH₂O) (subgroup analysis: method of measurement)



Footnotes
 (1) Incentive spirometer
 (2) Threshold device
 (3) Pflex device

Risk of bias legend
 (A) Bias arising from the randomization process
 (B) Bias due to deviations from intended interventions
 (C) Bias due to missing outcome data
 (D) Bias in measurement of the outcome
 (E) Bias in selection of the reported result
 (F) Overall bias


Analysis 2.19. Comparison 2: IMT vs control/sham, Outcome 19: Laboratory exercise test: VO₂ peak




Footnotes

- (1) Pflflex device
- (2) Threshold device

Analysis 2.20. Comparison 2: IMT vs control/sham, Outcome 20: Respiratory muscle endurance: respiratory muscle endurance pressure (P_{thmax}) (cmH₂O)




Analysis 2.21. Comparison 2: IMT vs control/sham, Outcome 21: Respiratory muscle endurance time: T_{lim} (seconds)




Footnotes

- (1) Breathing against 66% of P_Imax
- (2) Breathing against ≈80% of P_Imax
- (3) Breathing against 70% of P_Imax (resistive device) Incentive spirometer
- (4) Breathing against 70% of P_Imax (Threshold device)
- (5) Breathing against 50-60% of P_Imax
- (6) Breathing against 50% of Poesmax
- (7) Breathing against 60% of P_Imax
- (8) Breathing against 80% of P_Imax
- (9) Sustained ventilation at 66% or 75% of MVV

Analysis 2.22. Comparison 2: IMT vs control/sham, Outcome 22: Maximal voluntary ventilation (MVV)




**Analysis 2.23. Comparison 2: IMT vs control/sham, Outcome 23:
Respiratory function: forced expiratory volume at 1 second (FEV1) (%pred)**



Footnotes

- (1) Threshold device
- (2) Pflex device

**Analysis 2.24. Comparison 2: IMT vs control/sham, Outcome 24:
Respiratory function: forced expiratory volume at 1 second (FEV1) (Liters)**



Footnotes

- (1) Threshold device
- (2) Pflex device

APPENDICES

Appendix 1. Database search strategies

Database/search platform/date of last search	Search strategy	Results
Airways Register (via Cochrane Register of Studies) Date of most recent search: 19 October 2022	1 MeSH DESCRIPTOR Pulmonary Disease, Chronic Obstructive Explode All AND INSEGMENT	Oct 2020=937
	2 MeSH DESCRIPTOR Bronchitis, Chronic AND INSEGMENT 3 (obstruct*) near3 (pulmonary or lung* or airway* or airflow* or bronch* or respirat*) AND INSEGMENT	Aug 2021=44
	4 COPD:MISC1 AND INSEGME	Oct 2022=265 (together with CENTRAL)
	5 (COPD OR AECOPD):TI,AB,KW AND INSEGMENT	
	6 #1 OR #2 OR #3 OR #4 OR #5 AND INSEGMENT	
	7 MESH DESCRIPTOR Breathing Exercises AND INSEGMENT	
	8 MESH DESCRIPTOR Respiratory Muscles EXPLODE ALL AND INSEGMENT	
	9 MESH DESCRIPTOR Exercise Tolerance AND INSEGMENT	
	10 (IMT or RMT):ti,ab AND INSEGMENT	
	11 ((inspiratory or ventilat* or respiratory) NEAR3 (muscle or resistance) NEAR3 (train* or strength* or endur*)) AND INSEGMENT	
	12 (threshold NEAR2 (load or device*)) AND INSEGMENT	
	13 (resist* NEAR2 device*) AND INSEGMENT	
	14 isocapnic hyperpnea AND INSEGMENT	
	15 #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #14 OR #13	
	16 #6 AND #15	
	17 INREGISTER	
	18 #16 AND #17	
	CENTRAL (via Cochrane Register of Studies) Date of most recent search: 19 October 2022	
2 MeSH DESCRIPTOR Bronchitis, Chronic AND CENTRAL:TARGET		Aug 2021=69
3 (obstruct*) near3 (pulmonary or lung* or airway* or airflow* or bronch* or respirat*) AND CENTRAL:TARGET		
4 COPD:MISC1 AND CENTRAL:TARGET		
5 (COPD OR AECOPD):TI,AB,KW AND CENTRAL:TARGET		
6 #1 OR #2 OR #3 OR #4 OR #5 AND CENTRAL:TARGET		
7 MESH DESCRIPTOR Breathing Exercises AND CENTRAL:TARGET		
8 MESH DESCRIPTOR Respiratory Muscles EXPLODE ALL AND CENTRAL:TARGET		
9 MESH DESCRIPTOR Exercise Tolerance AND CENTRAL:TARGET		
10 (IMT or RMT):ti,ab AND CENTRAL:TARGET		
11 ((inspiratory or ventilat* or respiratory) NEAR3 (muscle or resistance) NEAR3 (train* or strength* or endur*)) AND CENTRAL:TARGET		
12 (threshold NEAR2 (load or device*)) AND CENTRAL:TARGET		
13 (resist* NEAR2 device*) AND CENTRAL:TARGET		
14 isocapnic hyperpnea AND CENTRAL:TARGET		
15 #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #14 OR #13 AND CENTRAL:TARGET		
16 #6 AND #15 AND CENTRAL:TARGET		

(Continued)

- 17 CENTRAL:TARGET
18 #16 AND #17 AND CENTRAL:TARGET

MEDLINE (Ovid) ALL Date of most recent search: 20 October 2022	1 Lung Diseases, Obstructive/	Oct 2020=1609
	2 exp Pulmonary Disease, Chronic Obstructive/	
	3 emphysema\$.tw.	
	4 (chronic\$ adj3 bronchiti\$).tw.	
	5 (obstruct\$ adj3 (pulmonary or lung\$ or airway\$ or airflow\$ or bronch\$ or respirat\$)).tw.	Aug 2021=86
	6 (COPD or AECOPD or AECB).ti,ab.	
	7 or/1-6	Oct 2022=172
	8 Breathing Exercises/	
	9 exp Respiratory Muscles/	
	10 Exercise Tolerance/	
	11 Muscle Strength/	
	12 (IMT or RMT).ti,ab.	
	13 ((inspiratory or ventilat\$ or respiratory) adj3 (muscle or resistance) adj3 (train\$ or strength\$ or endur\$)).tw.	
	14 (threshold adj2 (load or device\$)).tw.	
	15 (resist\$ adj2 device\$).tw.	
	16 isocapnic hyperpnea.tw.	
	17 or/8-16	
	18 (controlled clinical trial or randomized controlled trial).pt.	
	19 (randomized or randomised).ab,ti.	
	20 placebo.ab,ti.	
	21 randomly.ab,ti.	
	22 trial.ab,ti.	
	23 groups.ab,ti.	
	24 or/18-23	
	25 Animals/	
	26 Humans/	
	27 25 not (25 and 26)	
	28 24 not 27	
	29 7 and 17 and 28	

Embase (Ovid) Date of most recent search: 20 October 2022	1 Chronic Obstructive Lung Disease/	Oct 2020=2093
	2 Obstructive Airway Disease/	
	3 Chronic Bronchitis/	
	4 Lung Emphysema/	
	5 emphysema\$.tw.	Aug 2021=146
	6 (chronic\$ adj3 bronchiti\$).tw.	
	7 (obstruct\$ adj3 (pulmonary or lung\$ or airway\$ or airflow\$ or bronch\$ or respirat\$)).tw.	Oct 2022=251
	8 (COPD or AECOPD or AECB).ti,ab.	
	9 or/1-8	
	10 breathing exercise/	
	11 breathing muscle/	
	12 exercise tolerance/	
	13 muscle strength/	
	14 (IMT or RMT).ti,ab.	
	15 ((inspiratory or ventilat\$ or respiratory) adj3 (muscle or resistance) adj3 (train\$ or strength\$ or endur\$)).tw.	
	16 (threshold adj2 (load or device\$)).tw.	
	17 (resist\$ adj2 device\$).tw.	
	18 isocapnic hyperpnea.tw.	
	19 or/10-18	
	20 Randomized Controlled Trial/	
	21 randomization/	
	22 controlled clinical trial/	
	23 Double Blind Procedure/	

(Continued)

- 24 Single Blind Procedure/
- 25 Crossover Procedure/
- 26 (clinica\$ adj3 trial\$).tw.
- 27 ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj3 (mask\$ or blind\$ or method\$)).tw.
- 28 exp Placebo/
- 29 placebo\$.ti,ab.
- 30 random\$.ti,ab.
- 31 ((control\$ or prospectiv\$) adj3 (trial\$ or method\$ or stud\$)).tw.
- 32 (crossover\$ or cross-over\$).ti,ab.
- 33 or/20-32
- 34 exp animals/ or exp invertebrate/ or animal experiment/ or animal model/ or animal tissue/ or animal cell/ or nonhuman/
- 35 human/ or normal human/ or human cell/
- 36 34 and 35
- 37 34 not 36
- 38 33 not 37
- 39 9 and 19 and 38

CINHAL (EBSCO)	S41 S7 AND S17 AND S40	Oct 2020=182
Date of most recent search: 19 October 2022	S40 S39 NOT S38	
	S39 S18 OR S19 OR S20 OR S21 OR S22 OR S23 OR S24 OR S25 OR S26 OR S27 OR S28 OR S29 OR S30 OR S31 OR S32	Aug 2021=22
	S38 S36 NOT S37	
	S37 (MH "Human")	
	S36 S33 OR S34 OR S35	Oct 2022=76
	S35 TI (animal model*)	
	S34 (MH "Animal Studies")	
	S33 (MH "Animals+")	
	S32 AB (cluster W3 RCT)	
	S31 MH (crossover design) OR MH (comparative studies)	
	S30 AB (control W5 group)	
	S29 PT (Randomized Controlled Trial)	
	S28 (MH "Placebos")	
	S27 MH ("sample size") AND AB (assigned OR allocated OR control)	
	S26 TI (trial)	
	S25 AB (random*)	
	S24 TI (randomised OR randomized)	
	S23 (MH "Cluster Sample")	
	S22 (MH "Pretest-Posttest Design")	
	S21 (MH "Random Assignment")	
	S20 (MH "Single-Blind Studies")	
	S19 (MH "Double-Blind Studies")	
	S18 (MH "Randomized Controlled Trials")	
	S17 S8 OR S9 OR S10 OR S11 OR S12 OR S13 OR S14 OR S15 OR S16	
	S16 isocapnic hyperpnea	
	S15 resist* N2 device*	
	S14 threshold* N2 (load or device*)	
	S13 (inspiratory or ventilat* or respiratory) N3 (muscle or resistance) N3 (train* or strength* or endur*)	
	S12 AB (IMT or RMT) or TI (IMT or RMT)	
	S11 (MH "Muscle Strength")	
	S10 (MH "Exercise Tolerance")	
	S9 (MH "Respiratory Muscles+")	
	S8 (MH "Breathing Exercises")	
	S7 S1 or S2 or S3 or S4 or S5 or S6	
	S6 COPD or COAD or COBD or AECD	
	S5 (pulmonary or lung* or airway* or airflow* or bronch* or respirat*) and (obstruct*) and (disease*)	
	S4 chronic bronchitis	
	S3 "emphysema**"	

(Continued)

 S2 (MH "Lung Diseases, Obstructive")
 S1 (MH "Pulmonary Disease, Chronic Obstructive+")

PsycINFO (Ovid)	1 exp Chronic Obstructive Pulmonary Disease/	Oct 2020=10
Date of most recent search: 19 October 2022	2 emphysema\$.tw.	August 2021=1
	3 (chronic\$ adj3 bronchiti\$).tw.	
	4 (obstruct\$ adj3 (pulmonary or lung\$ or airway\$ or airflow\$ or bronch\$ or respirat\$)).tw.	Oct 2022=3
	5 (COPD or AECOPD or AECB).ti,ab.	
	6 or/1-5	
	7 ((inspiratory or ventilat\$ or respiratory) adj3 (muscle or resistance) adj3 (train\$ or strength\$ or endur\$)).tw.	
	8 (IMT or RMT).ti,ab.	
	9 (threshold adj2 (load or device\$)).tw.	
	10 (resist\$ adj2 device\$).tw.	
	11 isocapnic hyperpnea.tw.	
	12 or/7-11	
	13 random\$.tw.	
	14 (clinical adj5 trial\$).tw.	
	15 (control\$ adj5 trial\$).tw.	
	16 ((clinical or control\$ or comparativ\$) adj5 (study or studies)).tw.	
	17 placebo\$.tw.	
	18 (single blind\$ or single-blind\$).tw.	
	19 (double blind\$ or double-blind\$).tw.	
	20 (triple blind\$ or triple-blind\$).tw.	
	21 or/13-20	
	6 and 12 and 21	
PEDro	Abstract & title: inspiratory muscle copd random*	Oct 2020=81
Date of most recent search: 20 October 2022	Topic: chronic respiratory disease	Aug 2021=1
	Method: Clinical trial	
	When searching: Match all search terms (AND)	Oct 2022=5
ClinicalTrials.gov	Study type: Interventional	Oct 2020=58
	Condition: COPD	
Date of most recent search: 20 October 2022	Intervention: inspiratory muscle training OR threshold load OR threshold device OR resistive device	Aug 2021=6

(Continued)

Oct 2022=12

WHO trials portal Date of most recent search: 20 October 2022	Condition: COPD Intervention: inspiratory muscle training OR threshold load OR threshold device OR resistive device	Oct 2020=347
---	--	--------------

Aug 2021=2

Oct 2022=9

WHAT'S NEW

Date	Event	Description
9 January 2023	Amended	Search date corrected in Abstract and PLS to be 20 October 2022

HISTORY

Protocol first published: Issue 11, 2020

Review first published: Issue 1, 2023

CONTRIBUTIONS OF AUTHORS

Conception of the review: OA/WF

Co-ordination of the review: OA

Search and collection of studies for inclusion in the review: OA/WF

Data extraction: OA/AK

Assessment of the risk of bias in the included studies: OA/TL

Analysis of data: OA/AK

Assessment of certainty in the body of evidence: OA/TL

Interpretation of data: WF/SK

Writing the review: OA

Clinical and statistical comments: RG/AR

Contributions of editorial team

Sally Spencer (Co-ordinating Editor) edited the review; advised on methodology, interpretation and content; and approved the review prior to publication.

Rebecca Fortescue (Co-ordinating Editor) checked the data in the review.

Anne Holland (Contact Editor): edited the review; and advised on methodology, interpretation and content.

Emma Dennett (Deputy Co-ordinating Editor): advised on methodology, interpretation and content; and edited the review.

Emma Jackson (Managing Editor): co-ordinated the editorial process; conducted peer review; obtained translations; and edited the plain language summary and reference sections of the review.

Elizabeth Stovold (Information Specialist): designed the search strategy; ran the searches; and edited the search methods section.

Vittoria Lutje (Freelance Information Specialist): ran the updated search.

Kayleigh Kew (Freelance Editor): edited the review and advised on methodology.

DECLARATIONS OF INTEREST

OA: none known

RG: none known

AK: none known

WF: none known

Inspiratory muscle training, with or without concomitant pulmonary rehabilitation, for chronic obstructive pulmonary disease (COPD) (Review)

231

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

TL: none known
AR: none known
SK: none known

SOURCES OF SUPPORT

Internal sources

- University of Sfax, Faculty of Medicine, Tunisia

Omar Ammous is supported by the University of Sfax, Faculty of Medicine, to conduct this systematic review as his medical thesis.

External sources

- National Institute for Health and Care Research (NIHR), UK

Provision of Cochrane Infrastructure funding to Cochrane Airways

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Because few studies used the Borg scale and reported it at different levels of effort, we decided to replace it with P_Imax for subgroup analysis. The 6MWD and P_Imax are respectively clinical and physiological outcomes, and we believed that analysing both outcomes together would present an opportunity to check for the possible correlation between them.

Because many RCTs were included, we conducted a sensitivity analysis by keeping just the studies at an overall low risk of bias.

We conducted a sensitivity analysis by removing (Preusser 1994) because we noticed large differences in baseline data between the two groups.

We did not use Robvis to generate weighted bar plots because it was possible to create them with Revman Web, and finally we kept just the PRISMA flow diagram and funnel plots.

INDEX TERMS

Medical Subject Headings (MeSH)

*Breathing Exercises; Dyspnea [rehabilitation]; Muscles; *Physical Therapy Modalities; *Pulmonary Disease, Chronic Obstructive [therapy]; Quality of Life

MeSH check words

Humans