

ORIGINAL ARTICLE



# Does Inspiratory Muscle Training Affect Respiratory Parameters? Different Effects of Diaphragm Muscle Focused Respiratory Exercise

<sup>1</sup>Mehmet Vural , <sup>1</sup>Mustafa Ozdal , <sup>2</sup>M. Berk Demiryol \*, <sup>3</sup>C. Didem Eyipinar

<sup>1</sup>Coaching Education Department, Faculty of Sport Science, Gaziantep University, Gaziantep, Turkey.

<sup>2</sup>Sport Management Department, Faculty of Sport Science, Gaziantep University, Gaziantep, Turkey.

<sup>3</sup>Physical Education and Sports Teaching Department, Faculty of Sport Science, Gaziantep University, Gaziantep, Turkey.

Submitted April 17, 2024; Accepted in final form June 21, 2024.

## ABSTRACT

**Background.** Considering that breathing strategies are important for athletic performance, there isn't enough scientific data available to fully understand how they operate just yet. **Objectives.** This study aimed to investigate the effect of conscious diaphragm muscle-based respiratory muscle exercise on respiratory muscle strength and function. **Methods.** For that, four groups (IMT-DF (inspiratory muscle training using diaphragm), IMT (inspiratory muscle training), IMT-SHAM (training without intensity) and control) were formed in which 24 healthy sedentary participants were randomly divided. Inspiratory muscle training was organized as three days a week for 4 weeks and measurements of respiratory muscle strength (measured with a pressure gauge) and function (measured with a spirometer) were obtained one day before and one day after the 4-week period. The data obtained were normally distributed and one-way ANOVA analysis of variance and Paired sample t-test were applied for statistical analysis. In respiratory muscle strength parameters maximal inspiratory pressure and maximal expiratory pressure (MIP-MEP), significant differences were found in favor of IMT-DF and IMT groups. **Results.** Significance was found in favor of IMT-DF in the analysis of the difference between means and percentage change in the comparison of pre-post test data between groups. In the pulmonary function tests, a significant difference was found between the IMT-DF group and the IMT group in favor of the IMT-DF group in the forced expiratory volume in the first second (FEV1) parameter; a significant difference was found between the IMT-SHAM group and the IMT-DF group in favor of the IMT-SHAM group and no significant difference was detected in respiratory functions ( $p>0.05$ ). **Conclusion.** As a result, performed respiratory muscle training specific to the diaphragm muscle may provide more positive improvements in respiratory muscle strength and function compared to normal training.

**KEYWORDS:** *Diaphragm, Respiratory, Exercise.*

## INTRODUCTION

Respiration is an activity needed every second of life for every living creature to carry out its vital activities in a healthy way. When a person breathes, oxygen, the basic element that cells need, reaches the cells and the

necessary need is met. In the process, oxygen passes through various filtration processes, binds to hemoglobin in the blood, and is delivered to the appropriate organ, tissue, and cell (1, 2).

\*. Corresponding Author:

M. Berk Demiryol, Ph.D.

E-mail: [demiryolmete@gmail.com](mailto:demiryolmete@gmail.com)

Almost all people do not pay attention to all the physiological processes which mentioned before, do not try to control them and the process continues as it is. Because it is known that this involves an autonomous process and that the respiration of the organism takes place smoothly and completely according to its own coding (3, 4). This is followed by a number of exercises and health practices aimed at improving the respiratory performance of individuals (5, 6). Here, the implementation of respiratory muscle-strengthening exercise programs for sedentary people and athletes comes to the fore looking at the literature, it is possible to find many studies in which respiratory muscles were strengthened and contributed to the improvement of muscle strength and respiratory function (5, 7, 8). Sometimes, studies have been carried out on a group of patients to overcome the respiratory difficulties and difficulties caused by the disease, and sometimes, research has been designed to achieve better capacities both in normal life (9, 10) and in sports performance (5), focusing on improving the existing healthy capacity to better levels. However, when the studies are carefully examined, there is not much attention or detailed explanation of how respiratory muscle training is done outside the protocol (8, 11-14).

Although many benefits are obtained from respiratory muscle training (6, 11), there is a detail that is not emphasized (how to perform breathing exercises mechanically). When training for the respiratory muscle was performed, it was determined that the diaphragm muscle was involuntary activated and loaded while inhaling and exhaling during the training (15). People are only given an inspiratory muscle training device and told how the training protocol will be on it and how long the rest will be (16, 17). We designed that muscle-focused training should be prepared so that the diaphragm muscle can perform stronger and more effective training and the capacity of this muscle, which is mainly responsible for breathing, can be increased at a more effective level. Based on this information, respiratory muscle training was carried out in addition to a working process for consciously starting and maintaining breathing in the diaphragm area by activating the respiratory muscle (diaphragm). Our hypothesis is that conscious respiratory

muscle training for the diaphragm muscle will affect respiratory function and muscle strength more positively. The aim is to reach a conscious level in the use of the diaphragm and to achieve more and more effective results in respiratory muscle training. In this direction, our study is important both to eliminate such an important deficiency in the literature and to carry out training programs in a more conscious state in future research on respiratory muscles.

## MATERIALS AND METHODS

**Experimental Design and Participants.** Our study was designed as a randomized controlled pre-posttest experimental design. Participants visited the research site a total of 2 times and underwent a 4-week training period. Three groups (IMT-SHAM, IMT, IMT-DF) performed respiratory muscle exercises to the accompaniment of researchers, 2 sets of 30 repetitions (1 min rest between repetitions) 3 days a week for each week during this 4-week training period. Participants were assigned to study groups according to the full randomization. Prior to the 4-week training period (first visit), participants' respiratory muscle strength maximal inspiratory pressure (MIP), and maximal expiratory pressure (MEP) parameters and respiratory functions were measured, and other anthropometric characteristics of the participants were recorded (Table 1). The intensity of the training sessions was based on 40% of the participants' MIP values. A-priori power analysis of the study was calculated using the G\*Power 3.1.9.7 program. As a result of the calculation performed using the research data with a total sample size of 24 (The sample size was estimated on the basis of data published by Kim, Park & Lee (18)), the effect size value of the research was found to be 0. With a 5% error rate ( $\alpha=0.05$ ), the power of the study ( $1-\beta$ ) was calculated as 0.95. A total of 24 healthy sedentary males, who did not have any respiratory disease, did not use medication, did not smoke or consume alcohol, and no leisure-time physical activity, with a mean age of  $19.63\pm 1.173$  years, a mean height of  $177.25\pm 3.90$  centimeters, mean weight of  $74\pm 4.59$  kilograms and a mean body mass index of  $23.6\pm 0.68$  kilograms/square meter, participated in our study. Inclusion criteria are not having any respiratory disease, not smoking or drinking alcohol, and not using any medication regularly. No participant withdrew from the study.

**Table 1. Descriptive parameters of subjects (n=24)**

	Minimum (n=24)	Maximum (n=24)	Mean±Std.Dev (n=24)
Age (Years)	18	21	19.63±1.173
Height (cm)	175	182	177.25±3.90
Weight (kg)	69	80	74±4.59
BMI (kg/m <sup>2</sup> )	22.5	24.2	23.6±0.68

BMI: Body Mass Index.

Participants were divided into 4 groups, with 6 people in each group. In addition, the participants randomly assigned to the IMT-DF group (n=6) were shown breathing exercises focused on that area in order to perform conscious respiratory muscle training for the diaphragm muscle, and the participants were asked to perform the training by paying attention to the specified points. The other groups were asked to perform respiratory muscle training without any intervention (the control group (n=6) was not subjected to any training). IMT-SHAM application group (n=6) was trained

without any exercise intensity (at %0 of MIP), IMT group (n=6) was only informed about the repetition and rest periods of the respiratory muscle training and performed the training for 4 weeks (19, 20). At the end of the 4-week training period, the respiratory muscle strength and function of the participants were measured as the final test (Table 2; Table 3). The necessary permission was obtained from the Gaziantep University Social and Human Sciences Ethics Committee for the realization of the study (Protocol number: 361976).

**Table 2. Descriptive Statistics of Pre-test and Post-test Respiratory Muscle Strength Values of Groups**

Groups (n=24)		MIP (cmH <sub>2</sub> O)		MEP (cmH <sub>2</sub> O)	
		$\bar{x}$	SD	$\bar{x}$	SD
IMT-DF (n=6)	Pre	61.83	21.55	83.83	27.68
	Post	108.50	13.46	125.83	21.78
IMT (n=6)	Pre	87.17	18.51	118.83	29.61
	Post	100.33	22.97	131.67	27.12
IMT-SHAM (n=6)	Pre	76.00	14.21	103.33	25.34
	Post	79.67	13.94	110.33	28.44
Control (n=6)	Pre	81.33	12.97	127.83	11.0
	Post	114.33	22.04	132.50	22.46

MIP: Maximal Inspiratory Pressure; MEP: Maximal Expiratory Pressure; IMT-DF: diaphragm focused inspiratory muscle training; IMT: Inspiratory Muscle Training; IMT-SHAM: Inspiratory Muscle Training-Sham; Control: Non-Inspiratory Muscle Training.

**Table 3. Descriptive Statistics of Pre-test and Post-test Respiratory Function Values of the Groups**

Groups (n=24)		FEV (lt)		FVC (lt)		MVV (lt/min)		VC (lt)		TV (lt)		IVC (lt)	
		$\bar{x}$	SD	$\bar{x}$	SD	$\bar{x}$	SD	$\bar{x}$	SD	$\bar{x}$	SD	$\bar{x}$	SD
IMT-DF (n=6)	Pre	3.23	0.98	4.19	1.29	112.98	34.45	4.48	2.02	0.60	0.19	2.61	1.09
	Post	3.87	0.68	4.89	0.80	135.58	23.73	6.09	0.77	0.87	0.51	2.94	0.50
IMT (n=6)	Pre	3.70	0.44	4.49	0.62	129.43	15.46	3.97	1.13	0.83	0.31	2.40	0.43
	Post	4.54	0.98	4.99	0.99	160.32	31.50	5.26	0.42	1.26	0.46	2.93	0.42
IMT-SHAM (n=6)	Pre	2.60	1.19	3.98	1.14	112.05	30.52	2.39	1.13	0.64	0.14	1.53	0.62
	Post	3.06	1.02	4.33	0.43	125.13	16.17	3.79	1.17	0.75	0.35	2.03	0.51
Control (n=6)	Pre	4.23	0.25	4.41	0.32	142.88	15.48	4.67	1.06	0.69	0.40	2.42	0.78
	Post	3.88	0.52	4.30	0.89	135.93	18.23	4.62	1.06	1.08	0.52	2.67	0.60

FEV: Forced Expiratory Volume; FVC: Forced Vital Capacity; MVV: Maximal Voluntary Ventilation; VC: Vital Capacity; TV: Tidal Volume; IVC: Inspiratory Vital Capacity.

**Training Protocol and Data Measurement Tools.** The groups performed the inspiratory muscle training program for 3 days a week for a

month. The participants breathed 30\*2 times at 40% MIP (1-min rest between repetitions) using an inspiratory muscle training device. For the IMT-

DF and IMT groups, MIP was measured on the first training day of every week. The same procedure was used in the control group; however, the intensity was at 0% MIP with IMT-SHAM (6). At every inspiratory muscle training session, oxygen saturation was followed by pulse oximetry to avoid a hypoxic condition in participants.

For the diaphragm-focused group, diaphragm-strengthening exercise methods were shown before the study in order to perform breathing more consciously. While doing respiratory training, the diaphragm muscle was made to contract consciously and the training was carried out in that way. Here, the conscious initiation of breathing through the diaphragm muscle and not through the rib cage was an important factor.

**MIP-MEP Measurement.** The MIP and MEP were measured using an electronic pulmonary pressure meter (Pocket Spiro MPM-100, Medical Electronic Construction R&D, Brussels, Belgium) in accordance with the American Thoracic Society and European Pulmonary Society recommendations from 2002. Subjects did maximal expiration first, followed by maximal inspiration for 1-3 seconds. Subjects completed maximal inspiration for MEP measurement before being asked to execute maximal expiration for 1-3 seconds. The lab tests were taken again till the difference between the best two values reached 5 cmH<sub>2</sub>O, at which point the best result was noted in cmH<sub>2</sub>O (16).

**Respiratory Function Measurement.** The pulmonary function was measured using a spirometer (Pocket Spiro USB-100, Medical Electronic Construction R&D, Brussels, Belgium). Slow and forced vital capacity tests were selected and measured according to the 2002 guidelines of the American Thoracic Society and European Respiratory Society. Variables such as forced vital capacity (FVC), forced expiratory volume in one second (FEV<sub>1</sub>), maximal voluntary ventilation (MVV), vital capacity (VC), tidal volume (TV), inspiratory vital capacity (IVC) were recorded using pulmonary function test (20). Subjects performed twice all of the respiratory function measurements, and we have recorded the measurements with higher data.

**Statistical Analysis.** SPSS 22.0 (SPSS Inc., Chicago, IL, USA) program was used for statistical analysis. Values were represented as minimum, maximum, mean, standard deviation, and significance was set at 0.05. Shapiro–Wilk test was performed to assess normality, and One Way

ANOVA and LSD significant test were performed for differences of pre-posttest (intergroup), and percentage differences between groups. Paired Sample T-test was used for pre and post-tests.

## RESULTS

In Table 4, the analysis of statistical changes in respiratory muscle strength in different breathing practice groups is presented. The data obtained from the practice groups were compared before and after the 4-week training period. When the pre-post test data of each group were analyzed, a significant difference was found in favor of the post-test for the IMT-DF group ( $t=-6.019$ ,  $p=0.002$ ), IMT-SHAM group ( $t=-4.825$ ,  $p=0.005$ ) and MIP in the control group ( $t=4.807$ ,  $p=0.005$ ) ( $p<0.05$ ). When a comparison was made for the MEP parameter of the same groups, a significant difference was found in favor of the post-test only in the IMT-DF group ( $t=-4.376$ ,  $p=0.007$ ,  $p<0.05$ ). When the difference between the pre-test and post-test data was compared between the groups, a significant difference was found between the IMT-DF group and other groups for the MIP parameter ( $F=4.242$ ,  $p=0.018$ ,  $p<0.05$ ). The significant difference was found to be in favor of the IMT-DF groups. A significant difference was detected in the comparison of the difference of pre-post test data for MEP between groups ( $F=3.546$ ,  $p=0.033$ ,  $p<0.05$ ). This significant difference was found to be between the IMT-DF group and the IMT group in favor of the IMT-DF group. When the change between the pre and post-test data of the groups was analyzed in percentage, a significant difference was found in the MIP parameter ( $F=4.180$ ,  $p=0.019$ ,  $p<0.05$ ). The significant difference was between the IMT-DF group, SHAM group, and control group and resulted in favor of the IMT-DF group. In the MEP parameter, when the percentage change between the pre-post test data was analyzed, a significant difference was found ( $F=5.028$ ,  $p=0.009$ ,  $p<0.05$ ). The significant difference was between the IMT-DF group and IMT group, SHAM group, and control group and resulted in favor of the IMT-DF group.

In Table 5, statistical analysis of the changes in pulmonary function in different breathing treatment groups is demonstrated. When the data were examined, in the pre-post test results of the groups, a significant difference was found in the IMT group in favor of the post-test in the averages of FEV<sub>1</sub> (lt) ( $t=-2.790$ ,  $p=0.038$ ), MVV (lt/min) ( $t=-3.368$ ,  $p=0.020$ ) and VC (lt) ( $t=-3.183$ ,


p=0.024) (p<0.05). In the IMT-SHAM group, a significant difference was found in the mean VC (lt) in favor of the post-test (t=-3.261, p=0.022, p<0.05). In the control group, a significant difference was found in the mean TV (lt) in favor of the post-test (t=-4.094, p=0.009, p<0.05). In the comparison of the difference between the pre-post test data in pulmonary function parameters between the groups, a significant difference was found between the IMT-DF group and the IMT

group in favor of the IMT-DF group in the FEV1 parameter, and between the IMT group and the control group in favor of the IMT group (F=3.546, p=0.033, p<0.05); When the change between the pre-test and post-test data of the groups was analyzed in percentage, no significant difference was detected in respiratory functions (p>0.05). The changes in other respiratory parameters for the groups are shown in Figure 1.

**Table 4. Analysis of Changes in Respiratory Muscle Strength Parameters**

		IMT-DF (n=6)	IMT (n=6)	IMT-SHAM (n=6)	Control (n=6)
<b>MIP (cmH<sub>2</sub>O)</b>	Pre-test	61.83±21.55	87.17±18.51	76.00±14.21	81.33±12.97
	Post-test	108.50±13.46 <sup>a</sup>	100.33±22.97	79.67±13.94 <sup>a</sup>	114.33±22.04 <sup>a</sup>
	Diff. (lt)	42.00±23.51 <sup>bcd</sup>	12.83±26.21	7.00±12.05	4.67±17.34
	Diff. (%)	98.28±88.76 <sup>cd</sup>	19.37±35.86	5.08±3.15	41.54±21.69
<b>MEP (cmH<sub>2</sub>O)</b>	Pre-test	83.83±27.68	118.83±29.61	103.33±25.34	127.83±11.00
	Post-test	125.83±21.78 <sup>a</sup>	131.67±27.12	110.33±28.44	132.50±22.46
	Diff. (lt)	0.64±0.68 <sup>b</sup>	0.84±0.74 <sup>b</sup>	0.46±0.72	-0.35±0.57
	Diff. (%)	61.50±50.78 <sup>bcd</sup>	13.78±24.61	6.78±11.93	3.51±13.16

a: significant difference between pre-post test; b: significant difference with IMT; c: Significant difference with IMT-SHAM; d: significant difference with control group; MIP: maximal inspiratory pressure; MEP: maximal expiratory pressure; IMT-DF: intervention group 1 (a group that received inspiratory muscle training using diaphragm); IMT: intervention group 2 (a group that received chest-focused inspiratory muscle training); IMT-SHAM: intervention group 3 (a group that received without inspiratory muscle training); Control: control group 4 (a group without inspiratory muscle training).



**Figure 1.** The Percentage Analysis of Change in Respiratory Parameters of Groups.

Table 5. Analysis of Changes in Respiratory Functions

		IMT-DF (n=6)	IMT (n=6)	IMT-SHAM (n=6)	Control (n=6)
FEV1 (lt)	Pre-test	3.23±0.98	3.70±0.44	2.60±1.19	4.23±0.25
	Post-test	3.87±0.68	4.54±0.98 <sup>a</sup>	3.06±1.02	3.88±0.52
	Diff. (lt)	0.64±0.68 <sup>b</sup>	0.84±0.74 <sup>d</sup>	0.46±0.72	-0.35±0.57
	Diff. (%)	25.66±29.11	22.35±20.45	23.55±31.29	-8.05±13.48
FVC (lt)	Pre-test	4.19±1.29	4.49±0.62	3.98±1.14	4.41±0.32
	Post-test	4.89±0.80	4.99±0.99	4.33±0.43	4.30±0.89
	Diff. (lt)	0.70±1.33	0.50±0.74	0.35±0.93	-0.11±1.01
	Diff. (%)	27.09±47.29	11.02±16.56	16.31±32.93	-1.70±22.43
MVV (lt/min)	Pre-test	112.98±34.45	129.43±15.46	112.05±30.52	142.88±15.48
	Post-test	135.58±23.73	160.32±31.50 <sup>a</sup>	125.13±16.17	135.93±18.23
	Diff. (lt)	22.60±23.79	30.88±22.46	13.08±27.60	-6.95±22.03
	Diff. (%)	25.76±29.18	23.58±17.66	17.64±34.72	-4.04±15.24
VC (lt)	Pre-test	4.48±2.02	3.97±1.13	2.39±1.13	4.67±1.06
	Post-test	6.09±0.77	5.26±0.42 <sup>a</sup>	3.79±1.17 <sup>a</sup>	4.62±1.06
	Diff. (lt)	1.61±1.64	1.29±1.00	1.40±0.84	-0.06±0.89
	Diff. (%)	69.99±93.49	41.37±39.83	74.29±66.19	-0.16±18.02
TV (lt)	Pre-test	0.60±0.19	0.83±0.31	0.64±0.14	0.69±0.40
	Post-test	0.88±0.51	1.26±0.46	0.75±0.35	1.08±0.52 <sup>a</sup>
	Diff. (lt)	0.28±0.42	0.43±0.68	0.11±0.32	0.40±0.30
	Diff. (%)	51.14±77.54	85.11±132.52	19.51±61.54	71.49±74.45
IVC (lt)	Pre-test	2.61±1.09	2.40±0.43	1.53±0.62	2.42±0.78
	Post-test	2.94±0.50	2.93±0.42	2.03±0.51	2.67±0.60
	Diff. (lt)	0.33±1.01	0.53±0.78	0.50±0.36	0.25±0.52
	Diff. (%)	28.29±50.49	26.69±31.57	39.42±31.91	15.28±29.91

a: significance between pre-post test; b: significant difference with IMT; c: Significant difference with IMT-SHAM; d: significant difference with control group FVC: forced vital capacity; FEV1: forced expiratory volume in 1 sec; MVV: maximal voluntary ventilation; VC: vital capacity; TV: tidal volume; IVC: inspiratory vital capacity; lt liters; sec seconds; min minutes. IMT-DF: intervention group, group that respiratory muscle training was performed using the diaphragm; IMT: intervention group, respiratory muscle training focused on chest breathing. IMT-SHAM: intervention group, a group that received without inspiratory muscle training. Control: control group, a group without inspiratory muscle training.

## DISCUSSION

The purpose of this study was to investigate the effect of conscious diaphragm muscle-based respiratory muscle exercise on respiratory muscle strength and respiratory function. For this purpose, respiratory muscle training was applied in different ways with healthy sedentary individuals, and respiratory muscle strength and functions were tested. In our study, significant differences were found in the respiratory muscle strength data in favor of the post-test. Especially in the MIP parameter, there was a significant difference in the IMT-DF group, IMT group, and control group while in the MEP parameter, a significant increase was achieved only in the IMT-DF group. One of the important findings that draws attention here is the significant changes that occurred in the control group. When the data of the control group was examined, it was determined that there was a significant difference in the respiratory muscle strength parameter (MIP) in favor of the post-test. Possible mechanisms for the significant difference

obtained here can be explained as follows; The reason for this is that the control group could not achieve a sufficient level of motivational test concentration during the pre-test. Thus, this may cause the pre-test data to be very low compared to the data obtained in the post-test. Another mechanism can be said that the individuals in the control group were physiologically more ready during the post-test measurement compared to the pre-test and felt better during the post-test. In terms of respiratory functions, the significant difference in TV (tidal volume) capacity in favor of the last test can be explained by this parallel. However, when the analysis of the change in the pre- and post-test was analyzed both as a percentage and as average values, no significant change occurred in the control group.

When the changes in respiratory muscle strength were examined in general, significant changes were realized in favor of the IMT-DF group in both the pre-posttest and the difference and percentage differences between the tests. In respiratory functions, along with the changes in

some parameters in IMT-SHAM and control groups, a significant change was observed in FEV1 in favor of the post-test in the IMT group. Many studies have been conducted to date on respiratory functions and respiratory muscle strength of strengthening exercises and breathing exercises for respiratory muscles (6, 15-17, 21, 22). However, it has been observed that the respiratory muscle exercises performed by consciously focusing on the diaphragm muscle were either not mentioned or it is thought that the studies were performed by ignoring that correct and effective breathing should be performed diaphragm centered (8, 11, 14, 15). There are studies in which significant changes in respiratory muscle strength and respiratory functions have been recorded, especially in the patient (eg. COPD) and disabled (eg. Down Syndrome) groups, where significant improvements have been recorded to alleviate the negative outcomes of disability and disease (1, 23). Although the studies show that significant and significant changes have been made in these parameters, it has been observed that no attention has been drawn to diaphragm-focused conscious breathing. In our study, diaphragm-focused inspiratory muscle training positively affected respiratory muscle strength and function. It is known that this positive and significant change is due to the increase in the contraction and relaxation capacity of the respiratory muscles, especially the diaphragm muscle (24). In addition, the significant change in respiration can be explained by physiological changes such as improvement in nervous conduction in respiratory muscles, increased contraction capacity, and increased metabolic enzyme activity required during contraction (25-27). Healthy skeletal muscles contribute to the control of many important functions during breathing. The reason why these contributions are so important is that the input needed for life is mostly provided by the airway and respiration (28). Muscle strength and functions will be positively affected as the performance increase provided consciously in respiratory muscles provides significant changes in the respiratory tract (29). When current studies are examined, it is shown that respiratory training or respiratory training significantly affects respiratory functions in patients, disabled, and healthy groups (17, 30-33). However, as mentioned before, it has not been reported that inspiratory muscle training was

performed consciously with a focusing the diaphragm. In our study, it is seen that conscious inspiratory muscle training focused on the diaphragm muscle provides significant changes in muscle strength and function. There are several limitations in this study. The first limitation of this study is that the sample does not consist of athletes. The second limitation is that the number of participants in the groups in the study is not sufficient. Therefore, in future studies, the application of the diaphragm-focused breathing exercise technique, especially in sports branches where respiratory muscle strength comes to the force (swimming, diving, underwater rugby, water polo), will play an important role in further detailing this technique and making its application widespread.

## CONCLUSION

In conclusion, inspiratory muscle training can significantly affect respiratory function and muscle strength, as in our study. The critical point to be emphasized in our study is that conscious respiratory muscle training focused on the diaphragm muscle can significantly affect respiratory muscle strength and function in healthy sedentary individuals.

## APPLICABLE REMARKS

- Although breathing plays a crucial part in sportive performance, there is currently not enough scientific information to paint a complete and accurate picture of how it works.
- The study's findings imply that diaphragm-specific respiratory muscle exercise affects healthy individual's ability to use their respiratory muscles and their strength.
- These findings are important in terms of supporting the development and popularizing the use of new training methods that can contribute to sports performance by integrating the diaphragm-focused technique into respiratory muscle training.

## AUTHORS' CONTRIBUTIONS

Study concept and design: Mehmet Vural, Mustafa Ozdal. Acquisition of data: M. Berk Demiryol, C. Didem Eyipinar. Analysis and interpretation of data: Mehmet Vural, Mustafa Ozdal. Drafting the manuscript: Mehmet Vural, M. Berk Demiryol, C. Didem Eyipinar. Critical revision of the manuscript for important intellectual content: Mustafa Ozdal. Statistical

analysis: Mehmet Vural, Mustafa Ozdal.  
Administrative, technical, and material support:  
Mustafa Ozdal. Study supervision: Mustafa Ozdal.

## CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

## REFERENCES

1. Borg M, Thastrup T, Larsen KL, Overgaard K, Hilberg O, Løkke A. Free diving-inspired breathing techniques for COPD patients: A pilot study. *Chron Respir Dis.* 2021;18: 14799731211038673. [doi:10.1177/14799731211038673] [PMid:34399604]
2. Burki, Nausherwan K. et al. "The ability of asthmatics to detect added resistive loads. *Am. Rev. Respir. Dis.* 2015;117 (1): 71-75.
3. Romer LM, McConnell AK. Specificity and reversibility of inspiratory muscle training. *Med Sci Sports Exerc.* 2003; 35(2), 237-244. [doi:10.1249/01.MSS.0000048642.58419.1E] [PMid:12569211]
4. Jin J, Sánchez-Sinencio E. A home sleep apnea screening device with time-domain signal processing and autonomous scoring capability. *IEEE Trans. Biomed. Circuits Syst.* 2014; 9(1), 96-104. [doi:10.1109/TBCAS.2014.2314301] [PMid:25486649]
5. Volianitis S, McConnell AK, Koutedakis Y, et al. Inspiratory muscle training improves rowing performance. *Med. Sci. Sports Exerc.* 2001; 33(5), 803–809. [doi:10.1097/00005768-200105000-00020] [PMid:11323552]
6. Kilding AE, Brown S, McConnell AK. Inspiratory muscle training improves 100 and 200 m swimming performance. *Eur. J. Appl. Physiol.* 2010; 108: 505-511. [doi:10.1007/s00421-009-1228-x] [PMid:19841931]
7. Romer LM, McConnell AK, Jones DA. Effects of inspiratory muscle training on time-trial performance in trained cyclists. *J. Sports Sci.* 2002; 20(7), 547-590. [doi:10.1080/026404102760000053] [PMid:12166881]
8. Lomax M, Grant I, Corbett J. Inspiratory muscle warm-up and inspiratory muscle training: separate and combined effects on intermittent running to exhaustion. *J. Sports Sci.* 2011; 29(6), 563-569. [doi:10.1080/02640414.2010.543911] [PMid:21347970]
9. Enright SJ, Unnithan VB. Effect of inspiratory muscle training intensities on pulmonary function and work capacity in people who are healthy: a randomized controlled trial. *Phys. Ther.* 2011; 91(6), 894-905. [doi:10.2522/ptj.20090413] [PMid:21493747]
10. Seixas MB, Almeida LB, Trevizan PF, et al. Effects of inspiratory muscle training in older adults. *Respir. Care.* 2020; 65(4), 535-544. [doi:10.4187/respcare.06945] [PMid:31662444]
11. Enright SJ, Unnithan VB, Heward C, et al. Effect of high-intensity inspiratory muscle training on lung volumes, diaphragm thickness, and exercise capacity in subjects who are healthy. *Phys. Ther.* 86(3), 2006; 345-354. [doi:10.1093/ptj/86.3.345] [PMid:16506871]
12. Liaw MY, Wang YH, Tsai YC, et al. Inspiratory muscle training in bronchiectasis patients: a prospective randomized controlled study. *Clin Rehabil.* 2011; 25:524-36. [doi:10.1177/0269215510391682] [PMid:21285287]
13. Archiza B, Andaku DK, Caruso FCR. Effects of inspiratory muscle training in professional women football players: a randomized sham-controlled trial. *J. Sports Sci.* 2018; 36(7), 771-780. [doi:10.1080/02640414.2017.1340659] [PMid:28622081]
14. Bostancı Ö, Mayda H, Yılmaz C, et al. Inspiratory muscle training improves pulmonary functions and respiratory muscle strength in healthy male smokers. *RESPNB.* 2019; 264, 28-32. [doi:10.1016/j.resp.2019.04.001] [PMid:30953791]
15. Lacasse Y, Goldstein R, Lasserson TJ, et al. Pulmonary rehabilitation for chronic obstructive pulmonary disease (review). *Cochrane Database Syst Rev.* 2006; 4(4). [doi:10.1002/14651858.CD003793.pub2]
16. Polkey MI, Green M, Moxham J. Measurement of respiratory muscle strength. *Thorax.* 1995; 50(11), 1131. [doi:10.1136/thx.50.11.1131] [PMid:8553266]
17. Klefbeck B, Nedjad JH. Effect of inspiratory muscle training in patients with multiple sclerosis. *Arch Phys Med Rehabil.* 2003; 84: 994-999. [doi:10.1016/S0003-9993(03)00133-3] [PMid:12881823]
18. Kim M, Park S, Lee M. The Comparison of Changes in Pulmonary Function and Respiratory Muscle Strength in Young Adults According to the Abdominal Breathing Exercise Methods. *Ann Appl Sport Sci.* 2023; 11(4). [doi:10.61186/aassjournal.1193]



19. Bostancı Ö, Özdal M, Hakan Mayda H et al. The effect of preparation period trainings on respiratory muscle strength of hearing impaired judokas. *Arch Budo Sci Mariia*. 2017; (13);97-102.
20. Vural M, Özdal M, Pancar Z. Effects of inspiratory muscle training on respiratory functions and respiratory muscle strength in Down syndrome: A preliminary study. *Isokinet. Exerc. Sci*. 2019; 27(4), 283-288. [doi:10.3233/IES-193127]
21. Miller MR, Hankinson J, Brusasco V, et al. ATS/ERS task force: standardisation of spirometry. *Eur Respir J*. 2005; 26: 319-338. [doi:10.1183/09031936.05.00034805] [PMid:16055882]
22. Pastuty, R. Effect of diaphragm breathing exercise on reducing anxiety level of pregnant women. *Journal of Maternal and Child Health Sciences (JMCHS)*. 2021; 1(2), 105-110. [doi:10.36086/jakia.v1i2.990]
23. Da Silva MR, Peruzzolo CC, Canizelli J, et al. Acute effects of breathing exercises on diaphragmatic mobility and respiratory muscles activity in healthy adults. *Cardiorespiratory Physiotherapy, Critical Care and Rehabilitation*. 2022; 1, 0-0. [doi:10.4322/2675-9977.cpcr.42701]
24. Wells GD, Pyley M, Thomas S, et al. Effects of concurrent inspiratory and expiratory muscle training on respiratory and exercise performance in swimmers. *Eur Journal Applied Physiology*. 2002; 200: 527-540. [doi:10.1007/s00421-005-1375-7] [PMid:15942767]
25. Ranatunga KW, Sharpe B, Turnbull B. Contractions of human skeletal muscle at different temperatures. *J Physiol*. 1987; 390(1): 383-395. [doi:10.1113/jphysiol.1987.sp016707] [PMid:3443940]
26. Proske U, Morgan DL, Gregory JE. Thixotropy in skeletal muscle and in muscle spindles: a review. *Prog Neurobiol*. 1993; 41(6): 705-721. [doi:10.1016/0301-0082(93)90032-N] [PMid:8140258]
27. Karvonen J, Lemon PWR. *Medicine in Sports Training and Coaching*. Karger Pub. 1992; 190-213.
28. Mehrotra PK, Varma N, Tiwari S, et al. Pulmonary functions in Indian sportsmen playing different sports. *IJPP*. 1998; 42: 412-6.
29. Joshi LN, Joshi VD, Gokhale LV. Effect of short term pranayam, practice of breathing rate and ventilatory functions of lung. *Indian J Physiol Phamscol*. 1992; 36(2): 105-108.
30. Laoutaris I, Dritsas A, Brown MD, et al. Inspiratory muscle training using an incremental endurance test alleviates dyspnea and improves functional status in patients with chronic heart failure. *Eur J Cardiovasc Prev Rehabil*. 2004; 11(6): 489-496. [doi:10.1097/00149831-200412000-00008] [PMid:15580060]
31. Hulzebos EH, Helder PJ, Favié NJ, et al. Preoperative intensive inspiratory muscle training to prevent postoperative pulmonary complications in high-risk patients undergoing CABG surgery: a randomized clinical trial. *JAMA*. 2006; 296(15): 1851-7. [doi:10.1001/jama.296.15.1851] [PMid:17047215]
32. Carmichael C, Sapienza Ramig L. Expiratory muscle strength training and the Lee Silverman treatment program, in Parkinson's disease. *Conference of Speech Motor Control*. March, Austin, Texas. 2006.
33. Özdal, M. Acute effects of inspiratory muscle warm-up on pulmonary function in healthy subjects. *Respir Physiol Neurobiol*. 2016; 227, 23-26. [doi:10.1016/j.resp.2016.02.006] [PMid:26903486]